

# Open Research Online

---

The Open University's repository of research publications and other research outputs

## Correlations between SO<sub>2</sub> flux, seismicity, and outgassing activity at the open vent of Villarrica volcano, Chile

### Journal Item

How to cite:

Palma, José Luis; Calder, Eliza S.; Basualto, Daniel; Blake, Stephen and Rothery, David A. (2008). Correlations between SO<sub>2</sub> flux, seismicity, and outgassing activity at the open vent of Villarrica volcano, Chile. *Journal of Geophysical Research: Solid Earth*, 113(B10) B10201.

For guidance on citations see [FAQs](#).

© [\[not recorded\]](#)

Version: [\[not recorded\]](#)

Link(s) to article on publisher's website:  
<http://dx.doi.org/doi:10.1029/2008JB005577>

---

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's [data policy](#) on reuse of materials please consult the policies page.

---

1 **Correlations between SO<sub>2</sub> flux, seismicity and**  
2 **outgassing activity at the open vent of Villarrica**  
3 **volcano, Chile**

José Luis Palma<sup>1,2</sup>, Eliza S. Calder<sup>3</sup>, Daniel Basualto<sup>4</sup>, Stephen Blake<sup>1</sup>, and  
David A. Rothery<sup>1</sup>

---

J. L. Palma, Department of Earth Sciences, The Open University, Walton Hall, Milton Keynes,  
MK7 6AA, U.K. (j.l.palma@open.ac.uk)

<sup>1</sup>Department of Earth Sciences, The Open  
University, Milton Keynes, U.K.

<sup>2</sup>Now at Department of Geological and  
Mining Engineering and Sciences, Michigan  
Technological University, Houghton,  
Michigan, U.S.

<sup>3</sup>Department of Geology, State University  
of New York at Buffalo, Buffalo, New York,  
U.S.

<sup>4</sup>Southern Andes Volcano Observatory,  
Servicio Nacional de Geología y Minería,  
Temuco, Chile.

**Abstract.**

The characteristics of the open-vent activity of Villarrica volcano, Chile, were studied in detail by integrating visual observations of the lava lake, analysis of the seismic tremor and measurements of SO<sub>2</sub> flux. The outgassing activity comprises a persistent gas plume emission from the bottom of the crater as well as frequent explosive events. Three main styles of bubble bursting were identified at the surface of the active lava lake: seething magma, small short-lived lava fountains and strombolian explosions. Seething magma consists of continual burst of relatively small bubbles (a few meters in diameter) with varying strength over the entire surface of the lava lake. Small lava fountains, seen as a vigorous extension of seething magma, commonly have durations of 20-120 s and reach 10-40 m high above the lava lake. Correlations between seismicity and visual observations indicate that the seismic tremor is mostly caused by the explosive outgassing activity. Furthermore, for different periods between 2000-2006, during which the activity remained comparable, the RSAM and SO<sub>2</sub> emission rates show a very good correlation. Higher SO<sub>2</sub> emissions appeared to be related to higher levels of the lava lake, stronger bubble bursting activity and changes in the morphology and texture of the crater floor. Background (low) levels of activity correspond to a lava lake located >80 m below the crater rim, small and/or blocky morphology of the roof, seismic amplitude (RSAM) lower than 25 units, few volcanotectonic earthquakes, and daily averages of SO<sub>2</sub> emissions lower than 600 Mg/day.

## 1. Introduction

26 Villarrica volcano (39.42 S, 71.93 W, 2847 m a.s.l) is the most active volcano in the  
27 southern Andes. Like other open-vent volcanic systems (e.g. Stromboli, Ripepe [1996],  
28 Bertagnini *et al.* [2003]; Mount Erebus, Rowe *et al.* [2000], Aster *et al.* [2003]; and Masaya,  
29 Duffell *et al.* [2003], Williams-Jones *et al.* [2003]), Villarrica is characterized by persistent  
30 degassing and sustained seismicity [Calder *et al.*, 2004]. Since the last eruption in 1984-  
31 1985, it has shown persistent gas plume emission and bubble-burst activity at the surface of  
32 an active lava lake typically located less than 200 meters below the crater rim [Fuentealba  
33 *et al.*, 2000; Calder *et al.*, 2004]. Open-vent volcanoes of low silica composition (e.g.  
34 basalts, phonolite) commonly exhibit activity that ranges from sluggish and slowly moving  
35 lava lakes, as seen at Erta Ale's summit caldera [Oppenheimer and Yirgu, 2002; Harris  
36 *et al.*, 2005], to intermittent explosive strombolian eruptions from multiple vents, as seen  
37 at Stromboli volcano [Ripepe, 1996; Ripepe *et al.*, 2005; Patrick *et al.*, 2007]. In the  
38 former, the formation of a cooled lava crust on top of a convecting degassed magma is  
39 a common occurrence, with sporadic bubble-bursting also taking place. More explosive  
40 activity involves gas slugs rising through a magma-filled conduit and rupturing at the  
41 magma free-surface [Ripepe *et al.*, 2002; Aster *et al.*, 2003]. Although not sustained for  
42 long periods of time, other types of activity such as lava fountains (e.g. Vergnolle [1996];  
43 Andronico *et al.* [2005]) and abnormally strong (paroxysmal) explosions (e.g. Calvari  
44 *et al.* [2006]) can occur during eruptions or periods of elevated activity. Despite the  
45 importance of understanding the characteristics of these eruption styles and the evolution

46 of the activity with time (e.g. Andronico *et al.* [2005]; Harris and Ripepe [2007a]), at  
47 Villarrica these aspects have not been investigated in detail.

48 In this study the main characteristics and variations of the continuous strombolian  
49 activity of Villarrica volcano have been identified, and the relationship between seismicity,  
50 degassing and observed activity at the summit has been established. This was achieved  
51 through new measurements of SO<sub>2</sub> flux, seismic data and visual observations pertaining to  
52 the period November 2004 to February 2006. Total SO<sub>2</sub> fluxes measured at the summit and  
53 around the volcano were correlated with volcanic tremor. Video recordings, photographs,  
54 and direct observations of the lava lake activity within the crater allowed us to identify  
55 different styles of bubble bursting, and to describe some distinct activity which we consider  
56 unique to this particular volcano with its visible active lava lake.

## 2. Background

57 Villarrica is an active stratovolcano located in the southern Andes of Chile (33°- 46° S)  
58 (Figure 1). The snow-covered cone is located on the northwest side of an 6.5 x 4.2 km  
59 elliptical caldera that was created during the Late Pleistocene (ca. 95 ka, Moreno *et al.*  
60 [1994]; Clavero and Moreno [2004]). The interior of the crater has a funnel shape and steep  
61 inner walls, with an internal diameter of approximately 150±10 meters at the crater rim  
62 (Figure 2). The predominant composition of lavas and pyroclastic deposits is basaltic to  
63 basaltic andesite (50-57 wt% SiO<sub>2</sub>) [Moreno *et al.*, 1994; Witter *et al.*, 2004; Hickey-Vargas  
64 *et al.*, 2004]. The tectonic setting has been described in López-Escobar *et al.* [1995]; Lavenu  
65 and Cembrano [1999]; Ortiz *et al.* [2003], the geology and composition of the products in  
66 Moreno *et al.* [1994]; Witter *et al.* [2004]; Clavero and Moreno [2004]; Hickey-Vargas *et al.*  
67 [2004], gas plume composition in Witter *et al.* [2004]; Witter and Calder [2004]; Shinohara

68 and Witter [2005], and the recent eruptive activity in Fuentelba *et al.* [2000]; Ortiz *et al.*  
69 [2003] and Calder *et al.* [2004].

70 Volcanic tremor is one type of seismic signal that characteristically accompanies the ac-  
71 tivity of open-vent volcanoes. It may last for minutes, days, or even months [McNutt, 2000;  
72 Zobin, 2003; Ripepe, 1996; Falsaperla *et al.*, 2005]. Several studies have shown a correla-  
73 tion between tremor amplitude and volcanic activity (e.g. Mount Etna, Falsaperla *et al.*  
74 [2005]; Kilauea, Koyanagi *et al.* [1987]). For instance, the continuous tremor recorded on  
75 Etna during the eruption of July-August 2001 showed amplitude and frequency variations  
76 that have been linked to the level and style of activity: phreatomagmatic explosions, lava  
77 fountains, lava effusion and strombolian explosions [Falsaperla *et al.*, 2005]. Since the start  
78 of the seismic monitoring of Villarrica volcano in 1982, permanent tremor and variable  
79 amounts of long-period events have been the dominant seismicity recorded [Fuentelba  
80 and Peña, 1998]. Other types of signals present in Villarrica seismicity have been classi-  
81 fied as explosion events, hybrid signals and volcano-tectonic earthquakes [Fuentelba and  
82 Peña, 1998; Calder *et al.*, 2004]. The volcano observatory of the southern Andes (OV-  
83 DAS), of the Chilean geological survey (Servicio Nacional de Geología y Minería), uses a  
84 Real-time Seismic-Amplitude Measurement system (RSAM, Endo and Murray, 1991) as  
85 one of the main tools for tracking changes in the volcanic activity at Villarrica.

86 Measurements of SO<sub>2</sub> concentrations using correlation spectroscopy have proved to be  
87 a valuable tool in volcano monitoring, and in the investigation of the dynamics of magma  
88 degassing [Stoiber *et al.*, 1983; Fischer *et al.*, 2002; Young *et al.*, 2003]. At Villarrica,  
89 there are only a few periods with SO<sub>2</sub> flux measurements. They were mostly carried out  
90 by Witter *et al.* [2004] during early 2000 and 2001 to constrain the total gas emission and

91 analyze the dynamics of degassing within the system. Other measurements have been  
92 also carried out by OVDAS as part of their monitoring program. Daily averages of SO<sub>2</sub>  
93 flux typically revealed moderate emissions ca. 100-700 Mg/day. Combining SO<sub>2</sub> emission  
94 rates with measurements of sulphur and other gas species within the gas plume (as molar  
95 ratios) and glass inclusions, Witter *et al.* [2004] and Shinohara and Witter [2005] showed  
96 that: i) magma ascending to the surface is almost completely degassed, ii) the relative  
97 abundance of gas species in the gas plume remains constant during spattering of lava at  
98 the vent, which implies that bubbles bursting at the lava lake are in equilibrium with the  
99 magma, and that iii) on average, ca. 2.2 m<sup>3</sup>/s of magma is degassed. They concluded that  
100 convection within the conduit is the most appropriate model of degassing for Villarrica  
101 volcano.

102 In this paper we make a distinction between the terms magma degassing and outgassing.  
103 We recognize three main processes responsible for the transfer of magmatic gas from deep  
104 magma chambers to the atmosphere: exsolution of gas from the melt, gas segregation and  
105 outgassing. Gas exsolution involves bubble nucleation and bubble growth by diffusion of  
106 gas into it [Sparks, 2003]. Once the gas has been separated from the melt and formed bub-  
107 bles of sufficiently large size, it migrates upwards through a magma-filled plumbing system  
108 in the process that is commonly termed ‘gas segregation’. Continuous bubble growth by  
109 decompression, gas diffusion, and coalescence occurs during this stage. ‘Degassing’ is a  
110 general term often implying any or all of these processes (e.g. Sparks [2003]). The term  
111 ‘outgassing’ has been previously used by Gerlach [1986]; Ryan [1995]; Adams *et al.* [2006];  
112 Lautze and Houghton [2007], among others, to describe the escape of gas from magma.  
113 However, despite a consensus on the main idea, a clear definition has not been provided.

114 Accordingly, magma outgassing is here defined as the escape of gas from the magma ei-  
115 ther directly to the atmosphere, or to the permeable country rock or hydrothermal system  
116 surrounding a magma body. Gas may escape as a segregate gas phase or by diffusion of  
117 gas from the edge of the magma body. For instance, gas may escape to the atmosphere  
118 by gas diffusion at the surface of a magma column. Hence, outgassing activity includes  
119 all the processes of bubble bursting, non-explosive gas emission, and sustained explosion  
120 by which gas escapes to the atmosphere.

### 3. Recent activity of Villarrica volcano

#### 3.1. Background activity

121 Since the eruption of 1984-1985, continuous shallow magmatic activity has been seen  
122 inside the crater at the summit of the volcano. These observations account for persistent  
123 outgassing that involve mild strombolian activity taking place at the surface of a highly  
124 dynamic lava lake, which represents the top of the magma column within the main conduit.  
125 Inside the main crater, the lava lake is commonly located at depths ranging from >150 to  
126 about 50 m below the summit. Because of the changing morphology of the crater floor,  
127 the lava lake is not always visible from the crater rim (Figure 2). Descriptions of the  
128 activity at the lava lake by Fuentealba *et al.* [2000] and Calder *et al.* [2004] mention the  
129 rapid crusting-over of a relatively tranquil lava surface as well as vigorous ~5-30 m high  
130 fountaining.

131 Within this scenario of volcanic activity, background (low to moderate) levels of activity  
132 are commonly characterized by the surface of the lava lake being located more than  
133 90 meters below the crater rim. Gentle roll-over with brief moments of quiescence of the  
134 lava lake (as observed by Calder *et al.* [2004]) is likely to mark the lowest strength in

135 activity of the visible lava lake. Strombolian explosions rarely reach 100 meters above the  
136 lava free surface. In addition, during low levels of activity the tremor amplitude rarely  
137 reaches values above 20 RSAM units (persisting for more than one day), and the seismicity  
138 lacks considerable amounts of volcano tectonic-type earthquakes.

139 Although these characteristics constrain the predominant volcanic activity observed  
140 since 1985, Villarrica volcano displays continuous variations in seismicity, activity observed  
141 at the crater, and amount and style of outgassing.

### 3.2. The 1999 and 2000 crisis

142 Three different episodes of high activity have been observed recently: in 1999, 2000 and  
143 2005. These periods developed with abnormal types of observed and seismic activity but  
144 did not culminate in eruptions. The first two are documented by Calder *et al.* [2004] and  
145 Ortiz *et al.* [2003], respectively, whereas the activity exhibited in 2005 is described in this  
146 paper (Section 5).

147 Between August and December 1999, Villarrica showed a significant increase in seismic  
148 activity, a rise in the level of the lava lake, several large but discrete explosions, and  
149 morphological changes of the crater floor [Calder *et al.*, 2004]. Several episodes of sudden  
150 increase in seismic amplitude (two or three-fold) lasting for several hours occurred in  
151 August, November and December. As a result of the explosive activity, scoria bombs up  
152 to 50 cm in diameter were found on the crater rim and tephra fall deposits extended up  
153 to 5 km from the vent [Calder *et al.*, 2004].

154 During September 2000, several tectonic earthquakes occurred in the region, including  
155 a magnitude 3.8 earthquake that took place less than 70 km from the volcano on 20  
156 September. Subsequently, the tremor spectrum exhibited a frequency shift of its dominant

157 peak from 1 to 2 Hz [Ortiz *et al.*, 2003]. This seismic activity was associated with a sudden  
158 decrease in the fumarolic activity observed at the crater, and an apparent crusting-over  
159 of the lava lake which concealed the nature of magmatic activity. Reestablishment of  
160 the activity at the crater was accompanied by pahoehoe flows on the crater floor and  
161 the construction of a spatter cone (end of October–beginning of November). During  
162 October and November 2000 the seismicity showed peaks at higher frequencies (up to 5  
163 Hz), returning to normal at the beginning of 2001.

### 3.3. Morphology of the crater floor: the spatter roof

164 Cooling at the surface of the lava lake creates partly solidified patches of crust that  
165 stay afloat temporarily on the lava surface. At Villarrica, slowly moving crust plates,  
166 as seen at Erta Ale (e.g. Oppenheimer and Yirgu [2002]; Harris *et al.* [2005]), are not  
167 observed. Instead, continually ejected spatter adheres to the inner walls of the vent and  
168 forms a spatter roof that grows by accretion and agglutination of the pyroclastic material  
169 (Figures 2, 3a). The spatter roof can partly or completely conceal the activity of the lava  
170 lake. The location of this roof also effectively defines the depth of the observable crater  
171 floor. The roof is commonly unstable and experiences frequent collapses depending on the  
172 intensity of the lava lake activity underneath; when the explosive activity increases, the  
173 roof thickens by accumulation of material on its upper surface, eventually creating ter-  
174 races (relatively flat surfaces, Figure 3b), or even forming a small scoria cone (Figure 3c).  
175 The collapsing of the roof or terraces may occur mainly due to the increasing overburden,  
176 but also by thermal erosion of its lower surface by the hot magma. Terraces can dis-  
177 play concentric fractures which subsequently accommodate collapses. In turn, the outer  
178 concentric fracture observed on 16 January (Figure 3b) was the site of a small collapse

179 that occurred within two days after the photograph was taken (16-18 January 2005). On  
180 26 January 2005, another small collapse was witnessed by mountain guides. This time,  
181 pyroclastic material that had accumulated around an elongated aperture in the spatter  
182 roof, not more than 10 m wide, collapsed and left an almost circular opening about 25 m  
183 in diameter. When the magma column withdraws and the free surface height lowers, the  
184 unstable roof usually collapses often within a time span of hours.

## 4. Observed outgassing styles

### 4.1. Continuous outgassing

185 As mentioned earlier, Villarrica volcano is characterized by the continuous emission of  
186 a gas plume from the summit. This ‘passive’ gas release is the background outgassing  
187 activity observed at the crater. It has persisted since the end of the last eruption in 1985,  
188 although a gas plume has been observed recurrently since the end of the XIX century  
189 [Casertano, 1963].

190 Direct visual observations from the crater rim along with measurements of SO<sub>2</sub> path-  
191 length concentrations evidence variations in the emissions from the bottom of the crater.  
192 Rough estimations indicate that fluctuations in the gas flux exceed 50% of total emissions.  
193 Several factors can contribute to these variations, such as the accumulation of gas inside  
194 the lava lake-spatter roof cavity, and the sudden release of higher amounts of gas through  
195 explosive activity (in which the gas is vented at higher speeds). Thus, small gas puffs  
196 can be seen rising with irregular periodicity from the crater floor. Wind entering the  
197 crater intensifies the gas circulation and contributes to further variations in the gas fluxes  
198 observed outside the crater. For example, Bluth *et al.* [2007] showed measurements of SO<sub>2</sub>  
199 fluxes of the gas plume of Villarrica volcano, up to 3500 m away from the crater, with

200 fluctuations that ranged over a factor of three (between 197 and 640 Mg/d; see Figure 2  
201 in Bluth *et al.* [2007]).

## 4.2. Bubble burst activity

202 Mild strombolian explosions are, perhaps, the most common explosive activity observed  
203 at the crater. However, five distinct types of bubble burst have been observed during  
204 periods with different levels of activity (see color photos and videos in auxiliary material).  
205 Here, the term ‘bubble bursting’ refers to the processes involved in gas-bubble rupture at  
206 the magma free-surface, fragmentation and subsequent ejection of pyroclastic material,  
207 and includes:

208 1. Seething magma: This distinctive bubble bursting style is distinguishable only when  
209 the lava free-surface is visible. The activity at the surface of the lava lake resembles  
210 the dynamics of boiling water, as medium-size bubbles (~0.5-2 m in diameter) burst  
211 continuously across the magma free surface (Figure 4). Bubbles can be seen to rupture  
212 at the same location within the space of a few seconds. This style of activity induces the  
213 magma surface to experience continuous wave-like undulations. The vigour of seething  
214 magma is variable: on the lower end only a few bubble bursts occur per minute and gentle  
215 roll-over of the magma surface can be easily identified; at the more vigorous end of the  
216 spectrum, there are single bubble bursts almost every second ejecting pyroclasts more  
217 than 5 m above the lava lake.

218 2. Small lava fountains: Fountains of lava occur through a relatively wide roof opening.  
219 Compared to the more classic hawaiian style of activity observed at Kilauea volcano  
220 (e.g. Parfitt [2004]), in which fountains reach tens to a few hundred meters high, these  
221 lava fountains are very small and sustained only briefly (Figure 5). They normally last

222 between 20 seconds and 2 minutes, and reach 10 to 40 meters high. To some extent,  
223 a small lava fountain resembles a very strong variety of seething magma, with a much  
224 greater concentration of bubbles continuously reaching the surface. Although difficult to  
225 quantify, it is apparent that most of the pyroclastic material is centimeter-to-decimeter-  
226 sized clots of magma. Small lava fountains are less common than both seething magma  
227 and strombolian explosions.

228 3. Strombolian explosions: This type of bubble bursting, named after the activity at  
229 Stromboli volcano, has been described as mild explosions that occur from the rise and  
230 rupture of large gas bubbles or slugs at the top of the magma column [Vergnolle and  
231 Mangan, 2000; Parfitt, 2004]. The magnitude and frequency of strombolian explosions at  
232 Villarrica volcano are both variable. Strong explosions ejecting pyroclastic material over  
233 100 m above the vent are mostly seen when the general level of both observed and seismic  
234 activity is high, whereas during periods of reduced activity, when the level of magma is  
235 low within the crater, it is very rare to see such explosions. The duration of an explosion  
236 ranges from a fraction of a second, involving a single strong burst, to more than 15 seconds  
237 when the explosion is composed of rapid sequences of pyroclastic ejections. Due to the  
238 morphology of the vent, some explosions do not exhibit pyroclastic ejection through the  
239 orifice in the spatter roof, but a sudden and relatively rapid gas emanation that ascends  
240 as a distinctive thermal plume. Sometimes, when no spatter is emitted, the spurt of gas is  
241 the only evidence of explosions or a strong bubble burst. Strombolian explosions observed  
242 at Villarrica are similar to Stromboli type 1 eruptions described by Patrick *et al.* [2007],  
243 essentially by virtue of the dominance of coarse particles and lack of a dense ash plume.  
244 There have been observations of atypical explosions at Villarrica that are accompanied

245 by a brownish ash plume. They are associated with partial collapses of the spatter roof  
246 or avalanches from the inner crater walls, which is concurrent with the idea of backfilling  
247 of loose material for the type 2 eruptions at Stromboli [Patrick *et al.*, 2007].

248 4. Gas jetting: Gas jets are strong exhalations of gas and relatively fine (ash-lapilli  
249 size) pyroclastic material. Their durations are normally longer than that of strombolian  
250 explosions. None of them have been observed directly generated from the lava lake surface,  
251 but instead they originate through an opening in the spatter roof or spatter cone. If the  
252 hole in the roof is small, relative to the surface area of the lava lake underneath, it is likely  
253 that during an explosion or small lava fountain only the fine fraction of the ejecta vents  
254 through the hole, generating the impression of an exhalation of gas with only a small  
255 amount of coarse material. There have been observations of near simultaneous explosions  
256 and gas jetting events at two adjacent openings (less than 20 meters apart). In that case,  
257 the gas jet occurs in the smaller hole (in effect, a blow-hole), and commonly lasts as long  
258 as the explosion. Gas jet-like activity has also been observed during periods of elevated  
259 activity, when a scoria cone has built up on top of the spatter roof, and whose opening is  
260 generally narrow. In these instances, the gas jet resembles an energetic narrow fountain  
261 whose spatter, ash to bomb in size, can reach a hundred or more meters in height.

262 5. Splashing lava: A fifth type of activity, indirectly related to bubble bursting, has been  
263 observed occurring particularly when the spatter roof covers a big part of the lava lake.  
264 In this case, the roof prevents the explosions from sending pyroclastic material out of the  
265 vent. Often, shortly after an explosion is heard or a gas spurt is observed, a considerable  
266 amount of spatter is expelled through the roof orifice. It is characteristically coarse spatter  
267 that fragments on exit and accumulates around the vent. We believe that this material is

268 not derived from primary fragmentation of magma during bubble rupturing, but is caused  
269 by subsequent splashing of lava associated with waves generated on the lava free-surface  
270 in the aftermath of an explosion.

271 These descriptions recognize different mechanisms for the explosive events. Whilst  
272 seething magma, strombolian explosions and small lava fountains represent types of pri-  
273 mary bubble bursting activity at the surface of the lava lake, gas jetting appears to result  
274 from a combination of bubble burst activity and subsequent interaction with the spatter  
275 roof. This interaction modifies the development of the bursting activity and ejection of  
276 pyroclastic material as it leaves the vent. This would explain why the size distribution  
277 of pyroclasts ejected during gas jetting appears skewed towards smaller fractions com-  
278 pared to that generated as a result of the primary fragmentation. Spatter generated  
279 from splashing magma activity is derived directly from the lava lake, although again the  
280 fragmentation is not caused by bubble bursting directly.

## 5. The activity during November 2004-April 2005

### 5.1. Chronology

281 From the end of 2004 and until June 2005, the volcano showed an increase in activity as  
282 recorded by seismicity as well as visual observations. A summary of the chronology of the  
283 principal events and dates is given in Table 1. The rise in activity levels was accompanied  
284 by frequent changes in the morphology of the crater floor (Figure 3). In November 2004  
285 the bottom of the crater was >90 m below the crater rim, from where the lava lake was  
286 out of direct view. Spatter from small strombolian explosions could be seen but rarely  
287 reached more than 20 m above the spatter roof. By the end of November, a new small  
288 spatter roof had formed at the bottom of the crater (Figure 3a). Subsequently, the activity

289 increased gradually and, by the middle of December, some explosions were sending bombs  
290 and spatter up to 100 m above the crater floor. Increasing amounts of new tephra were  
291 observed around the vent and also on the crater rim.

292 During January 2005 the intensity of the strombolian activity continued to increase,  
293 accumulating abundant material on top of the spatter roof and crater walls. Although it  
294 was unusual to see large explosions during the time spent at the summit of the volcano  
295 (usually between 1.5-3.5 hours), by the middle of January it was more common to find new  
296 pyroclastic material (lapilli-bomb sized) on the north-east side of the crater rim. During  
297 this period, the morphology of the crater floor was evolving rapidly and showed evidence  
298 of repeated construction and partial collapse cycles. A prominent upper terrace formed  
299 only ~50 meters below the crater rim, suggesting the rise of the magma free surface. The  
300 second half of January was characterized by a rapid increase in the level of activity and,  
301 by the end of month, explosions were ejecting centimeter-sized pyroclasts up to ca. 100  
302 meters above the bottom of the crater (~50 m above the crater rim).

303 There were no further substantial changes until late February and early March when  
304 the upper terrace increased in thickness by accumulation of pyroclastic material. By the  
305 end of March, a small spatter cone started to grow on top of the roof with an orifice less  
306 than 10 m in diameter on its top (Figure 3c). During April the roof morphology kept  
307 changing, notably the size of the cone and diameter of its vent. The small vent acted  
308 as a nozzle generating narrow, vertically-directed jet-like explosions. The most vigorous  
309 explosions observed from the crater rim had durations of the order of 3-10 seconds and  
310 ejected bomb-sized pyroclasts at least 150 m above the crater floor. This explosive activity  
311 continued through May and explosions throwing material above the crater rim were seen

312 until July. By the end of July–early August the lava roof (and magma free–surface) had  
313 retracted to about 70 m below the crater rim, which was accompanied by a decline of the  
314 activity, marking the end of the 2004-2005 episode.

315 Pyroclasts derived from explosions exhibit vesicles with a broad size distribution. Two  
316 examples of typical pyroclasts found on the crater rim of Villarrica volcano are shown  
317 in (Figure 6). Reticulite is common among this material (Figure 6a). General textural  
318 characteristics of scoria found on the crater rim include high vesicularity (>60% in the  
319 most vesicular samples) with vesicle radii up to a few centimeters (Figure 6b), irregular and  
320 iridescent surface with adhered Pele’s hair, or spatter with cowpat-like form (elongated  
321 and flat) in some products of big explosions. Often after an explosion, it is only the  
322 more dense material that remains on the crater rim as the highly vesicular ejecta is easily  
323 dispersed by the wind. Observations of millimeter to centimeter-sized vesicles in scoria  
324 samples, along with observations of the meter-sized bubbles bursting at the lava lake,  
325 are an indication of a very broad bubble-size distribution in the gas phase reaching the  
326 magma free–surface.

## 5.2. Characteristics of the seismic tremor

327 During November 2004-April 2005, two short-period vertical component seismic sta-  
328 tions were operating near the volcano, at 3.7 km (station VNVI) and 19 km (station  
329 CVVI) to the NW and W of the crater, respectively (Figure 1). Both are part of the  
330 volcano monitoring seismic array operated by the Southern Andes Volcano Observatory  
331 (OVDAS-SERNAGEOMIN). In this study, we utilize data from the VNVI station, which  
332 is the closest station to the volcano. We do not correct for the frequency response of the  
333 seismometer, which is flat above the corner frequency of 1 Hz. This limits the quantitative

334 analysis of frequency peak amplitude and dominance below 1 Hz, but it does not affect  
335 the temporal analysis of amplitude and frequency variations.

336 Tremor at Villarrica contributes more than 90% of the total seismic energy. It is com-  
337 monly a continuous, irregular and low amplitude seismic signal. Its waveform generally  
338 has a pulsating pattern in which short tremor bursts of higher amplitude occur as often as  
339 once per minute (Figure 7). The frequency of occurrence of the higher amplitude bursts  
340 varies with the level of activity (Calder *et al.* [2004] reported 10 events per hour in 1999).  
341 Periods of elevated observed volcanic activity have an overall higher occurrence of these  
342 events. If viewed as individual discrete events, these higher amplitude bursts normally  
343 last less than 50 seconds, are generally characterized by emergent starts and ends (gradual  
344 increase and decrease in amplitude, respectively), and show a wide range of amplitudes  
345 and durations (Figure 7). They have been previously described as low-frequency events  
346 generated by strombolian explosions [Fuentelba and Peña, 1998; Fuentelba *et al.*, 2000].

347 Banded tremor, as described in the literature of Kilauea volcano [Koyanagi *et al.*, 1987],  
348 is also characteristic of the seismicity of Villarrica volcano. Such higher amplitude tremor  
349 can last minutes to days, but commonly has a duration of a few hours. Typically, it  
350 preserves the spectral features of the lower amplitude tremor preceding it. Banded tremor  
351 repeats during the period of study and has been described during the 1999 crisis [Calder  
352 *et al.*, 2004].

353 One of the main tools employed by the Southern Andes Chilean Volcano Observatory  
354 (OVDAS), to routinely monitor the activity of Villarrica volcano, is the amplitude of the  
355 seismic signal, which is measured as RSAM units [Endo and Murray, 1991]. Changes and  
356 trends in seismic amplitude (RSAM) for the period November 2004-April 2005 correlate

357 well with the volcanic activity described above (Table 1 and Figure 8a). RSAM started  
358 increasing in December and continued increasing until the end of January, reaching values  
359 of between 40 and 50 RSAM counts. There was a decrease in amplitude at the beginning  
360 of February, although remaining higher than 20 counts, and a further sudden decrease that  
361 is followed by a slow increase starting the last week of February and reaching values of 30  
362 counts by the middle of March. During the second half of March and April, the RSAM  
363 values remained fairly constant between 20 and 30 counts. In general, these variations are  
364 manifested by the total range and maximum values of the RSAM amplitude (Figure 8a).  
365 In addition to the long monthly trend in seismic amplitude there are some short episodes,  
366 normally just a few days, where the overall RSAM amplitude changes abruptly to higher  
367 or lower values (e.g. 14-19 and 29-30 January 2005). One of these episodes, at the middle  
368 of January, was correlated with increasing gas emissions (Section 6.2).

369 Throughout the analyzed period most of the seismic energy is concentrated within the  
370 frequency range 1-7.2 Hz, with the highest peaks commonly between 1-2.15 Hz (Figures 8b,  
371 9). Along with the increasing activity from November 2004 to January 2005, the associated  
372 tremor displayed gradual variations in the relative amplitude of the dominant peaks in the  
373 frequency domain: the amplitude in the frequency band 1.65-2.15 Hz increases whereas in  
374 the frequencies bands 1-1.35 and 2.55-7.2 Hz the relative amplitude decreases (Figures 8b,  
375 9). Sharp changes in amplitude within the range 2.55-5.5 Hz, occurring on 14-19 and 29-30  
376 January 2005, strongly contribute to the sudden increases in RSAM values (Figures 8, 9,  
377 10). These variations last for a few hours or days and do not represent individual transient  
378 events. The relationship between these short periods of higher amplitude and the activity  
379 observed at the summit of the volcano is not clear. Although environmental effects such

380 as high winds cannot be ruled out, weather reports and observations in the field did not  
381 point to any particular meteorological conditions that would coincide with the timing of  
382 these variations. Moreover, a closer look at the signal reveals different amplitudes and  
383 start times of these peaks at low and high frequencies (Figure 10). The onset of the tremor  
384 with highest amplitudes in the band 2.15-7.2 Hz coincides with a decrease in the amplitude  
385 of the band 1-2.15 Hz. Only a few hours later, when the amplitude of the relatively high  
386 frequency band (2.15-7.2 Hz) is in a waning stage, the low frequency band (1-2.15 Hz)  
387 recovers its previous amplitude (Figure 10). Unfortunately, detailed visual observations  
388 of the crater on some of these days (16 and 18 January 2005) were not sufficient to allow  
389 the correlation of these variations with the volcanic activity.

390 During periods of elevated seismic activity, such as in December 2004 and January 2005,  
391 the daily RSAM exhibited a broad range of values with a fast increase and subsequent  
392 slow fall in amplitude (Figure 11a). These fluctuations had a periodicity of about 1.9 to 5  
393 hours, but the most prominent saw-tooth cycles commonly had a duration between 2 and  
394 3.5 hours. Seismic traces coincident with high RSAM units evidence a higher frequency  
395 and amplitude of the tremor bursts (Figure 11b), although the frequency content of the  
396 tremor is similar on both RSAM peaks and RSAM troughs (Figure 11c). During periods  
397 where the RSAM was low ( $< 20$  units), these features were absent or appeared more  
398 erratic and less frequent.

### 5.3. Correlation between outgassing activity and tremor magnitude

399 Simple experiments were carried out between January-February 2006 in order to cor-  
400 relate the seismic signal with the observed activity at the summit of Villarrica volcano.  
401 From the crater rim, we made timed observations of the explosions and other outgassing

402 related events (Figure 12). During this period the level of activity was considered low,  
403 with the bottom of the crater located more than 80 m below the crater rim. The di-  
404 ameter of the orifice in the roof was no greater than 15 m and so the lava lake was not  
405 directly visible. A second small hole was present on the west side of the roof and showed  
406 continuous gas emission with only sporadic pyroclastic activity. Owing to morphological  
407 restrictions not every bursting event could be observed. In spite of that, the results show  
408 a good correlation between the timing of the observed bubble bursts and that of higher  
409 amplitude tremor transients (Figure 12a). Moreover, the frequency content of the seismic  
410 signal, evaluated as the relative contribution of bands 0.95-2.15, 2.15-3.35 and 3.35-5.5 Hz,  
411 was observed to change slightly with time. This was particularly evident on high ampli-  
412 tude tremor bursts (Figure 12a). However, despite the apparent higher component of  
413 low frequency energy on tremor peaks, there was no consistent variation in the frequency  
414 content of low and high amplitude tremor (Figure 12b). This characteristic has been  
415 also observed in other periods, such as December 2004-January 2005. Hence, the higher  
416 amplitude tremor transients are not distinguishable based upon their frequency content  
417 alone.

#### 5.4. Statistics of the tremor

418 In order to assess the fluctuations in amplitude of the seismic tremor, three statistical  
419 parameters have been calculated from a high resolution RSAM of the seismic signal: 1)  
420 RSAM mean ( $\overline{rsam}$ ), 2) RSAM standard deviation ( $\sigma$ ), and 3) rate of high amplitude  
421 tremor burst ( $\#events$ ). Details of the procedure to calculate them can be found in Ap-  
422 pendix A. One of the advantages of processing the seismicity based on a high resolution  
423 RSAM time series, obtained from time-windows of 10 seconds duration, is that it allowed

424 the identification and counting of the tremor events. In addition, the RSAM mean and  
425 standard deviation yielded information about the difference in amplitude of the back-  
426 ground low amplitude tremor and higher amplitude events, as well as the occurrence of  
427 exceptional bigger events. As shown in the example of Figure 13, all three parameters  
428 vary greatly,  $\overline{rsam}$  from 16 to 36,  $\sigma$  from 4 to more than 8, and #events from 1 to 2.4 per  
429 minute. Also, the combination of their values varies continuously with time. These vari-  
430 ations indicate that, unlike the more steady frequency content of the tremor (Figure 11),  
431 its amplitude can change considerably over a short time span. Further, the combination  
432 of these parameters can describe relevant characteristics of the tremor waveform. For  
433 instance, low events rate along with low RSAM mean and high standard deviation (e.g.  
434 area (a) in Figure 13) represent relatively less frequent high amplitude discrete events  
435 with well defined starts and ends (not overlapping with each other). It is noteworthy  
436 that if the duration of the window used in the calculations of the statistics is short, one  
437 single big event can increase the standard deviation substantially (as observed in area (b)  
438 of Figure 13). The opposite arrangement, with high event rate and very low standard  
439 deviation (e.g. area (c) in Figure 13), represents tremor with a steady envelope in which  
440 events of similar amplitude occur more frequently; the amplitude of these events can be  
441 determined by the magnitude of the RSAM mean.

442 Although the overall trend of events per minute might display a rough correlation with  
443 the amplitude of the RSAM (e.g. Figure 13), during November 2004–April 2005 neither  
444 the RSAM standard deviation nor the events rate showed a consistent correlation with  
445 the RSAM amplitude. An example of this is shown in Figure 14. During the distinct  
446 changes in RSAM amplitude in January 2005, both the standard deviation and the event

447 rate showed patterns different to that of the RSAM mean. Some noteworthy features  
448 are those that occurred on the 14–15 and 29–30 January (areas (a) and (c) in Figure 14,  
449 respectively), in which the number of events per hour increased and subsequently de-  
450 creased sharply, and correlated with the inverse fluctuations in standard deviation; the  
451 RSAM displayed a rather different behavior. In turn, the peaks in RSAM amplitude and  
452 standard deviation denote the occurrence of relatively high amplitude tremor events on  
453 a time span of a few hours (e.g. 9-12hrs 14 January, 15-16hrs 30 January). The trough  
454 in standard deviation accompanied by higher RSAM denotes banded tremor (e.g. 29–30  
455 Jan). On the 20 January (area (b) in Figure 14), however, the tremor event rate followed  
456 the decrease in RSAM mean amplitude, in the same manner as all frequency bands did  
457 (Figure 10), denoting a considerable drop in seismic activity.

458 Hence, the three statistical parameters presented here yield meaningful information that  
459 can be used in the analysis of seismic variations and its relation with volcanic activity.

### 5.5. Volcano-tectonic earthquakes

460 Volcano-tectonic (VT) earthquakes, also called high-frequency events [McNutt,  
461 2000, 2005], are uncommon in the seismicity of Villarrica volcano. Only a few of these  
462 events are reported every month during periods of low (background) activity (between  
463 1-3, Calder *et al.* [2004]). During the elevated volcanic activity of December 2004–July  
464 2005, however, as many as 10 and 36 VT earthquakes were identified in March and April,  
465 respectively (Table 2). The small number of stations in the local seismic network operat-  
466 ing during that time period did not allow the location of these events. Nevertheless, three  
467 major VT earthquakes that occurred on 6 April 2005 were clearly recorded by the seismic  
468 network located at neighboring active volcanoes (Llaima, Lonquimay and Calbuco volca-

469 noes) as well as in the city of Temuco. Analysis of data from 9 stations located between  
470 approximately 4 to 220 km away from the volcano, using the Hypo 71 algorithm under the  
471 software SEISAN, allowed the localization of these events within 8 km to the northeast  
472 of Villarrica crater. These results support the idea of a change in the stress state of the  
473 volcanic system. To better constrain the evolution of such changes, however, an adequate  
474 seismic array is needed around Villarrica volcano.

## 6. Magma degassing

### 6.1. SO<sub>2</sub> flux measurements

475 Despite the persistent passive degassing exhibited for more than 20 years, measurements  
476 of gas composition and emission rates have not been done regularly at Villarrica volcano.  
477 Witter *et al.* [2004] and Witter and Calder [2004] contributed most of the data available  
478 on SO<sub>2</sub> fluxes and on the gas composition of the plume. They measured sulphur dioxide  
479 fluxes using ground-based correlation spectroscopy (COSPEC) in early 2000 and 2001. For  
480 this work, the new ultraviolet spectrometer (UVS) known as FLYSPEC [Horton *et al.*,  
481 2006] was utilized. The hardware design of this UVS is based on fore-optics, electronics,  
482 the USB2000 ultraviolet spectrometer from Ocean Optics, and a computer for recording  
483 and processing of the acquired spectra. See Galle *et al.* [2002]; Horton *et al.* [2006]; Elias  
484 *et al.* [2006]; Edmonds *et al.* [2003] and references therein for more details on the design  
485 and capabilities of these instruments. The procedure for gas concentration retrieval is  
486 based on Differential Optical Absorption Spectroscopy (DOAS) [Platt, 1994], and has  
487 been described in Horton *et al.* [2006] and Elias *et al.* [2006]. Software designed specially  
488 for FLYSPEC allows the user to calibrate the instrument in the field and to get real-time  
489 SO<sub>2</sub> path-length concentrations.

490 In order to estimate the SO<sub>2</sub> flux from Villarrica volcano, ground-based traverses were  
491 carried out on the roads that surround the volcano, between 10 and 20 km away from the  
492 crater (Figure 1). These traverses were performed on six days between November 2004 and  
493 January 2005, and on three days in January-February 2006 (Table 3). Measurements on  
494 the southeast side of the volcano, that coincide with the direction of the prevailing winds,  
495 could not be completed owing to the unsuitable conditions of the road for traverses, and  
496 so those measurements were not included in this paper. The calibration of the FLYSPEC  
497 instrument was performed using a pair of calibration cells with concentrations 438 and  
498 1504 ppm m and, on some occasions, was additionally complemented with calibration  
499 cells of 170 and 704 ppm m.

500 The estimation of plume velocity is one of the primary sources of error in SO<sub>2</sub>  
501 flux measurements [Stoiber *et al.*, 1983]. For this work, the National Centers for  
502 Environmental Prediction (NCEP) Reanalysis data [Kalnay *et al.*, 1996], provided  
503 by NOAA-CIRES (Climate Diagnostics Center, Boulder, Colorado, USA; available at  
504 <http://www.cdc.noaa.gov/>), were used for wind velocity estimates. These estimates are  
505 generated by using a state-of-the-art analysis system to perform data assimilation using  
506 past data from numerical models as well as from direct observations of atmospheric con-  
507 ditions [Kalnay *et al.*, 1996]. Comparison of our observations of plume dispersion and  
508 wind direction obtained from the NCEP Reanalysis data show very good agreement. At  
509 Villarrica volcano, a few estimations of plume speed were performed observing at the  
510 plume up to 5 km away from the crater. The comparison of these estimates with the  
511 interpolated wind field extracted from the NCEP Reanalysis data yielded differences of  
512 less than 50% in wind speed. However, it is not possible to accurately quantify the error

513 using this method. Therefore, for our results we maintain the overall error of 10%-40%  
514 considered in previous studies that performed similar methodologies for SO<sub>2</sub> flux measure-  
515 ments [Stoiber *et al.*, 1983; Kyle *et al.*, 1994; Williams-Jones *et al.*, 2003; Witter *et al.*,  
516 2004; Rodríguez *et al.*, 2004]. It is worth noting that, in any reported gas flux estimate  
517 lacking direct measurement of plume speed, this error can be much higher [Rodríguez  
518 *et al.*, 2004; Williams-Jones *et al.*, 2006].

## 6.2. Results and correlation between SO<sub>2</sub> flux and RSAM

519 Sulfur dioxide emitted from the crater of Villarrica volcano during November 2004  
520 to January 2005, on individual traverses, ranged between 180 and 1500 metric tones  
521 per day (Mg/d) (Table 3, Figure 15). Daily averages ranged between 260 and 1300  
522 Mg/d; that is a 5-fold difference. This range includes values that are clearly higher than  
523 those obtained in previous studies (in 2000-2001 and 2003; Witter *et al.* [2004]; Mather  
524 *et al.* [2004]). During the period January–February 2006 the gas fluxes were considerably  
525 lower, with daily averages no higher than 300 Mg/d. These new results highlight the  
526 variability of gas emissions at Villarrica volcano. Further, they were obtained at periods  
527 with different levels of activity, which contributes to a more representative analysis and  
528 correlation with seismicity and the observable activity of the lava lake. The highest SO<sub>2</sub>  
529 flux measured coincides with high levels of activity on 15–17 January 2005 and high values  
530 of RSAM. Conversely, the lowest fluxes were measured during periods of background (low)  
531 levels of volcanic activity: February–March 2000, February 2003 and January–February  
532 2006 (Figure 15). For comparison, considering SO<sub>2</sub> emission rates of 200-500 Mg/d as  
533 representative of background levels of volcanic activity, degassing at Villarrica is lower  
534 compared to similar open-vent systems such as Masaya (350–1800 Mg/d, Duffell *et al.*

535 [2003]), Stromboli (300–1200 Mg/d, Allard *et al.* [1994]), Yasur (216–1665 Mg/d, Bani  
536 and Lardi [2007]), and Pacaya (350–2400 Mg/d, Rodríguez *et al.* [2004]), and higher than  
537 Erebus (16–71 (up to 230) Mg/d, Kyle *et al.* [1994]). (Andres and Kasgnoc [1998] reported  
538 time-averaged values of 790, 730, 900, 510 and 79 Mg/d for Masaya, Stromboli, Yasur,  
539 Pacaya and Erebus, respectively.) It is noteworthy that the total emission of sulfur dioxide  
540 is subject to the variations in SO<sub>2</sub> fluxes that these volcanoes can manifest during periods  
541 with different levels of activity. For instance, at Yasur and Villarrica the emission of sulfur  
542 dioxide can exhibit more than a 5-fold increase between low and high activity phases.

543 Measured SO<sub>2</sub> emission rates of individual traverses exhibit a high variability on any  
544 particular day (Table 3). These fluctuations are the consequence of the combination  
545 of four phenomena: 1) the unsteady emission from the lava lake, in part owing to the  
546 concurrence of ‘passive’ and explosive outgassing, 2) accumulation of gas between the  
547 magma free surface and the spatter roof, 3) the gas plume-air turbulent mixing and  
548 circulation caused by the combination of the rise of the thermal plume and wind velocity  
549 fluctuations inside the crater, and 4) non-uniform dispersion of the gas plume between  
550 the crater and the location where it is measured. In addition, a single traverse spans  
551 a time period in which the transport and dispersion of the gas plume continues and,  
552 therefore, the final gas column-abundance profile measured might not be representative  
553 of the actual plume cross-section at a specific time. All these variables contribute to the  
554 scattered results of daily SO<sub>2</sub> flux measurements.

555 SO<sub>2</sub> flux data acquired herein (2004-2006) complemented by available measurements  
556 from previous studies (Witter *et al.* [2004] and Witter and Calder [2004], in 2000-2001;  
557 Mather *et al.* [2004], in 2003) show a good positive correlation with seismic amplitude

558 (RSAM) (Table 3, Figure 15). Similar to gas flux measurements, the one-hour averaged  
 559 RSAM (calculated from 9 to 21 hours, local time) exhibits daily fluctuations. These  
 560 variations in seismic amplitude are explained by the occurrence of banded tremor and  
 561 changes in the frequency, length and amplitude of the higher amplitude seismic transients.  
 562 In particular, on days when the volcano showed elevated levels of activity (e.g. 13, 15, 17  
 563 January 2005), recognized by visual observations and high RSAM values, the emission of  
 564 SO<sub>2</sub> was consistently higher (Figure 15a). This relation establishes a strong link between  
 565 the generation of seismic tremor and outgassing activity. Similar correlations between  
 566 SO<sub>2</sub> emissions and seismic amplitude have been found at Mount Etna (Leonardi *et al.*  
 567 [2000]) and Yasur (Bani and Lardi [2007]) volcanoes.

Selected time periods in which Villarrica volcano exhibited sustained levels of activ-  
 ity allowed a better correlation between degassing and seismicity (Figure 15b, Table 4).  
 During these periods, the activity observed at the crater remained unchanging along with  
 the RSAM values. Assuming a uniform emission of SO<sub>2</sub> for each one of these periods,  
 the average of daily results improves the precision of the measurements. Hence, a linear  
 relationship SO<sub>2</sub> flux-RSAM is obtained (Figure 15b):

$$SO_2 [Mg/d] = 32.8 \cdot RSAM - 181 \quad (1)$$

568 The negative value of the ordinate-intercept (second value on the right hand side) of this  
 569 linear equation implies a lateral displacement of the regression line to the right, meaning  
 570 an abscissa-intercept (at SO<sub>2</sub>=0 Mg/d) of 5.5 RSAM units. Even though this value  
 571 represents very low levels of volcano seismicity, it is within 30% of the RSAM values  
 572 measured during background activity. Some plausible explanations for this displacement  
 573 are: 1) tremor is not only caused by outgassing activity but also by another volcano-

574 related or non-volcanic phenomenon, 2) towards very low levels of outgassing and tremor  
575 magnitude the relationship turns non-linear and the curve goes through the origin, or 3)  
576 this relationship is non-linear but there is not enough data (or it is not accurate enough)  
577 to define it. As a first approximation, this relationship certainly serves as an estimation  
578 of the gas emitted from Villarrica volcano based on measurements of tremor magnitude.

## 7. Discussion

### 7.1. Volcanic activity and crater morphology

579 Currently, the interior of the crater is characterized by a spatter roof built above the  
580 surface of the lava lake, which constitutes the bottom of the crater. The descriptions  
581 presented herein illustrate how the depth and morphology of the spatter roof in the  
582 crater of the volcano change according to the degree of volcanic activity. Elevated or  
583 increasing activity levels are characterized by scoria terraces, spatter cones or pahoehoe  
584 lavas flooding the crater. The roof itself is very unstable and can experience major changes  
585 in morphology on a time span of hours to days. Low levels (or decreasing levels) of  
586 volcanic activity are commonly linked to a partly or totally collapsed roof, deep crater floor  
587 (perhaps with no roof), or a deep funnel shaped crater with no magmatic activity directly  
588 visible from the crater rim. The latter case implies that the top of the magma column  
589 is more than 100-150 m deep below the crater rim. Hence, the position, morphology and  
590 texture of the roof can give important information regarding changes in the depth of the  
591 magma column, and the intensity and type of activity taking place during the previous  
592 few days to weeks.

593 The visibility of the lava lake is not directly related to the level of activity but instead  
594 to the depth of the lava free-surface, morphology of the spatter roof, and the occurrence

595 of roof collapses. Although strong explosions can affect the stability of the overlying roof,  
596 visual observations suggest a poor correlation between its collapses and the magnitude  
597 of explosions. However, the morphology of the spatter roof does have an effect on the  
598 apparent magnitude of explosions: they look stronger when the opening is big, by allowing  
599 pyroclastic material to be ejected freely out of the vent. Explosions also look stronger  
600 when a narrow cone-shape vent is built on top of the spatter roof during elevated levels  
601 of activity, probably owing to the high level of the lava lake.

602 The persistence and variability in outgassing styles observed at the lava lake is also  
603 evidenced by infrasound recording [Johnson *et al.*, 2004]. At Villarrica, the infrasound  
604 trace shows continuous pressure oscillations, including impulsive transients and small  
605 amplitude waveforms that repeat along the whole time series (see Figure 3 in Johnson  
606 *et al.* [2004]). The continuous bubble bursting activity observed at the surface of the  
607 lava lake reflects the highly dynamic convection of magma developed at shallow depths  
608 in the conduit. Convection of magma at the lava lake, as observed during the persistent  
609 seething magma activity, inhibits the development of a continuous crust over the lake  
610 surface. This convection is caused by the density difference between the hot and relatively  
611 gas-rich magma rising from deep levels in the magma column (or magma chamber), and  
612 the degassed and cooler magma that has reached the uppermost part of the system, which  
613 must sink back owing to its higher density [Kazahaya *et al.*, 1994; Stevenson and Blake,  
614 1998]. In addition, the intensity of the convection is enhanced by the growing bubbles  
615 that form above gas supersaturation levels, by displacing the degassed melt sideways and  
616 downward as they pass upwards. Hence, near the surface the convection is governed by  
617 bubbly to churn-type flow regime whose strength is dependent on the ascending gas flux

618 [Mudde, 2005; Xu *et al.*, 1999]. We are not aware of this very dynamic flow regime, as  
619 represented by seething magma and small lava fountains, having been observed elsewhere.  
620 The continual arrival of bubbles to the surface observed during seething magma activity,  
621 with strombolian explosions and small lava fountains taking place intermittently, suggest  
622 that bubbles ascend freely within the conduit. Coalescence of bubbles at the top of  
623 a reservoir before entering in to the upper conduit is unlikely to occur at Villarrica.  
624 Considering the conical shape and elevation of the volcanic edifice, approximately 2000 m,  
625 we suggest that the upper plumbing system of Villarrica volcano consists of a near vertical  
626 conduit of cylindrical shape, with a length greater than 2000 m.

## 7.2. The relationship between outgassing and seismicity

627 Although we have no geophysical data appropriate to study the seismic source of the  
628 tremor, we have presented evidence that supports the relationship between the seismicity  
629 and volcanic activity observed at the crater:

- 630 1. the tight correlation between visual observations of bursting events and rapid changes  
631 in tremor amplitude (transient events) (Figure 12a),
- 632 2. the predominance of low frequencies within the tremor (Figures 8,10),
- 633 3. the consistent frequency content of the low and higher amplitude tremor (Fig-  
634 ure 12b), and
- 635 4. the positive correlation between RSAM and SO<sub>2</sub> flux (Table 3, Figure 15).

636 Much work carried out at other basaltic open-vent volcanoes supports the link between  
637 tremor and outgassing activity [Ripepe *et al.*, 1996; McGreger and Lees, 2004; Ripepe  
638 *et al.*, 2001; Métaixian and Lesage, 1997; Ripepe, 1996; Ripepe *et al.*, 2002]. For instance,  
639 by using small aperture seismic arrays, Métaixian and Lesage [1997] analysed the wavefield

640 of the tremor at Masaya volcano. At the time of their study, lava lake activity was  
641 characterised by minor fountaining and audible outgassing. They concluded that the  
642 source of the tremor was seismic surface waves linked to superficial magmatic activity in  
643 Santiago crater. Ripepe *et al.* [2001] combined infrasound and seismicity to study a phase  
644 of vigorous strombolian activity at Mt Etna, and concluded that the tremor was generated  
645 by a superposition of small point sources, lasting 1-2 s, due to pressure flux instability  
646 induced by magma degassing in the shallow portion of the magma column. At Stromboli,  
647 combination of seismicity with infrasound and thermal recording has demonstrated that  
648 the source of tremor resides in the explosive outgassing activity [Ripepe *et al.*, 1996;  
649 Ripepe and Gordeev, 1999; Ripepe *et al.*, 2002].

650 The results presented herein on the activity of Villarrica volcano yield complementary  
651 insights into the results obtained from the analysis of tremor source at similar open-vent  
652 volcanoes. We suggest that the tremor at Villarrica volcano, in particular the higher  
653 amplitude transients, is produced by seismic surface waves resulting from gas ascent and  
654 outgassing activity taking place at the top of the magma column.

### 7.3. Relationship between bubble bursting styles

655 A schematic diagram of the relationship between different explosive outgassing events  
656 observed at Villarrica is proposed on the basis of their duration and strength (Figure 16).  
657 (Here, the strength of the bubble bursting activity is estimated as the height above the  
658 surface of the lava lake that pyroclastic material can reach, which is related to the mass  
659 fraction of volatiles, overpressure and magma rise velocity [Wilson, 1980].) This diagram  
660 is qualitative and is intended to clarify the identification of new styles of bubble burst

661 events described herein that, in a general sense, have been described in previous work as  
662 ‘Strombolian activity’ [Ortiz *et al.*, 2003; Witter *et al.*, 2004].

663 At one end, seething magma is considered the activity with the longest duration and low  
664 strength. Although its duration has not been constrained, on the basis of direct visual  
665 observations of the lava lake it is believed that this activity is continuous in time but  
666 with fluctuations in strength: from mild seething magma, whose lower end corresponds  
667 to gently roll-over of the lava surface, to vigorous seething magma, whose upper end is  
668 lava fountaining. Seething magma has been observed to evolve into a small lava fountain  
669 in a time span of a few seconds (see video in auxiliary material). Occasionally, the  
670 activity falls in between the two extremes, appearing as a vigorous seething magma or as  
671 a very small lava fountain (Figure 16). The occurrence of the small lava fountain activity  
672 at or near the surface of the lava lake, involving a higher amount of bubbles bursts  
673 than seething magma, indicates higher gas contents rising in packets at slightly higher  
674 velocities determined by the density of the melt-bubbles mixture. These fluctuations in  
675 gas (bubble) content in the vertical column of magma can be interpreted on the basis  
676 of waves of bubbles [Manga, 1996], in which bubbles ascending within the conduit tend  
677 to concentrate into layers (or clusters) owing to the decreasing rise speed of the bubbles  
678 as their concentration increases. Consequently, gas bubbles reach the surface in packets  
679 rather than with an uniform vertical distribution. A similar mechanism has been invoked  
680 to explain gas puffing activity at Stromboli volcano [Ripepe *et al.*, 2002]. The persistent  
681 gas puffing activity, characteristic of Stromboli volcano [Ripepe *et al.*, 2002; Harris and  
682 Ripepe, 2007b], is also included in the sketch of Figure 16.

683 At the other extreme, strombolian explosions are relatively short and display a broad  
684 range of behavior, from relatively mild single bubble bursts to major explosions that  
685 can send pyroclastic material hundreds of meters above the lava lake. The link between  
686 strombolian explosions and seething magma is not a continuum. Strombolian explosions  
687 are considered isolated events that interrupt the seething magma activity.

688 Although Hawaiian style lava fountaining is not part of the common activity seen during  
689 periods of volcanic quiescence, in which the lava is confined inside the crater, it is included  
690 in Figure 16 to map the relation with respect to the other types of primary bubble burst  
691 seen at Villarrica. For instance, strombolian explosions with very high bursting frequency  
692 can look similar to a Hawaiian style lava fountain (e.g. Chester *et al.* [1985], p. 126–  
693 130). Parfitt and Wilson [1995] stated that it is the combination of the rise speed of the  
694 magma and the degree of bubble coalescence that defines the respective style of activity,  
695 where the magma gas content and viscosity influence only the conditions in which the  
696 transition between these two styles occurs. Hawaiian-style lava fountaining requires a  
697 relatively high magma rise speed and little coalescence of bubbles, which cause a highly  
698 vesicular melt that disrupts at deep levels (high pressure), whereas strombolian explosions  
699 are caused by a slow rise of the magma in which coalescence plays a fundamental role  
700 in the generation of large bubbles that disrupt the magma at the magma-atmosphere  
701 surface. A two-fold change in magma rise speed might be sufficient for the transition  
702 between one style of activity to the other [Parfitt and Wilson, 1995]. Changes in the  
703 physical properties of magma are plausible within the open system of Villarrica volcano.  
704 They can be a consequence of either entry of new gas-rich batches of magma into the  
705 system, or heterogeneous magma distribution and convection in a magma chamber, that

706 could eventually vary the properties of the magma rising into the upper plumbing system.  
707 Hence, entry of magma with higher gas content and lower density into the system will  
708 exhibit a higher rise speed and, consequently, increase the gas flow rate within the upper  
709 part of the conduit. Depending on the extent of these changes, the outgassing activity  
710 can evolve from seething magma to lava fountaining of specific height and extrusion rate.

#### 7.4. Volcano monitoring

711 Seismic tremor is the most common and continuous type of seismicity related to the  
712 activity of Villarrica volcano. Its amplitude, evaluated as RSAM units, has been shown  
713 to correlate well with the level of the activity observed at the summit. For instance,  
714 the different features shown in Figure 8 between January and April 2005 correlate with  
715 the increase in activity and evolution of the spatter roof (Table 1). Changes in seismic  
716 amplitude were also part of the elevated activity of 1999 [Calder *et al.*, 2004], as well as  
717 of the rapid changes in the level of the magma column observed at the end of 2000 [Ortiz  
718 *et al.*, 2003].

719 The abnormal sequence of events that took place in 2000 [Ortiz *et al.*, 2003] is an  
720 example of variations in the type of volcanic activity accompanied by clear changes in  
721 the frequency distribution of the seismicity. These results and our analysis of the activity  
722 during November 2004–April 2005 show that the amplitude and frequency distribution of  
723 the tremor can be good indicators of the variations in the activity of Villarrica volcano.  
724 However, during this time the relationship between the variations in frequency content  
725 and type or evolution of the volcanic activity was not straightforward. We hypothesize  
726 that elevated high frequencies (2.15–5.5 Hz) are caused by a rise in the level of the lava  
727 lake, perhaps interacting strongly with the spatter roof. Support for this idea comes from

728 the positive correlation between gas flux and seismic amplitude that is in accordance  
729 with stronger levels of degassing, and therefore, with vertical expansion of the magma  
730 column. Moreover, during one of the periods that exhibit high frequencies, 15-19 January,  
731 instability and collapse of the roof was directly observed.

732 Volcano-tectonic (VT) earthquakes are uncommon in the background seismicity at Vil-  
733 larrica volcano. VT-type seismicity is the result of rock failure caused by an increase in  
734 pressure within the magmatic system that is transferred to the country rock [McNutt,  
735 2005]. Such a pressure increase can occur due to new inputs of magma into the sys-  
736 tem and/or increasing degassing. The eruption in 1971, which involved the opening of  
737 a fracture in the upper part of the edifice and the generation of fissure-fed lava flows,  
738 is an example of this pressure rise and subsequent magma intrusion. If this fracturing  
739 occurs at deeper levels, it can lead to emplacement of magma as dikes or sills, which may  
740 reach the surface and initiate a flank eruption. Furthermore, because of the open state  
741 of the system, the occurrence of a shallow magma intrusion or increasing degassing will  
742 be accompanied by variations in the activity observed at the summit. The elevated SO<sub>2</sub>  
743 flux measured during January and the appearance of VT earthquakes during March-April  
744 2005, were strong evidence of an increase in the pressure of the system related to stronger  
745 degassing, perhaps caused by the supply of volatile-rich magma. It is not possible to  
746 know, however, how close the volcano was from starting a renewed eruptive episode. This  
747 anomalous episode did not last for long, as the VT earthquakes vanished in June and  
748 towards the end of July the activity at the crater declined to background levels (Table 4).

749 In addition to seismic characteristics, visual observations of the activity in the crater give  
750 valuable information regarding the level of the magma column and its variations, as well

751 as the style of activity taking place. Some important parameters worth considering are: a)  
752 the position and morphology of the spatter roof, b) the texture of its surface, c) presence  
753 and texture of new material on the crater rim, d) frequency of the big explosions, and e) the  
754 height above the crater floor that ejected pyroclasts can reach. Furthermore, continual  
755 measurements of SO<sub>2</sub> fluxes would be a significant contribution to the monitoring of  
756 degassing levels at Villarrica. The correlation of these gas data with seismicity and visual  
757 observations of activity in the crater offer a fuller appreciation of the current state and  
758 variations of the volcanic activity, and help to understand the evolution of the magmatic  
759 system.

## 8. Conclusions

760 Villarrica is a basaltic to basaltic-andesite open-vent stratovolcano with a very active  
761 lava lake and persistent gas emission. Three primary styles of bubble bursting have  
762 been recognized at the magma free surface: seething magma, small lava fountains, and  
763 strombolian explosions. Seething magma is a distinctive activity of Villarrica volcano,  
764 which may contrast to the activity observed at other open-vent volcanoes, and involves  
765 the continuous arrival and bursting of gas bubbles with radii generally lower than 5 m.  
766 Strombolian explosions can be variable in duration, from a single bubble burst (although  
767 rare) to a bursting sequence (>15 s) derived from the ascent of major gas slugs. Small lava  
768 fountains are intermediate in style, length and strength of the bursting event, between  
769 seething magma and strombolian explosions. Long duration lava fountains (>30 s) must  
770 result from large gas pockets that reach the surface of the lava lake. The interaction  
771 of bubble bursting activity with the spatter roof can be manifested by two other types  
772 of events: gas jetting and 'splashing' magma. The strength and style of the outgassing

773 activity are associated with variations in the height of the magma column, leaving an  
 774 imprint in the position, morphology and texture of the spatter roof.

775 The activity at the crater of Villarrica volcano shows fluctuations on the scale of months  
 776 to years. Slow increase in the activity at Villarrica, as observed during November 2004 and  
 777 May 2005, exhibited a rise in the magma column reflected in the position of the spatter  
 778 roof (at least 60 m), higher SO<sub>2</sub> fluxes (we measured up to 1500 tons/day), and an increase  
 779 in the frequency and strength of strombolian explosions. This evolution was also evidenced  
 780 in the higher tremor amplitude as well as the presence of VT-earthquakes. Hence, typical  
 781 background (low) levels of activity correspond to a lava lake located >80 m below the  
 782 crater rim, small and/or blocky morphology of the roof, seismic amplitude (RSAM) lower  
 783 than 25 units, few VT-earthquakes, and daily averages of SO<sub>2</sub> emissions lower than 600  
 784 tons/day. Elevated levels of activity show fluctuations in the height of the magma column,  
 785 changes in the morphology of the spatter roof, and changes in the characteristics of the  
 786 bubble bursting activity. Available SO<sub>2</sub> flux data also suggest an increase in gas emission  
 787 rates during periods of elevated activity.

## Appendix A: Details of the calculation of the tremor statistics

The Real Time Seismic Amplitude (RSAM) is calculated over the whole time series by means of a moving average window in the time domain. For a particular time,  $t$ , the RSAM is calculated as follow:

$$RSAM(t) = \frac{1}{T} \sum_{l=t-\frac{T}{2}}^{t+\frac{T}{2}} |s(l)| \quad (2)$$

788 where  $T$  is the duration of the moving time-window and  $|s(t)|$  corresponds to the absolute  
 789 value of the seismic signal. The length of the resulting RSAM time series depends on

790 the overlap (or time step) of the moving window. The selection of both the duration  
791 ( $T$ ) and time step determines the resolution of the RSAM and affects the results and  
792 meaning of the analysis. In particular, it is desirable to be able to identify each of the  
793 higher amplitude transients (or events) of the tremor, thereby the rate of events can be  
794 calculated. Accordingly, values of 10 and 7.5 seconds (25% overlap) for the duration of  
795 the window and time step, respectively, have been found appropriate for the statistical  
796 analysis of tremor amplitude at Villarrica volcano (Figures 13-14).

797 Once the RSAM( $t$ ) has been obtained, another moving window is applied over the  
798 RSAM time series in which the mean ( $\overline{rsam}$ ) and standard deviation ( $\sigma$ ) are calculated.  
799 The duration ( $\Delta T$ ) and time step ( $\delta t$ ) of this window are dependent on the objective of  
800 the analysis. For instance, the study of the characteristics and waveform of the tremor,  
801 as well as the hourly fluctuations in its amplitude, were carried out using  $\Delta T = 300s$  and  
802  $\delta t = 180s$  (Figure 13). To study long-term variations,  $\Delta T = 900s$  and  $\delta t = 600s$  were  
803 chosen (Figure 14).

804 A third parameter, the rate of higher amplitude events, was calculated by counting  
805 the amount of these events over a time-window of duration equivalent to  $\Delta T$ . An event  
806 corresponds to a peak in the RSAM time series whose magnitude is above the average  
807 ( $\overline{rsam}$ ) calculated for that window. To identify the peaks, a forward finite difference was  
808 performed beforehand in the RSAM time series, that was used to find those elements of  
809 the series with a preceding positive derivative and followed by a negative derivative.

### 810 **Acknowledgments.**

811 We acknowledge the Southern Chile Volcano Observatory (OVDAS-SERNAGEOMIN)  
812 for their support and seismic data. Keith Horton is thanked for helping with the setting

813 up of the FLYSPEC instrument, and significant assistance with the software for processing  
814 of the spectral data.

## References

- 815 Adams, N. K., Houghton, B. F., and Hildreth, W. (2006). Abrupt transitions during  
816 sustained explosive eruptions: examples from the 1912 eruption of Novarupta, Alaska.  
817 *Bulletin of Volcanology*, **69**, 189–206.
- 818 Allard, P., Carbonnelle, J., Mètrich, N., Loyer, H., and Zettwoog, P. (1994). Sulphur  
819 output and magma degassing budget of stromboli volcano. *Nature*, **368**, 326–329.
- 820 Andres, R. and Kasgnoc, A. (1998). A time-averaged inventory of subaerial volcanic sulfur  
821 emissions. *Journal of Geophysical Research*, **103**(D19), 22,251–25,261.
- 822 Andronico, D., Branca, S., Calvari, S., Burton, M., Caltabiano, T., Corsaro, R. A., Carlo,  
823 P. D., Garfi, G., Lodato, L., Miraglia, L., Murè, F., Neri, M., Pecora, E., Pompilio,  
824 M., Salerno, G., and Spampinato, L. (2005). A multi-disciplinary study of the 2002–03  
825 Etna eruption: insights into a complex plumbing system. *Bulletin of Volcanology*, **67**,  
826 314–330.
- 827 Aster, R., Mah, S., Kyle, P., McIntosh, W., Dunbar, N., Johnson, J., Ruiz, M., and  
828 McNamara, S. (2003). Very long period oscillations of Mount Erebus volcano. *Journal*  
829 *of Geophysical Research*, **108**(B11), 2522–2544.
- 830 Bani, P. and Lardi, M. (2007). Sulphur dioxide emission rates from Yasur volcano,  
831 Vanuatu archipelago. *Geophysical Research Letters*, **34**(L20309).
- 832 Bertagnini, A., Mètrich, N., Landi, P., and Rosi, M. (2003). Stromboli volcano (Aeolian  
833 Archipelago, Italy): An open window on the deep-feeding system of a steady state

- 834 basaltic volcano. *Journal of Geophysical Research*, **108**(B7), 2336–2350.
- 835 Bluth, G., Shannon, J. M., Watson, I., Prata, A., and Realmuto, V. (2007). Development  
836 of an ultra-violet digital camera for volcanic SO<sub>2</sub> imaging. *Journal of Volcanology and*  
837 *Geothermal Research*, **161**, 47–56.
- 838 Bohm, M., Luth, S., Echtler, H., Asch, G., Bataille, K., Bruhn, C., Rietbrock, A., and  
839 Wigger, P. (2002). The Southern Andes between 36 and 40 S latitude: seismicity and  
840 average seismic velocities. *Tectonophysics*, **356**, 275–289.
- 841 Calder, E., Harris, A., Peña, P., Pilger, E., Flynn, L., Fuentealba, G., and Moreno, H.  
842 (2004). Combined thermal and seismic analysis of the Villarrica volcano lava lake,  
843 Chile. *Revista Geológica de Chile*, **31**(2), 259–272.
- 844 Calvari, S., Spampinato, L., and Lodato, L. (2006). The 5 April 2003 vulcanian parox-  
845 ysmal explosion at Stromboli volcano (Italy) from field observations and thermal data.  
846 *Journal of Volcanology and Geothermal Research*, **149**(1-2), 160–175.
- 847 Casertano, L. (1963). Activity of Villarrica volcano in the current century. *Boletín*  
848 *Universidad de Chile*, **40**, 22–28, 48–54. (in spanish).
- 849 Chester, D., Duncan, A., Guest, J., and Kilburn, C. (1985). *Mount Etna: the anatomy of*  
850 *a volcano*. Chapman and Hall Ltd.
- 851 Clavero, J. and Moreno, H. (2004). Evolution of Villarrica volcano. In L. E. Lara and  
852 J. Clavero, editors, *Villarrica volcano (39.5° S), Southern Andes, Chile*, Servicio Na-  
853 cional de Geología y Minería, Boletín No. 61, pages 17–27.
- 854 Duffell, H. J., Oppenheimer, C., Pyle, D. M., Galle, B., McGonigle, A. J., and Burton,  
855 M. R. (2003). Changes in gas composition prior to a minor explosive eruption at Masaya  
856 volcano, Nicaragua. *Journal of Volcanology and Geothermal Research*, **126**, 327–229.

- 857 Edmonds, M., Herd, R., Galle, B., and Oppenheimer, C. (2003). Automatic, high time-  
858 resolution measurements of SO<sub>2</sub> flux at Soufrière Hills volcano, Montserrat. *Bulletin*  
859 *of Volcanology*, **65**, 578–586.
- 860 Elias, T., Sutton, A. J., Oppenheimer, C., Horton, K. A., Garbeil, H., Tsanev, V., Mc-  
861 Gonigle, A. J. S., and Williams-Jones, G. (2006). Comparison of COSPEC and two  
862 miniature ultraviolet spectrometer systems for SO<sub>2</sub> measurements using scattered sun-  
863 light. *Bulletin of Volcanology*, **68**, 313–322.
- 864 Endo, T. and Murray, T. (1991). Real-time seismic amplitude measurement (RSAM): A  
865 volcano monitoring and prediction tool. *Bulletin of Volcanology*, **53**, 533–545.
- 866 Falsaperla, S., Alparone, S., D’Amico, S., Granzia, G., Ferrari, F., Langer, H., Sgroi,  
867 T., and Spampinato, S. (2005). Volcanic tremor at Mt. Etna, Italy, preceding and  
868 accompanying the eruption of July-August, 2001. *Pure and Applied Geophysics*, **162**,  
869 2111–2132.
- 870 Fischer, T. P., Roggensack, K., and Kyle, P. R. (2002). Open and almost shut case for  
871 explosive eruptions: vent processes determined by SO<sub>2</sub> emission rates at Karymsky  
872 Volcano, Kamchatka. *Geology*, **30**(12), 1059–1062.
- 873 Fuentealba, G. and Peña, P. (1998). Instalación y puesta en funcionamiento del sistema  
874 de monitoreo sismológico del volcán Villarrica, durante el primer semestre de 1998.  
875 Unpublished, Servicio Nacional de Geología y Minería. 13p.
- 876 Fuentealba, G., Peña, P., and Calder, E. (2000). Sustained tremor, open system degassing  
877 and annual perturbations at the Villarrica volcano lava lake, Chile. In *IX Congreso*  
878 *Geológico Chileno*.

- 879 Galle, B., Oppenheimer, C., Geyer, A., McGonigle, A., Edmonds, M., and Horrocks, L.  
880 (2002). A miniaturised ultraviolet spectrometer for remote sensing of SO<sub>2</sub> fluxes: a new  
881 tool for volcano surveillance. *Journal of Volcanology and Geothermal Research*, **119**(1),  
882 241–254.
- 883 Gerlach, T. M. (1986). Exsolution of H<sub>2</sub>O, CO<sub>2</sub>, and S during eruptive episodes at Kilauea  
884 volcano, Hawaii. *Journal of Geophysical Research*, **91**, 12177–12185.
- 885 Harris, A. and Ripepe, M. (2007a). Synergy of multiple geophysical approaches to unravel  
886 explosive eruption conduit and source dynamics ñ A case study from Stromboli. *Chemie  
887 der Erde Geochemistry*, **67**, 1–35.
- 888 Harris, A. and Ripepe, M. (2007b). Temperature and dynamics of degassing at Stromboli.  
889 *Journal of Geophysical Research*, **112**(B03205).
- 890 Harris, A. J., Carniel, R., and Jones, J. (2005). Identification of variable convective  
891 regimes at Erta Ale lava lake. *Journal of Volcanology and Geothermal Research*, **142**,  
892 207–223.
- 893 Hickey-Vargas, R., López-Escobar, L., Moreno, H., Clavero, J., Lara, L., and Sun, M.  
894 (2004). Magmatic evolution of the villarrica volcano. In L. E. Lara and J. Clavero, edi-  
895 tors, *Villarrica volcano (39.5° S), Southern Andes, Chile*, Servicio Nacional de Geología  
896 y Minería, Boletín No. 61, pages 39–45.
- 897 Horton, K., Williams-Jones, G., Garbeil, H., Elias, T., Sutton, A., Mouginiis-Mark, P.,  
898 Porter, J., and Clegg, S. (2006). Real-time measurement of volcanic SO<sub>2</sub> emissions:  
899 Validation of a new UV correlation spectrometer (FLYSPEC). *Bulletin of Volcanology*,  
900 **68**, 323–327.

- 901 Johnson, J. B., Aster, R. C., and Kyle, P. R. (2004). Volcanic eruptions observed with  
902 infrasound. *Geophysical Research Letters*, **31**, L14604.
- 903 Kalnay, E., M. Kanamitsu, R. K., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha,  
904 S., White, G., Woollen, J., Zhu, Y., A. Leetmaa and, B. R., Chelliah, M., Ebisuzaki,  
905 W., Higgins, W., Janowiak, J., Mo, K., Ropelewski, C., Wang, J., Jenne, R., and  
906 Joseph, D. (1996). The ncep/ncar 40-year reanalysis project. *Bulletin of the American  
907 Meteorological Society*, **77**(3), 437–471.
- 908 Kazahaya, K., Shinohara, H., and Saito, G. (1994). Excessive degassing of Izu-Oshima  
909 volcano: magma convection in a conduit. *Bulletin of Volcanology*, **56**, 207–216.
- 910 Koyanagi, R., Chouet, B., and Aki, K. (1987). Origin of volcanic tremor in Hawaii, I,  
911 Compilation of seismic data from the Hawaiian Volcano Observatory, 1972 to 1985.  
912 *U.S. Geological Survey Professional Paper*, **1350**, 1221–1258.
- 913 Kyle, P., Sybeldon, L., McIntoch, W., Meeker, K., and Symonds, R. (1994). Sulfur dioxide  
914 emission rates from Mount Erebus, Antarctica. In P. Kyle, editor, *Volcanological and  
915 Environmental Studies of Mount Erebus, Antarctica.*, volume 66 of *Antartic Research  
916 Series*, pages 69–82.
- 917 Lautze, N. C. and Houghton, B. F. (2007). Linking variable explosion style and magma  
918 textures during 2002 at Stromboli volcano, Italy. *Bulletin of Volcanology*, **69**, 445–460.
- 919 Lavenu, A. and Cembrano, J. (1999). Compressional and transpressional-stress pattern  
920 for Pliocene and Quaternary brittle deformation in fore arc and intra-arc zones (Andes  
921 of Central and Southern Chile). *Journal of Structural Geology*, **21**, 1669–1691.
- 922 Leonardi, S., Gresta, S., and Mulargia, F. (2000). Cross-correlation between volcanic  
923 tremor and SO<sub>2</sub>: Flux data from Mount Etna volcano, 1987-1992. *Physics and Chem-*

- 924 *istry of the Earth (A)*, **25**(9-11), 737–740.
- 925 López-Escobar, L., Cembrano, J., and Moreno, H. (1995). Geochemistry and tectonics  
926 of the Chilean Southern Andes basaltic Quaternary volcanism (37°-46° S). *Revista*  
927 *Geológica de Chile*, **22**(2), 219–234.
- 928 Manga, M. (1996). Waves of bubbles in basaltic magmas and lavas. *Journal of Geophysical*  
929 *Research*, **101**(B8), 17,457–17,465.
- 930 Mather, T. A., Tsanev, V. I., Pyle, D. M., McGonigle, A. J. S., Oppenheimer, C.,  
931 and Allen, A. G. (2004). Characterization and evolution of tropospheric plumes from  
932 Láscar and Villarrica volcanoes, Chile. *Journal of Geophysical Research (Atmospheres)*,  
933 **109**(D18), D21303.
- 934 McGreger, A. D. and Lees, J. M. (2004). Vent discrimination at Stromboli volcano, Italy.  
935 *Journal of Volcanology and Geothermal Research*, **137**, 169–185.
- 936 McNutt, S. (2000). Seismic Monitoring. In H. Sigurdsson, J. Stix, B. Houghton, S. R.  
937 McNutt, and H. Rymer, editors, *Encyclopedia of Volcanoes*, pages 1095–1119. Academic  
938 Press.
- 939 McNutt, S. R. (2005). Volcanic seismology. *Annual Review of Earth and Planetary*  
940 *Sciences*, **32**, 461–491.
- 941 Métaixian, J. and Lesage, P. (1997). Permanent tremor of Masaya volcano, Nicaragua:  
942 wave field analysis and source location. *Journal of Geophysical Research*, **102**(B10),  
943 22529–22545.
- 944 Moreno, H., Clavero, J., and Lara, L. (1994). Explosive post-glacial activity of Villarrica  
945 volcano, southern Andes (39°25' S). In *7° Congreso Geológico Chileno*. University of  
946 Concepción. (in Spanish).

- 947 Mudde, R. F. (2005). Gravity-driven bubbly flows. *Annual Review of Fluid Mechanics*,  
948 **37**, 393–423.
- 949 Oppenheimer, C. and Yirgu, G. (2002). Thermal imaging of an active lava lake: Erta  
950 ëAle volcano, Ethiopia. *International Journal of Remote Sensing*, **23**(22), 4777–4782.
- 951 Ortiz, R., Moreno, H., García, A., Fuentealba, G., Astiz, M., Peña, P., Sánchez, N., and  
952 Tárraga, M. (2003). Villarrica volcano (Chile): characteristics of the volcanic tremor  
953 and forecasting of small explosions by means of a material failure method. *Journal of*  
954 *Volcanology and Geothermal Research*, **128**, 247–259.
- 955 Parfitt, E. and Wilson, L. (1995). Explosive volcanism IX. the transition between  
956 Hawaiian-style lava fountaining and strombolian explosive activity. *Geophysical Journal*  
957 *International*, **121**, 226–232.
- 958 Parfitt, E. A. (2004). A discussion of the mechanisms of explosive basaltic eruptions.  
959 *Journal of Volcanology and Geothermal Research*, **134**(1-2), 77–107.
- 960 Patrick, M. R., Harris, A. J. L., Ripepe, M., Dehn, J., Rothery, D. A., and Calvari,  
961 S. (2007). Strombolian explosive styles and source conditions: insights from thermal  
962 (FLIR) video. *Bulletin of Volcanology*, **69**(7), 769–784.
- 963 Platt, U. (1994). Differential optical absorption spectroscopy (DOAS). In M. Sigrist,  
964 editor, *Air monitoring by spectroscopic techniques. Chemical Analysis Series, vol. 127*,  
965 pages 27–84. Wiley, NewYork.
- 966 Ripepe, M. (1996). Evidence for gas influence on volcanic seismic signal recorded at  
967 Stromboli. *Journal of Volcanology and Geothermal Research*, **70**, 221–233.
- 968 Ripepe, M. and Gordeev, E. (1999). Gas bubble dynamics model for shallow volcanic  
969 tremor at Stromboli. *Journal of Geophysical Research*, **104**, 10639–10654.

- 970 Ripepe, M., Poggi, P., Braun, T., and Gordeev, E. (1996). Infrasonic waves and volcanic  
971 tremor at Stromboli. *Geophysical Research Letters*, **23**(2), 181–184.
- 972 Ripepe, M., Coltelli, M., Provitera, E., Gresta, S., Moretti, M., and Piccinini, D. (2001).  
973 Seismic and infrasonic evidences for an impulsive source of the shallow volcanic tremor  
974 at Mt. Etna, Italy. *Geophysical Research Letters*, **28**(6), 1071–1074.
- 975 Ripepe, M., Harris, A. J., and Carniel, R. (2002). Thermal, seismic and infrasonic ev-  
976 idences of variable degassing rates at Stromboli volcano. *Journal of Volcanology and*  
977 *Geothermal Research*, **118**, 285–297.
- 978 Ripepe, M., Harris, A., and Marchetti, E. (2005). Coupled thermal oscillations in explosive  
979 activity at different craters of Stromboli volcano. *Geophysical Research Letters*, **32**,  
980 L17302. doi:10.1029/2005GL022711.
- 981 Rodríguez, L. A., Watson, I. M., Rose, W. I., Branan, Y. K., Bluth, G. J., Chigna, G.,  
982 Matías, O., Escobar, D., Carn, S. A., and Fischer, T. P. (2004). SO<sub>2</sub> emissions to the  
983 atmosphere from active volcanoes in Guatemala and El Salvador, 1999–2002. *Journal*  
984 *of Volcanology and Geothermal Research*, **138**, 325–344.
- 985 Rowe, C., Aster, R., Kyle, P., Dibble, R., and Schlue, J. (2000). Seismic and acoustic  
986 observations at Mount Erebus Volcano, Ross Island, Antarctica, 1994 ñ 1998. *Journal*  
987 *of Volcanology and Geothermal Research*, **101**, 105 ñ 128.
- 988 Ryan, S. (1995). Quiescent outgassing of Mauna Loa volcano 1958-1994. In J. Rhodes  
989 and J. P. Lockwood, editors, *Mauna Loa Revealed: Structure, Composition, History, and*  
990 *Hazards*, number 92 in Geophysical Monograph, pages 95–115. American Geophysical  
991 Union.

- 992 Shinohara, H. and Witter, J. (2005). Volcanic gases emitted during mild strombolian  
993 activity of Villarrica volcano, Chile. *Geophysical Research Letters*, **32**(20), L20308.1–  
994 L20308.5.
- 995 Sparks, R. (2003). Dynamics of magma degassing. In C. Oppenheimer, D. Pyle, and  
996 J. Barclay, editors, *Volcanic Degassing*, volume 213, pages 5–22. Geological Society,  
997 London.
- 998 Stevenson, D. and Blake, S. (1998). Modelling the dynamics and thermodynamics of  
999 volcanic degassing. *Bulletin of Volcanology*, **60**, 307–317.
- 1000 Stoiber, R., Lawrence, L., Malinconico, J., and Stanley, W. (1983). Use of the correlation  
1001 spectrometer at volcanoes. In H. Tazieff and J. Sabroux, editors, *Forecasting Volcanic*  
1002 *Events*, chapter 29, pages 425–444. Elsevier, Amsterdam.
- 1003 Vergnolle, S. (1996). Bubble size distribution in magma chambers and dynamics of  
1004 basaltic eruptions. *Earth and Planetary Science Letters*, **140**, 269–279.
- 1005 Vergnolle, S. and Mangan, M. (2000). Hawaiian and Strombolian eruptions. In H. Sig-  
1006 urdsson, J. Stix, B. Houghton, S. R. McNutt, and H. Rymer, editors, *Encyclopedia of*  
1007 *Volcanoes*, pages 447–461. Academic Press.
- 1008 Williams-Jones, G., Rymer, H., and Rothery, D. A. (2003). Gravity changes and passive  
1009 SO<sub>2</sub> degassing at the Masaya caldera complex, Nicaragua. *Journal of Volcanology and*  
1010 *Geothermal Research*, **123**, 137–160.
- 1011 Williams-Jones, G., Horton, K. A., Elias, T., Garbeil, H., Mouginis, P. J., Sutton, M.  
1012 A. J., and Harris, A. J. L. (2006). Accurately measuring volcanic plume velocity with  
1013 multiple UV spectrometers. *Bulletin of Volcanology*, **68**, 328–332.

- 1014 Wilson, L. (1980). Relationships between pressure, volatile content and ejecta velocity in  
1015 three types of volcanic explosion. *Journal of Volcanology and Geothermal Research*, **8**,  
1016 297–313.
- 1017 Witter, J. B. and Calder, E. S. (2004). Magma degassing at Villarrica volcano. In  
1018 L. E. Lara and J. Clavero, editors, *Villarrica volcano (39.5° S), Southern Andes, Chile*,  
1019 Servicio Nacional de Geología y Minería, Boletín No. 61, pages 46–52.
- 1020 Witter, J. B., Kress, V. C., Delmelle, P., and Stix, J. (2004). Volatile degassing, petro-  
1021 logy, and magma dynamics of the Villarrica Lava Lake, Southern Chile. *Journal of*  
1022 *Volcanology and Geothermal Research*, **134**, 303–337.
- 1023 Xu, J., Cheng, P., and Zhao, T. (1999). Gas-liquid two-phase flow regimes in rectangular  
1024 channels with mini/micro gaps. *International Journal of Multiphase Flow*, **25**, 411–432.
- 1025 Young, S., Voight, B., and Duffell, H. (2003). Magma extrusion dynamics revealed by high-  
1026 frequency gas monitoring at Soufrière Hills volcano, Montserrat. In C. Oppenheimer,  
1027 D. Pyle, and J. Barclay, editors, *Volcanic Degassing*. Geological Society, London, Special  
1028 Publication, 213. 219–230.
- 1029 Zobin, V. (2003). *Introduction to Volcanic Seismology*. Elsevier Science B.V., first edition.  
1030 290p.

**Figure 1.** Location of Villarrica volcano in the southern Andes volcanic zone. The inset map shows some of the major regional tectonic structures (dashed lines, after Bohm *et al.* [2002]): Liquiñe-Ofqui (LOFZ), Gastre and Bio-Bio fault zones. It also shows the cities of Valdivia and Temuco (squares), and the location of other Quaternary volcanoes (triangles). Villarrica volcano is part of a NW-SE volcanic chain that also includes, to the SE, Quetrupillán and Lanín volcanoes. Stage I and Stage II calderas of Villarrica were created *ca.* 100 ka (14 ka) and 3.7 ka BP, respectively [Moreno *et al.*, 1994; Clavero and Moreno, 2004]. The summit crater is *ca.* 200 m in external diameter with an active lava lake at the bottom. The roads surrounding the volcano on which the ground-based traverses were performed for SO<sub>2</sub> measurements are also shown. VNVI is the seismometer. Contours every 100 m.

**Figure 2.** (a) Near vertical view of the crater of Villarrica volcano in November 2004 (courtesy of C.J.Tanguy). North is towards the top of the photo; the arrow indicates people (black dots) walking on the snow-covered crater rim. The interior of the crater has a diameter of  $150 \pm 10$  meters, whereas the outer rim is *ca.* 200 m across. (b) Schematic cross-section of Villarrica crater. The active lava lake represents the top of the magma column. The spatter roof adheres to the inner crater walls by accumulation of ejected spatter. A terrace (with a relatively flat surface) can be formed by accumulation of tephra during periods of elevated explosive activity.

**Figure 3.** Sequence of photographs that shows the morphological evolution of the vent during the period November 2004–April 2005: spatter roof and terraces on the crater floor a) at the end of November 2004, b) middle of January 2005, and c) end of April 2005. The position of the roof was approximately 90 m, 50 m and 30 m below the crater rim, respectively. The roof adheres to the walls of the crater above the lava free surface. In (b) a set of three terraces with concentric fractures can be seen; the outer fracture was the site of a small collapse within the next two days. Scales are approximated.

**Figure 4.** Seething magma bursting style. a) Photograph of the orifice in the spatter roof that shows part of the lava lake’s free–surface with a bubble bursting on one side. The dashed black line indicates the margin of the lake. The scale is approximate. b) Schematic of the seething magma activity, in which undulations occur in a partly encrusted lava lake (thick dash line) due to the bubbles that constantly reach the surface. In this case, two bubble bursts occur in a time frame of four seconds.

**Figure 5.** Photographs of the small lava fountain activity. a) The beginning of the fountain displays a few bubbles bursting simultaneously. b) Six seconds later, the fountain reaches more than 25 m high. Scales are approximate.

**Figure 6.** Photographs of two characteristic samples found on the crater rim, generated by explosions that occurred in December 2004 and January 2005. (a) Reticulite scoria that was part of the pyroclastic material thrown >100 m above the crater floor (19/Dec/2004 17:24 local time); by the next day, not one scoria with that texture remained on the crater rim. (b) Scoria sample showing the wide vesicle-size distribution commonly found in Villarrica scoria samples (ice axe handle for scale).

**Figure 7.** One hour of seismicity from 03-Feb-2005 beginning at 8 am (GMT). The vertical axis shows the start of the signal at 5-minute increment. Individual bursts have a wide range of amplitudes and durations, as well as emergent starts and ends.

**Figure 8.** a) Real-time Seismic-Amplitude Measurements (RSAM) and b) spectrogram of the tremor for the period November 2004–April 2005. RSAM is calculated over a moving window of length and step of 10 minutes. The spectrogram is normalized to the maximum value of the energy for every 10 min window, so it shows the relative changes in frequency distribution and dominant spectral peaks. Empty spaces represent data gaps.

**Figure 9.** Spectra of six one-hour long seismic signals that show the changes in the amplitude of the predominant peaks between November 2004 and April 2005. Vertical lines divide the spectra at frequencies 0.95, 1.35, 1.65, 2.15, 2.55, 2.85, 3.35, 5.5, 6 and 7.2 Hz. These lines were chosen in order to highlight the variations in amplitude of particular frequency ranges. Gray areas correspond to frequency bands 0.95-1.35, 1.65-2.15 and 2.55-5.5 Hz.

**Figure 10.** Amplitude variations in the frequency of the tremor during January 2005. Each trace was obtained by calculating the Fast Fourier Transform (FFT) and averaging the spectral amplitude of a 30 minutes long window moving with a step of 15 minutes. Frequency bands were chosen based on the distribution of peaks and troughs in the spectra after inspection of the whole data set. Notice the different scales on the vertical axes. Although the amplitudes are in arbitrary units, values between bands can be compared.

**Figure 11.** (a) Fluctuation patterns in seismic amplitude (RSAM). This example is from 7-8 January 2005. Many of the fluctuations display a serrated shape, with a fast increase in amplitude followed by a slow decrease. (b) Waveform plots of one-hour of seismic signal coincident with an RSAM peak and RSAM trough (indicated with arrows in (a)). (c) Their associated frequency distribution (Fourier Transform, normalized amplitude).

**Figure 12.** Example of the correlation between visual observations of explosions and seismicity at Villarrica volcano. a) Seismic tremor and amplitude of three frequency bands are illustrated considering 40 minutes of data of 17 January 2006. Arrows on top of the figure mark the time of explosions observed from the summit of the volcano. The amplitude of the frequencies was calculated with a 15 second long moving window with a step of 10 seconds. b) Distribution of the frequency content for the same signal. Higher amplitude tremor is represented by darker and bigger triangles.

**Figure 13.** Statistics and event counting of the tremor. The top graph displays a high resolution RSAM (bars) calculated with a 10 seconds long window moving with a step of 7.5 seconds. The continuous line is the average RSAM. The bottom graph shows the number of higher amplitude transients (#events) per minute (continuous line) and the standard deviation ( $\sigma$ ) of the RSAM (divided by 5, dashed line). The number of events was counted every five minutes. The average and standard deviation of the RSAM were calculated with a moving window with a duration and step of 5 and 3 minutes, respectively. Grey areas indicate example time periods with (a) high  $\sigma$  and low #events, (b) relatively high  $\sigma$  and high #events, and (c) with low  $\sigma$  but high #events. This example is from 7 January 2005 (two examples of tremor waveform and their spectra in Figure 11).

**Figure 14.** RSAM amplitude (top), number of events per hour (middle) and RSAM standard deviation (bottom) of the tremor signal in January 2005. The RSAM was calculated with a moving window with duration and step of 10 and 7.5 seconds, respectively. RSAM average and standard deviation were calculated in a 15 minutes long window moving with a step of 10 minutes. In each case, a moving average (four data points) was applied (dark line). Grey areas mark time periods with abrupt changes in RSAM amplitude, and coincide with areas marked in Figure 10.

**Figure 15.** Correlation between RSAM and SO<sub>2</sub> flux observed at Villarrica volcano. (a) Daily averages; the vertical and horizontal bars show the minimum and maximum for individual traverses and hourly RSAM, respectively (Table 3). Square symbols from Witter *et al.* [2004] and triangles from Mather *et al.* [2004]. (b) Average seismic and gas data for different periods (of a few days to weeks) with available SO<sub>2</sub> flux measurements (Table 2).

**Figure 16.** Schematic diagram of bubble burst styles that shows the relationship between the duration and strength of the events. Continuous line connections denote the continuum between events, whereas dashed line connections denote the feasible occurrence, although not observed directly at Villarrica, of a continuum between strombolian explosions or small lava fountains, and the Hawaiian-style fountaining. The grey arrow denotes the increase in total gas emission as the duration and strength of the events increases.

**Table 1.** Summary of visual observations and seismicity during November 2004-April 2005.

Date	Visual Observations	Seismicity <sup>a</sup>
9-17 November	Lava lake >100 m below crater rim; not visible. Ejected spatter rarely reached more than 20 m above the crater floor.	RSAM 10-20 units (average 15).
23 November	New spatter agglutinated to the inner wall of the vent.	Slowly increasing RSAM; up to 20 units
12-19 December	Explosions send pyroclasts onto the crater rim (>100 m above the lava lake).	RSAM over 20 units and increasing. Banded tremor appears interlaced with pulsating tremor.
3-12 January	Apparent increase in frequency of explosions ejecting material out of the vent.	Highly variable RSAM: 20-35 units and up to 40 on 8 and 9 January.
14-18 January	Spatter roof ca. 40 m higher than in November. Terrace built up with pyroclastic material falling on top of the roof. Evidence of partial concentric collapses of the roof around the vent.	Rapid increase in seismic amplitude: RSAM up to 50 units. Seismic signal shows high frequency.
21-26 January	New accumulation of spatter at the interior of the central opening. Big explosions sending bombs above crater rim. Partial concentric collapses.	RSAM 20-35 units.
February	Morphology of the spatter roof is more stable. Explosive activity similar to January. By the end of February, increase in accumulation of pyroclastic material around the vent is evident.	Average RSAM ca. 25 by the middle of Feb, but decreases to 15-22 towards the end of the month.
5-11 March	Pyroclastic material accumulating on the NE-E side of the vent. Two holes in the spatter roof.	RSAM shows positive trend, recovering values >20 units (on average).
24-30 March	Opening in the roof has consolidated into one small (ca. 5 m) circular hole. Pyroclasts ejected >100 m above the crater rim.	RSAM more stable at 20-27 units on average. Increasing amounts of VT earthquakes.
April	A spatter cone is developed on top of the roof with a hole <15 m in diameter. Explosions send bombs >150 m above the crater rim. Spatter appears around the vent on 19-20 of April.	Average RSAM stable at 20-30 units. Amount of VT earthquakes well above background (36 in the whole month).

<sup>a</sup> RSAM calculated over a time window of 10 minutes.

**Table 2.** Monthly volcano-tectonic (VT) earthquakes detected at Villarrica volcano.

Month	Number of VT
December 2004	2
January 2005	4
February 2005	2
March 2005	10
April 2005	36
May 2005	5
June 2005	5
July 2005	0

**Table 3.** Summary of the results of ground-based traverses carried out to estimate SO<sub>2</sub> emissions [Mg/d] at Villarrica volcano, along with RSAM averages for the same periods.

Date	RSAM			SO <sub>2</sub> flux			No. runs
	min	mean	max	min	mean	max	
<i>February-March 2000</i> <sup>a</sup>							
29 Jan	14.7	15.9	17.6		178		1
30 Jan	12.3	13.6	14.8	327	353	379	2
02 Feb	13.5	14.9	16.2	203	274	338	6
05 Feb	14.6	17.6	19.5	95	209	352	11
07 Feb	14.9	16.5	19.8		121		1
13 Feb	14.8	16.4	17.9	177	204	231	2
14 Feb	13.4	16.1	20.1	126	151	195	7
03 Mar	12.9	14	15.1	127	149	163	5
07 Mar	11.9	13.7	17.8		174		1
18 Mar	11.5	13.1	15.8	241	374	614	3
21 Mar	10	12.2	14.3	429	694	1115	8
<i>April-June 2000</i> <sup>b</sup>							
01 Apr	12.3	15.4	18.3	152	269	333	9
04 May	9.3	11.2	12.7	47	171	298	7
01 Jun	9	10.2	11.6		80		1
<i>January-February 2001</i> <sup>a</sup>							
15 Jan	12.2	17.3	22.8	87	118	176	8
25 Jan	20.9	24.6	30.3	257	441	961	6
28 Jan	22.2	24.7	28.6	324	564	705	7
02 Feb	24.5	28.9	30.5	565	732	939	7
<i>February 2003</i> <sup>c</sup>							
08 Feb		15.1		277	397	518	2
12 Feb		15.1		190	281	363	4
<i>November 2004</i>							
10 Nov	12.0	14.5	17.8	178	261	356	10
<i>January 2005</i>							
13 Jan	23.4	28.3	33.4	553	735	993	6
15 Jan	32.4	37.6	42.9	1176	1299	1482	4
17 Jan	36.5	40	44.6	706	951	1080	4
19 Jan	21.8	26.5	30	384	603	742	6
24 Jan	22.6	26.7	34.2	772	996	1164	4
<i>January-February 2006</i>							
17 Jan	6.2	6.8	7.5	78	122	154	8
09 Feb	8.3	9.2	10	233	262	343	6
13 Feb	9.6	11.2	12.5	86	149	232	4

<sup>a</sup> Gas data from Witter *et al.* [2004]. They used a COSPEC.

<sup>b</sup> Gas data from Witter and Calder [2004]. They used a COSPEC.

<sup>c</sup> Gas data from Mather *et al.* [2004]. They used a mini-DOAS. RSAM values from 12 November 2004.

**Table 4.** Time averaged SO<sub>2</sub> emissions [Mg/d] at Villarrica volcano alongside RSAM averages for selected periods, and the number of days over which the average was made.

Period	RSAM	SO <sub>2</sub>	No. days
Feb-Mar 2000	14.9	263	12
Jan-Feb 2001	23.9	464	4
Feb 2003	15.1	339	2
Jan 2005	27.1	778	3
15-17 Jan 2005	38.8	1125	2
Jan-Feb 2006	9.1	178	3