



## Open Research Online

### Citation

Penson, K. A.; Blasiak, P.; Horzela, A.; Duchamp, G. H. E. and Solomon, A. I. (2009). Laguerre-type derivatives: Dobiński relations and combinatorial identities. *Journal of Mathematical Physics*, 50(8) 083512.

### URL

<https://oro.open.ac.uk/18816/>

### License

None Specified

### Policy

This document has been downloaded from Open Research Online, The Open University's repository of research publications. This version is being made available in accordance with Open Research Online policies available from [Open Research Online \(ORO\) Policies](#)

### Versions

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding

## Laguerre-type derivatives: Dobiński relations and combinatorial identities

K. A. Penson,<sup>1,a)</sup> P. Blasiak,<sup>2,b)</sup> A. Horzela,<sup>2,c)</sup> G. H. E. Duchamp,<sup>3,d)</sup> and A. I. Solomon<sup>4,e)</sup>

<sup>1</sup>Laboratoire de Physique Théorique de la Matière Condensée (LPTMC), Université Pierre et Marie Curie, CNRS UMR 7600, Tour 24-2ième ét., 4 pl. Jussieu, F 75252 Paris Cedex 05, France

<sup>2</sup>H. Niewodniczański Institute of Nuclear Physics, Polish Academy of Sciences, ul. Eliasza-Radzikowskiego 152, PL 31342 Kraków, Poland

<sup>3</sup>LIPN, CNRS UMR 7030, Institut Galilée, 99 Av. J.-B. Clement, F 93430 Villetaneuse, France

<sup>4</sup>Department of Physics and Astronomy, The Open University, Milton Keynes MK7 6AA, United Kingdom and Laboratoire de Physique Théorique de la Matière Condensée (LPTMC), Université Pierre et Marie Curie, CNRS UMR 7600, Tour 24-2ième ét., 4 pl. Jussieu, F 75252 Paris Cedex 05, France

(Received 2 April 2009; accepted 28 May 2009; published online 11 August 2009)

We consider properties of the operators  $D(r, M) = a^r (a^\dagger a)^M$  (which we call generalized Laguerre-type derivatives), with  $r = 1, 2, \dots$ ,  $M = 0, 1, \dots$ , where  $a$  and  $a^\dagger$  are boson annihilation and creation operators, respectively, satisfying  $[a, a^\dagger] = 1$ . We obtain explicit formulas for the normally ordered form of arbitrary Taylor-expandable functions of  $D(r, M)$  with the help of an operator relation that generalizes the Dobiński formula. Coherent state expectation values of certain operator functions of  $D(r, M)$  turn out to be generating functions of combinatorial numbers. In many cases the corresponding combinatorial structures can be explicitly identified. © 2009 American Institute of Physics. [DOI: 10.1063/1.3155380]

### I. INTRODUCTION

Among the many ways of generalizing the ordinary derivative  $d/dx$ , the notion of the so-called Laguerre derivative<sup>1</sup> seems to be particularly fruitful. The idea is to extend the operator  $d/dx$  to a simple homogeneous counterpart  $D_x$ , which we define as in Refs. 2 and 3 as [note that here we omit the factor  $(-1)$  present in these references]

$$D_x = \frac{d}{dx} x \frac{d}{dx}. \quad (1)$$

In Refs. 1–3 many important consequences of the replacement  $d/dx \rightarrow D_x$  in the integral transform methods and in the operational calculus were investigated. The link between  $D_x$  and the Laguerre polynomials becomes clear if one notices the operational relation [see Eq. (5) in Ref. 3], which is easy to get using the amusing identity  $D_x^n = (d/dx)^n x^n (d/dx)^n$ ,<sup>4</sup>

$$e^{y D_x} x^n = n! y^n L_n \left( -\frac{x}{y} \right), \quad (2)$$

<sup>a)</sup>Electronic mail: penson@lptl.jussieu.fr.

<sup>b)</sup>Electronic mail: pawel.blasiak@ifj.edu.pl.

<sup>c)</sup>Electronic mail: andrzej.horzela@ifj.edu.pl.

<sup>d)</sup>Electronic mail: ghed@lipn-univ.paris13.fr.

<sup>e)</sup>Electronic mail: a.i.solomon@open.ac.uk.

where  $L_n(z)$  are Laguerre polynomials. This justifies *a posteriori* the name Laguerre derivative for  $D_x$ . Using Eq. (2) we may obtain the action of  $e^{\lambda D_x}$  on various functions using the different generating functions of Laguerre polynomials listed in Sec. 5.11 of Ref. 5. In particular, using the well known ordinary generating function of  $L_n(x)$  [formula (5.11.2.1) for  $\alpha=0$  in Ref. 5], one obtains<sup>6</sup>

$$e^{\lambda D_x} e^{-bx} = \frac{1}{1+b\lambda} \exp\left(-\frac{bx}{1+b\lambda}\right) \quad (3)$$

valid for  $|b\lambda| < 1$ . Analogously, using formula (5.11.2.6) in Ref. 5, one gets

$$e^{\lambda D_x} {}_1F_1([b],[1],x) = \frac{1}{(1-\lambda)^b} {}_1F_1\left([b],[1],\frac{x}{1-\lambda}\right), \quad (4)$$

with  ${}_1F_1$  as the hypergeometric function<sup>7</sup> which for many values of  $b$  specializes to elementary or known special functions.<sup>5</sup> Note that for both these examples the action of  $e^{\lambda D_x}$  results in a substitution and a prefactor which is reminiscent of the so-called Sheffer-type operators.<sup>9</sup>

We now employ the operational equivalence

$$\left[ \frac{d}{dx}, x \right] = 1 \leftrightarrow [a, a^\dagger] = 1, \quad (5)$$

where  $a$  and  $a^\dagger$  are boson annihilation and creation operators, respectively, and rewrite  $D_x \leftrightarrow D$  as

$$D = aa^\dagger a. \quad (6)$$

By going one step further we extend Eq. (6) by defining the generalized Laguerre derivative  $D(r, M)$  as

$$D(r, M) = a^r (a^\dagger a)^M \sim D_x(r, M) = \left( \frac{d}{dx} \right)^r \left( x \frac{d}{dx} \right)^M, \quad r = 1, 2, \dots, \quad M = 0, 1, \dots \quad (7)$$

These operators are the object of our present study. Although the equivalence in Eq. (7) between  $D(r, M)$  and  $D_x(r, M)$  is formal since the domains of  $a$ ,  $a^\dagger$ , and  $d/dx, x$  are different, we shall show that it provides one with an effective calculational tool.

Since  $(a^\dagger a)^M$  conserves the number of bosons, the operators  $D(r, M)$  act as monomials in boson operators which annihilate  $r$  bosons. Recent experiments in quantum optics have shown how one may produce quantum states with specified numbers of photons. This, in turn, raises the interesting possibility of producing exotic coherent states; that is, states other than the standard ones which satisfy  $a|z\rangle = z|z\rangle$ .<sup>10</sup> The current work introduces operators whose eigenstates may be used to model new coherent states which have many of the features of the standard ones and still permit explicit analytical description. The explicit forms of these new generalized coherent states can be used to evaluate relevant physical parameters, such as the photon distribution, the Mandel parameter, squeezing factors, and signal-to-noise ratio.

Much theoretical work has been devoted to the description of nonstandard coherent states; for example, the so-called nonlinear coherent states,<sup>11</sup> multiphoton coherent states,<sup>12</sup> and  $q$ -deformed coherent states.<sup>13</sup> The structure embodied in Eq. (7) is a special case of the extension of boson operators proposed in the construction of nonlinear coherent states.<sup>11</sup> In this latter reference one defines the generalized boson annihilator  $b$  by

$$b = af(a^\dagger a), \quad (8)$$

choosing the  $f(x)$  that most suits the problem in question. Evidently, for this identification  $r=1$  and  $f(x)=x^M$ . In this case the commutator is equal to

$$[D(1, M), D^\dagger(1, M)] = (a^\dagger a + 1)^{2M+1} - (a^\dagger a)^{2M+1}. \quad (9)$$

This emphasizes the fact that although  $D(1, M)$  and  $D^\dagger(1, M)$  annihilate and create one boson, respectively, they are *not* canonical boson operators (unless  $M=0$ ). Equation (9) is a special case of

$$[D(r, M), D^\dagger(r, M)] = (a^\dagger a + r)^{2M} \left( \sum_{k=1}^{r+1} |\sigma(r+1, k)| (a^\dagger a)^{k-1} \right) - (a^\dagger a)^{2M} \left( \sum_{k=1}^r |\sigma(r, k)| (a^\dagger a)^k \right), \quad (10)$$

where the  $\sigma(r, k)$  are Stirling numbers of the first kind.<sup>14</sup>

Equation (10) was obtained by using the following two equations:

$$a^r (a^\dagger)^r = \prod_{p=1}^r (a^\dagger a + p) = \sum_{k=1}^{r+1} |\sigma(r+1, k)| (a^\dagger a)^{k-1}. \quad (11)$$

The first part of Eq. (11) is readily proved by induction. To prove the second part of Eq. (11) we use the generating function for  $|\sigma(r+1, k)|$  in the form<sup>15</sup>

$$\sum_{k=1}^{r+1} |\sigma(r+1, k)| x^{r+1-k} = \prod_{p=1}^r (1 + px), \quad (12)$$

from which, by substituting  $x=1/n$  and using the first part of Eq. (11), the second part of Eq. (11) follows.

The basic objective of this work is the investigation of arbitrary powers of  $D(r, M)$  which, in turn, will allow one to evaluate Taylor-expandable functions of  $D(r, M)$ . We achieve our goal following the recently developed methods of construction of normally ordered products.<sup>16-19</sup> As we shall show, results derived in this way have a combinatorial flavor and lend themselves to a combinatorial interpretation.

This paper is organized as follows. In Sec. II we introduce generalizations of the Stirling and Bell numbers, which are well known from classical combinatorics and relate them to the normally ordered powers of operators  $D(r, M)$ . These numbers, as shown in Sec. III, may be explicitly found using generalized Dobiński relations. In Sec. IV we compare calculations of purely analytical origin with those based on methods of graph theory and give a combinatorial interpretation of our results. Examples of various applications of our approach are presented in Sec. V, while Sec. VI summarizes the paper.

## II. NORMAL ORDERING: GENERALIZED STIRLING AND BELL NUMBERS

The *normally ordered* form of  $F(a, a^\dagger)$ , denoted by  $F_{\mathcal{N}}(a, a^\dagger)$ ,<sup>20</sup> is obtained by moving all annihilators to the right using the canonical commutation relation of Eq. (5). It satisfies  $F_{\mathcal{N}}(a, a^\dagger) = F(a, a^\dagger)$ . On the other hand, the *double dot* operation  $:G(a, a^\dagger):$  means that we are applying the same ordering procedure but without taking account of the commutation relation. Conventionally the solution to the normal ordering problem is obtained if a function  $G(a, a^\dagger)$  is found satisfying,

$$F_{\mathcal{N}}(a, a^\dagger) = :G(a, a^\dagger):. \quad (13)$$

A large body of research has been recently devoted to finding the solution of Eq. (13).<sup>21</sup> A general approach that facilitates a combinatorial interpretation of quantum mechanical quantities is to use the coherent state representation. Standard coherent states

$$|z\rangle = e^{-|z|^2/2} \sum_{n=0}^{\infty} \frac{z^n}{\sqrt{n!}} |n\rangle \quad (14)$$

with the number states  $|n\rangle$  satisfying  $a^\dagger a|n\rangle = n|n\rangle$ ,  $\langle n|n'\rangle = \delta_{n,n'}$ , and  $z$  complex, are eigenstates of the annihilation operator, i.e.,  $a|z\rangle = z|z\rangle$ . The latter eigenstate property shows that having solved the normal ordering problem [Eq. (13)] for an operator  $F(a, a^\dagger)$ , we immediately find

$$\langle z|F_{\mathcal{N}}(a, a^\dagger)|z\rangle = G(z, z^*). \quad (15)$$

An early observation on how to extract combinatorial content from normally ordered forms<sup>22</sup> was based on the formula  $e^{\lambda a^\dagger a} = :e^{a^\dagger a(e^\lambda - 1)}:$ .<sup>23</sup> It led to the identification

$$\langle z|(a^\dagger a)^n|z\rangle = B(n), \quad (16)$$

where the  $B(n)$  are conventional Bell numbers described in Ref. 14. Equation (16) may be taken as a *definition* of the Bell numbers. For Stirling numbers of the second kind, we have<sup>14</sup>

$$(a^\dagger a)^n = \sum_{k=0}^n S(n, k) (a^\dagger)^k a^k \quad (17)$$

(which may also be used as a practical definition) in terms of which one defines the Bell polynomials by

$$B(n, x) = \sum_{k=0}^n S(n, k) x^k. \quad (18)$$

We have extended and developed the coherent state methodology for operators other than  $a^\dagger a$  in Refs. 16–18.

After the seminal observation by Katriel,<sup>22</sup> combinatorial methods found widespread application in this context.<sup>16–19,24</sup> We apply these methods to  $F(a, a^\dagger) = [D(r, M)]^n$ ,  $n = 1, 2, \dots$

Formally, we write  $[D(r, M)]^n$  in normally ordered form as

$$[D(r, M)]^n = \left[ \sum_{k=0}^{Mn} S_r^{(M)}(n, k) (a^\dagger)^k a^k \right] a^{rn}. \quad (19)$$

Clearly, from Eq. (19) the integers  $S_r^{(M)}(n, k)$  are generalizations of the conventional Stirling numbers of the second kind (which are recovered for  $r=0$ ,  $M=1$ ). Analogous to Eq. (18), the numbers  $S_r^{(M)}(n, k)$  serve to define the generalized Bell polynomials,

$$B_r^{(M)}(n, x) = \sum_{k=0}^{Mn} S_r^{(M)}(n, k) x^k. \quad (20)$$

Finding the explicit form of these generalized Stirling numbers will give the normally ordered form of  $[D(r, M)]^n$ . We proceed to do this in Sec. III by use of a generalization of the famous Dobiński formula.

### III. GENERALIZED DOBIŃSKI FORMULA

We first write Eq. (19) in the derivative form as

$$[D_x(r, M)]^n = \left[ \sum_{k=0}^{Mn} S_r^{(M)}(n, k) x^k \left( \frac{d}{dx} \right)^k \right] \left( \frac{d}{dx} \right)^{rn}. \quad (21)$$

Acting with the right hand side of Eq. (21) on  $e^x$ , one obtains  $B_r^{(M)}(n, x)e^x$ . The action of the left hand side of Eq. (21) on  $e^x$  is obtained by acting with the generalized Laguerre derivatives on monomials  $x^p$ ,

$$D_x(r, M)x^p = p^r p^M x^{p-r}, \quad (22)$$

where  $p^r = p(p-1)\cdots(p-r+1)$  is the falling factorial, then extending it to the  $n$ th power

$$[D_x(r, M)]^n x^p = \left[ \prod_{j=0}^{n-1} (p-rj)^r (p-rj)^M \right] x^{p-rn} \quad (23)$$

and next summing up contributions for  $x^p/p!$ ,

$$\sum_{p=0}^{\infty} [D_x(r, M)]^n \frac{x^p}{p!} = \sum_{p=rn}^{\infty} \left[ \prod_{j=0}^{n-1} (p-rj)^r (p-rj)^M \right] \frac{x^{p-rn}}{p!}. \quad (24)$$

Upon simplifying Eq. (24) leads to the Dobiński-type representation of generalized Bell polynomials,<sup>19,25,26</sup>

$$B_r^{(M)}(n, x) = e^{-x} \sum_{l=0}^{\infty} \left[ \prod_{i=1}^n (l+ir) \right]^M \frac{x^l}{l!}, \quad (25)$$

verified by direct calculation of  $\langle z | [a^r (a^\dagger a)^M]^n | z \rangle$ . The classic Dobiński formula<sup>14</sup> corresponds to  $r=0$ ,  $M=1$ ,

$$B(n, x) = e^{-x} \sum_{l=0}^{\infty} \frac{l^n x^l}{l!}. \quad (26)$$

From Eq. (25) the generalized Stirling numbers are obtained by standard Cauchy multiplication of series

$$S_r^{(M)}(n, k) = \frac{1}{k!} \sum_{j=0}^k \binom{k}{j} (-1)^{k-j} \left[ \prod_{i=1}^n (j+ir) \right]^M, \quad k=0, 1, \dots, Mn. \quad (27)$$

We point out that Eqs. (25) and (27) are the central results we need for further calculations. For practical applications it is useful to note that the generalized Stirling and Bell numbers, as well as generalized Bell polynomials, can be expressed through generalized hypergeometric functions  ${}_pF_q$ . Below we quote some examples of such relations.

$$S_1^{(M)}(n, k) = \frac{(-1)^k (n!)^M}{k!} \cdot {}_{M+1}F_M \left( \underbrace{[-k, n+1, \dots, n+1]}_{M \text{ times}}, \underbrace{[1, \dots, 1]}_{M \text{ times}}, 1 \right), \quad (28)$$

$$B_1^{(M)}(n, x) = e^{-x} (n!)^M \cdot {}_M F_M \left( \underbrace{[n+1, \dots, n+1]}_{M \text{ times}}, \underbrace{[1, \dots, 1]}_{M \text{ times}}, x \right). \quad (29)$$

The numbers  $B_1^{(M)}(n) = B_1^{(M)}(n, 1)$  can be shown to be related to the numbers  $B_{p,p}(n)$  (introduced in Refs. 16 and 17) characterizing the normal order of  $[(a^\dagger)^p a^p]^n$  by the formula,

$$B_1^{(M)}(n) = B_{n,n}(M+1), \quad (30)$$

as seen by comparing Eq. (2.6) in Ref. 17 with Eq. (25) of the present work

$$\begin{aligned}
 B_2^{(M)}(n,x) &= \frac{1}{\pi^{M/2}} 2^{Mn} e^{-x} (n!)^M \cdot {}_M F_{M+1}(\underbrace{[n+1, \dots, n+1]}_{M \text{ times}}, \underbrace{[1, \dots, 1]}_{M \text{ times}}, 1/2, x^2/4) \pi^{M/2} \\
 &\quad + 2^M [\Gamma(n+3/2)]^M \cdot x \cdot {}_M F_{M+1}(\underbrace{[n+3/2, \dots, n+3/2]}_{M \text{ times}}, \underbrace{[3/2, \dots, 3/2]}_{M+1 \text{ times}}, x^2/4), \tag{31}
 \end{aligned}$$

$$\begin{aligned}
 B_3^{(M)}(n,x) &= \frac{1}{2^{M+1} [\pi \Gamma(2/3)]^M} e^{-x} \cdot (2^{M+1} 3^{Mn} [\pi n! \Gamma(2/3)]^M \\
 &\quad \cdot {}_M F_{M+2}(\underbrace{[n+1, \dots, n+1]}_{M \text{ times}}, \underbrace{[1, \dots, 1]}_{M \text{ times}}, 1/3, 2/3, x^3/27) + 2 \cdot 3^{M(n+3/2)} (\Gamma^2(2/3) \Gamma(n+4/3))^M \\
 &\quad \cdot x \cdot {}_M F_{M+2}(\underbrace{[n+4/3, \dots, n+4/3]}_{M \text{ times}}, \underbrace{[4/3, \dots, 4/3]}_{M+1 \text{ times}}, 2/3, x^3/27) + 3^{M(n+1)} [\pi \Gamma(n+5/3)]^M \\
 &\quad \cdot x^2 \cdot {}_M F_{M+2}(\underbrace{[n+5/3, \dots, n+5/3]}_{M \text{ times}}, \underbrace{[5/3, \dots, 5/3]}_{M+1 \text{ times}}, 4/3, x^3/27)). \tag{32}
 \end{aligned}$$

We conjecture that in general  $B_r^{(M)}(n,x)$  is a combination of  $r$  hypergeometric functions of type  ${}_M F_{M+r-1}$  of argument  $x^r/r^r$ .

Examples of numbers resulting from Eqs. (29)–(32) for  $n=0, \dots, 6, \dots$  are

$$M = 1, \quad B_1^{(1)}(n) = 1, 2, 7, 34, 209, 1546, 13\,227, \dots,$$

$$M = 2, \quad B_1^{(2)}(n) = 1, 5, 87, 2971, 163\,121, 12\,962\,661, \dots,$$

$$M = 3, \quad B_1^{(3)}(n) = 1, 15, 1657, 513\,559, 326\,922\,081, 363\,303\,011\,071, \dots,$$

$$M = 2, \quad B_2^{(2)}(n) = 1, 10, 339, 23\,395, 2\,682\,076, 457\,112\,571, 107\,943\,795\,145, \dots,$$

$$\begin{aligned}
 M = 3, \quad B_2^{(3)}(n) \\
 = 1, 37, 9415, 7\,063\,615, 11\,360\,980\,081, 33\,040\,809\,105\,661, 156\,151\,310\,977\,544\,887, \dots,
 \end{aligned}$$

$$\begin{aligned}
 M = 3, \quad B_3^{(3)}(n) = 1, 77, 39\,839, 62\,310\,039, 214\,107\,236\,041, 1\,358\,185\,668\,416\,501, \\
 14\,247\,249\,149\,298\,651\,007, \dots,
 \end{aligned}$$

$$\begin{aligned}
 M = 4, \quad B_3^{(4)}(n) \\
 = 1, 372, 1\,905\,633, 43\,249\,617\,004, 2\,805\,942\,285\,116\,705, 411\,223\,445\,534\,704\,016\,116, \\
 117\,428\,972\,441\,699\,060\,660\,584\,977, \dots, \tag{33}
 \end{aligned}$$

which are positive integers and as such admit combinatorial interpretation. The first two sequences in Eq. (33) may be identified as A002720 [which enumerates matching numbers of a perfect graph  $K(n,n)$ ] and A069948, respectively, in Ref. 27.

We note in passing that the numbers  $B_r^{(M)}(n)$  are solutions of the Stieltjes moment problem, i.e., they are the  $n$ th moments of positive weight functions on the positive half axis. This can be deduced from their Dobiński-type relation [Eq. (25)], whose form allows one to obtain the weight functions for any  $r$  and  $M$ . For the first two sequences in Eq. (33), the Stieltjes weights are given in Ref. 27 under their entries.

As a second illustration of our approach we shall apply it to  $D(r, 1)$ . Note that

$$[a^r(a^\dagger a)^M]^n = :B_r^{(M)}(n, a^\dagger a)a^{rn}:, \quad (34)$$

which upon using the Dobiński relation [Eq. (25)] for  $M=1$  leads to

$$e^{\lambda D(r,1)} = : \frac{1}{1 - \lambda r a^r} \exp\left(\frac{a^\dagger a}{(1 - \lambda r a^r)^{1/r}} - a^\dagger a\right) :. \quad (35)$$

The operator  $D(r, 1)$  is of Sheffer-type viewed through Hermitian conjugation (see Refs. 8 and 9), and Eq. (35) can also be obtained through the methods developed in Ref. 9 (see the Appendix). Consequently,

$$\langle z | e^{\lambda D(r,1)} | z \rangle = \frac{1}{1 - r\lambda} \exp\left(\frac{1}{(1 - r\lambda)^{1/r}} - 1\right) \quad (36)$$

$$\equiv \sum_{n=0}^{\infty} B_r^{(1)}(n) \frac{\lambda^n}{n!}, \quad (37)$$

where

$$\langle z | (a^r a^\dagger a)^n | z \rangle = B_r^{(1)}(n) = \sum_{p=1}^{n+1} |\sigma(n+1, p)| r^{n-p+1} B(p-1). \quad (38)$$

In Eq. (38)  $\sigma(n, k)$  are Stirling numbers of the first kind,  $B(n)$  are conventional Bell numbers, and in obtaining Eq. (38) we have again used Eq. (11).

Using Eqs. (19) and (29) we obtain for  $r=1$  the following formula in a compact notation



$$[D(1, M)]^n = (n!)^M : e^{-a^\dagger a} \cdot {}_M F_M(\underbrace{[n+1, \dots, n+1]}_{M \text{ times}}, \underbrace{[1, \dots, 1]}_{M \text{ times}}, a^\dagger a) a^n :. \quad (39)$$

The last formula can be used to normally order  $H(\lambda D(1, M))$  for any Taylor-expandable  $H(x)$ .


#### IV. COMBINATORICS OF NORMALLY ORDERED LAGUERRE DERIVATIVES

In Sec. III we considered the normal ordering of Laguerre derivatives for which the results heavily exploited combinatorial identities stemming from the underlying iterative character of the problem. Indeed, the reordering of the operators  $a$  and  $a^\dagger$  is a purely combinatorial task which can be interpreted in terms of graphs<sup>19,28,29</sup> and analyzed by the use of combinatorial constructors.<sup>30</sup> Briefly, to each operator in the normally ordered form  $H = \sum_{r,s} \alpha_{r,s} (a^\dagger)^r a^s$ , one associates a set of one-vertex graphs such that each vertex  $\bullet$  carries weight  $\alpha_{r,s}$  and has  $r$  outgoing and  $s$  incoming lines whose free ends are marked with white and gray spots, respectively. Multivertex graphs are built in a step-by-step manner by adding one vertex at each consecutive step and joining some of its incoming lines with some free outgoing lines of the graph constructed in the previous step. Additionally, one keeps track of the history by labeling each vertex by the number of steps in which it was introduced. As a result, one obtains a set of increasingly labeled multivertex graphs with some free incoming and outgoing lines. It can be shown that the normal ordering of powers of the operator  $H$  can be obtained by enumeration of such structures. Namely, the coefficient of  $(a^\dagger)^k a^l$  in the normally ordered form of the operator  $H^n$  is obtained by counting all possible graphs with  $n$  vertices  $\bullet$  and having  $k$  white and  $l$  gray spots, respectively. For illustration, we give two



(a)



$n=2:$



(b)



$n=2:$




FIG. 1. Building blocks (in the inset) and the associated graphs of order  $n=1,2$  for Laguerre derivatives: (a)  $H = D(1,1) = a^\dagger a^2 + a$  and (b)  $H = D(2,1) = a^\dagger a^3 + 2a^2$ .

examples of Laguerre derivatives  $D(1,1) = a a^\dagger a = a^\dagger a^2 + a$  and  $D(2,1) = a^2 a^\dagger a = a^\dagger a^3 + 2a^2$  and their graph representation leading to the solution of the normal ordering problem by simple enumeration (see Fig. 1).

One should compare these “graphical results” with the explicit formulas of Eqs. (27) and (25) or the expansion coefficients of the generating function in Eq. (36) for  $r=1,2$  and  $M=1$ . Thus, using Eq. (33), the coefficients multiplying the operators in Fig. 1(a) are the first two terms in  $B_1^{(1)}(n) = 2, 7, 34, 209$  for  $n=1, 2, \dots$  (A002720). Similar coefficients in Fig. 1(b) are the first two terms in  $B_2^{(1)}(n) = 3, 16, 121, 1179$  for  $n=1, 2, \dots$  (A121629).

## V. EXAMPLES

- (1) For  $r=M=1$ , i.e., for  $D(1,1)=aa^\dagger a$ , one obtains (see Ref. 31 for a similar calculation)

$$[D(1,1)]^n = n! :L_n(-a^\dagger a): a^n, \quad (40)$$

where  $L_n(y)$  are Laguerre polynomials and Eq. (40) is derived from Eq. (39) and using the definition of  $L_n(y)$  via the function  ${}_1F_1$ . Then

$$e^{\lambda D(1,1)} = \sum_{n=0}^{\infty} \frac{\lambda^n}{n!} [D(1,1)]^n = \sum_{n=0}^{\infty} L_n(-a^\dagger a) (\lambda a)^n =: \frac{1}{1-\lambda a} \exp\left(\frac{\lambda a^\dagger a^2}{1-\lambda a}\right):, \quad (41)$$

where in Eq. (41) we have used the ordinary generating function for the Laguerre polynomials.<sup>5</sup>

Using other generating functions listed on p.704 of Ref. 5, one can derive further formulas of type Eq. (41). [In (a) and (b) below:  $\lambda \neq 0$ ,  $p=1,2,\dots$ ]

- (a) Formula (5.11.2.6) in Ref. 5 for  $\alpha=0$  provides the normal ordering of

$${}_1F_1([b],[1],\lambda D(1,1)) =: \frac{1}{(1-\lambda a)^b} {}_1F_1\left([b],[1],\frac{\lambda a^\dagger a^2}{1-\lambda a}\right):, \quad (42)$$

which for  $b$  integer and half-integer can be written down in terms of known functions. Examples are

$${}_1F_1([3],[1],\lambda D(1,1)) =: \frac{1}{(1-\lambda a)^3} L_2\left(-\frac{\lambda a^\dagger a^2}{1-\lambda a}\right) \exp\left(\frac{\lambda a^\dagger a^2}{1-\lambda a}\right):, \quad (43)$$

$${}_1F_1\left(\left[\frac{3}{2}\right],[1],\lambda D(1,1)\right) =: \frac{1}{(1-\lambda a)^{3/2}} \exp\left(\frac{\lambda a^\dagger a^2}{2(1-\lambda a)}\right) \times \left(I_0\left(\frac{\lambda a^\dagger a^2}{2(1-\lambda a)}\right) \left(1 + \frac{\lambda a^\dagger a^2}{1-\lambda a}\right) + I_1\left(\frac{\lambda a^\dagger a^2}{2(1-\lambda a)}\right)\right):,$$

where  $I_0(y)$  and  $I_1(y)$  are the modified Bessel functions.

- (b) Similarly, we consider formula (5.11.2.8) in Ref. 5,

$$\sum_{n=0}^{\infty} \binom{n+p}{n} t^n L_{n+p}(x) = \frac{1}{(1-t)^{p+1}} e^{-[tx/(1-t)]} L_p\left(\frac{x}{1-t}\right). \quad (44)$$

Using Eq. (40) we obtain the normally ordered form of  $[\lambda D(1,1)]^p \exp(\lambda D(1,1))/p!$ ,

$$\frac{1}{p!} \sum_{n=0}^{\infty} \frac{[\lambda D(1,1)]^{n+p}}{n!} = \sum_{n=0}^{\infty} \frac{(n+p)!}{p!n!} :L_{n+p}(-a^\dagger a)(\lambda a)^{n+p}: \quad (45)$$

$$=: \frac{1}{(1-\lambda a)^{p+1}} e^{(\lambda a^\dagger a^2)/(1-\lambda a)} L_p\left(-\frac{a^\dagger a}{1-\lambda a}\right): (\lambda a)^p. \quad (46)$$

- (2) The normal order of the modified Bessel function of the first kind  $I_0(2(\lambda D(1,1))^{1/2})$  may be derived,

$$I_0(2(\lambda D(1,1))^{1/2}) = \sum_{n=0}^{\infty} \frac{L_n(-a^\dagger a)(\lambda a)^n}{n!} = :e^{\lambda a} J_0(2\sqrt{\lambda(-aa^\dagger)}) : = :e^{\lambda a} I_0(2\sqrt{\lambda a^\dagger a^2}) :, \quad (47)$$

where in the last line we have used the exponential generating function of Laguerre polynomials.<sup>5</sup> The analogous formula for  $J_0(2(\lambda D(1,1))^{1/2})$  reads as

$$J_0(2(\lambda D(1,1))^{1/2}) = \sum_{n=0}^{\infty} \frac{L_n(-a^\dagger a)(-\lambda a)^n}{n!} = :e^{-\lambda a} I_0(2\sqrt{\lambda a^\dagger a^2}) :. \quad (48)$$

- (3) We quote here the eigenfunctions of  $D_x(r, M)$  with eigenvalue 1 satisfying  $D_x(r, M)E(r, M; x) = E(r, M; x)$ , with the following  $r$  boundary conditions:

$$E(r, M; 0) = 1, \quad \left[ \frac{d^p}{dx^p} E(r, M; x) \right] \Big|_{x=0} = 0, \quad p = 1, \dots, r-1, \quad (49)$$

which are

$$E(r, M; x) = {}_0F_{M+r-1}([ ], [\underbrace{1/r, 2/r, \dots, (r-1)/r}_{r-1 \text{ times}}, \underbrace{1, \dots, 1}_{M \text{ times}}], x^r/r^{r+M}). \quad (50)$$

Useful normal ordering formulas can be obtained by applying the Dobiński relations to the eigenfunctions of  $D_x(1, M)$  with the argument taking operator values [see Eq. (50)], i.e.,  $E(1, M; D_x(1, M))$ . We briefly show the calculation, in boson notation, for  $E(1, 2; \lambda D(1, 2)) = {}_0F_2([ ], [1, 1], \lambda D(1, 2))$  [see Eq. (39)],

$$\begin{aligned} {}_0F_2([ ], [1, 1], \lambda D(1, 2)) &= \sum_{n=0}^{\infty} \frac{\lambda^n}{(n!)^3} [a(a^\dagger a)^2]^n = :e^{-a^\dagger a} \sum_{l=0}^{\infty} \frac{(a^\dagger a)^l}{l!} \sum_{n=0}^{\infty} \frac{((n+l)!)^2}{(l!)^3} (\lambda a)^n : \\ &= :e^{-a^\dagger a} \sum_{l=0}^{\infty} \frac{(a^\dagger a)^l}{l!} {}_2F_2([1+l, 1+l], [1, 1], \lambda a) :, \end{aligned} \quad (51)$$

and similarly

$$\begin{aligned} E(1, M; \lambda D(1, M)) &= {}_0F_M([ ], [\underbrace{1, 1, \dots, 1}_{M \text{ times}}], \lambda D(1, M)) \\ &= :e^{-a^\dagger a} \sum_{l=0}^{\infty} \frac{(a^\dagger a)^l}{l!} {}_M F_M([\underbrace{1+l, 1+l, \dots, 1+l}_{M \text{ times}}, \underbrace{1, 1, \dots, 1}_{M \text{ times}}], \lambda a) :, \end{aligned} \quad (52)$$

which indicates a pattern appearing in the course of this procedure.

Indeed, by evaluating the coherent state expectation value of Eq. (52) between  $\langle z=1 | \dots | z=1 \rangle$  in the spirit of Eq. (16) we obtain the hypergeometric generating functions of the numbers  $B_1^{(M)}(n)$  as then

$$e^{-1} \sum_{l=0}^{\infty} \frac{1}{l!} {}_M F_M([\underbrace{l+1, l+1, \dots, l+1}_{M \text{ times}}, \underbrace{1, 1, \dots, 1}_{M \text{ times}}], \lambda) = \sum_{n=0}^{\infty} B_1^{(M)}(n) \frac{\lambda^n}{(n!)^{M+1}}, \quad M = 1, 2, \dots \quad (53)$$

The hypergeometric generating function of  $B_r^{(M)}(n, x)$  for arbitrary  $r$  and  $M$  can also be obtained from Eq. (25) and reads

$$e^{-x} \sum_{l=0}^{\infty} \frac{x^l}{l!} {}_M F_M \left( \underbrace{[l/r+1, l/r+1, \dots, l/r+1]}_{M \text{ times}}, \underbrace{[1, 1, \dots, 1]}_{M \text{ times}}, r^M \lambda \right) = \sum_{n=0}^{\infty} B_r^{(M)}(n, x) \frac{\lambda^n}{(n!)^{M+1}}. \quad (54)$$

In spite of their apparent complexity the left hand side of the above equations can be straightforwardly handled by computer algebra systems.<sup>32</sup>

## VI. CONCLUSIONS AND OUTLOOK

We have found exact analytical expressions for the generalized Stirling numbers and generalized Bell polynomials, which appear in the normal ordering of powers of Laguerre-type derivative operators, and have provided a complete set of hypergeometric generating functions for these quantities. The combinatorial aspect of the problem was demonstrated by finding an exact mapping between the normal ordering and an enumeration of increasingly labeled, multivertex forests constructed according to a two-parameter  $(r, M)$  prescription. In this way, analytical, numerical, and combinatorial facets of this problem have been given a very complete treatment. We have also used generalized Dobiński relations to investigate the properties of these Laguerre-type differential operators. We provided a large number of operational formulas involving functions of Laguerre derivatives, which can alternatively be applied using the boson language. The framework developed above enables one to construct and analyze new coherent states relevant to nonlinear quantum optics, which will be the subject of forthcoming research.

Note that we have used the operators  $a$  and  $a^\dagger$  since this is the usual convention for physics. In our pure algebraic setting it has no particular significance, however, and one might as well use the operators  $a$  and  $b$  for  $b \neq a^\dagger$  [see Refs. 33 and 34 for many relevant algebraic identities which follow from  $ab - qba = p$  and which apply to our case also].

## ACKNOWLEDGMENTS

We wish to acknowledge support from the Agence Nationale de la Recherche (Paris, France) under Program No. ANR-08-BLAN-0243-2. Two of us (P.B. and A.H.) wish to acknowledge support from the Polish Ministry of Science and Higher Education under Grant Nos. N202 061434 and N202 107 32/2832 as well as kind hospitality of Laboratoire d'Informatique d'Université Paris-Nord, where part of this research was carried out.

## APPENDIX: SHEFFER-TYPE OPERATORS

We derive Eq. (35) with the help of methods developed in Refs. 8 and 9. First, observe that  $D(r, 1) = a^\dagger a^{r+1} + r a^r$  from which it follows that  $D^\dagger(r, 1) = (a^\dagger)^{r+1} a + r (a^\dagger)^r$  is an operator of Sheffer type:  $D^\dagger(r, 1) = v(a^\dagger) + q(a^\dagger)a$  with  $q(x) = x^{r+1}$  and  $v(x) = r x^r$ . The normally ordered form of  $\exp(\lambda D^\dagger(r, 1))$  is obtained by solving the linear differential equations [Eqs. (2) and (3) in Ref. 9] for  $T(\lambda, x)$  and  $g(\lambda, x)$  yielding

$$T(\lambda, a^\dagger) = \frac{a^\dagger}{(1 - \lambda r (a^\dagger)^r)^{1/r}} \quad (A1)$$

and

$$g(\lambda, a^\dagger) = \frac{1}{1 - \lambda r (a^\dagger)^r}. \quad (A2)$$

According to Eq. (29) in Ref. 9, the normally ordered form of  $e^{\lambda D(r, 1)}$  is

$$e^{\lambda D(r, 1)} = [e^{\lambda D^\dagger(r, 1)}]^\dagger = :g(\lambda, a) e^{a^\dagger(T(\lambda, a) - a)}:, \quad (A3)$$

which gives Eq. (35).

<sup>1</sup>G. Dattoli and P. Ricci, *Georgian Math. J.* **10**, 481 (2003).

- <sup>2</sup>G. Dattoli, P. E. Ricci, and I. Khomasuridze, *Integral Transforms Spec. Funct.* **15**, 309 (2004).
- <sup>3</sup>G. Dattoli, M. R. Martinelli, and P. E. Ricci, *Integral Transforms Spec. Funct.* **16**, 661 (2005).
- <sup>4</sup>N. Fleury and A. V. Turbiner, *J. Math. Phys.* **35**, 6144 (1994).
- <sup>5</sup>A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev, *Special Functions, Integrals and Series Vol. 2* (Gordon and Breach, New York, 1998).
- <sup>6</sup>G. Dattoli, A. M. Mancho, M. Quatromini, and A. Torre, *Radiat. Phys. Chem.* **61**, 99 (2001).
- <sup>7</sup>We use a convenient and self-explanatory notation for the hypergeometric functions of type  ${}_pF_q: {}_pF_q(\text{[List of } p \text{ upper parameters]}, \text{[List of } q \text{ lower parameters]}, x)$ .
- <sup>8</sup>G. Dattoli, P. L. Ottaviani, A. Torre, and L. Vásquez, *Riv. Nuovo Cim.* **20**, 3 (1997).
- <sup>9</sup>P. Blasiak, A. Horzela, K. A. Penson, G. H. E. Duchamp, and A. I. Solomon, *Phys. Lett. A* **338**, 108 (2005).
- <sup>10</sup>J. R. Klauder and E. C. G. Sudarshan, *Fundamentals of Quantum Optics* (Benjamin, New York, 1968); J. R. Klauder and B.-S. Skagerstam, *Coherent States: Application in Physics and Mathematical Physics* (World Scientific, Singapore, 1985).
- <sup>11</sup>R. L. de Matos Filho and W. Vogel, *Phys. Rev. A* **54**, 4560 (1996).
- <sup>12</sup>M. Rasetti, J. Katriel, A. I. Solomon, and G. D'Ariano, in *Squeezed and Nonclassical Light*, edited by P. Tombesi and E. P. Pike (Plenum, New York, 1989), p. 301.
- <sup>13</sup>A. I. Solomon, *Phys. Lett. A* **188**, 215 (1994).
- <sup>14</sup>L. Comtet, *Advanced Combinatorics* (Reidel, Dordrecht, 1974).
- <sup>15</sup>E. W. Weisstein, Stirling number of the first kind, from MathWorld, a Wolfram Web Resource (<http://mathworld.wolfram.com/StirlingNumberoftheFirstKind.html>).
- <sup>16</sup>P. Blasiak, K. A. Penson, and A. I. Solomon, *Phys. Lett. A* **309**, 198 (2003).
- <sup>17</sup>P. Blasiak, K. A. Penson, and A. I. Solomon, *Ann. Comb.* **7**, 127 (2003).
- <sup>18</sup>P. Blasiak, K. A. Penson, A. I. Solomon, A. Horzela, and G. H. E. Duchamp, *J. Math. Phys.* **46**, 052110 (2005).
- <sup>19</sup>M. A. Méndez, P. Blasiak, and K. A. Penson, *J. Math. Phys.* **46**, 083511 (2005).
- <sup>20</sup>W. H. Louisell, *Quantum Statistical Properties of Radiation* (Wiley, New York, 1990).
- <sup>21</sup>P. Blasiak, A. Horzela, K. A. Penson, A. I. Solomon, and G. H. E. Duchamp, *Am. J. Phys.* **75**, 639 (2007) (this pedagogical paper contains exhaustive list of references dealing with the boson normal ordering problem).
- <sup>22</sup>J. Katriel, *Lett. Nuovo Cim.* **10**, 565 (1974).
- <sup>23</sup>K. E. Cahill and R. J. Glauber, *Phys. Rev.* **177**, 1857 (1969) (the formula is attributed to J. Schwinger).
- <sup>24</sup>V. V. Mikhailov, *J. Phys. A* **16**, 3817 (1983); J. Katriel, *ibid.* **16**, 4171 (1983).
- <sup>25</sup>P. Blasiak, K. A. Penson, and A. I. Solomon, *J. Phys. A* **36**, L273 (2003).
- <sup>26</sup>P. Blasiak, A. Horzela, K. A. Penson, and A. I. Solomon, *J. Phys. A* **39**, 4999 (2006).
- <sup>27</sup>N. J. A. Sloane, Encyclopedia of Integer Sequences, <http://www.research.att.com/~njas/sequences> (2009).
- <sup>28</sup>P. Blasiak and A. Horzela, e-print arXiv:0710.0266.
- <sup>29</sup>P. Blasiak, A. Horzela, K. A. Penson, A. I. Solomon, and G. H. E. Duchamp, *J. Phys. A: Math. Theor.* **41**, 415204 (2008).
- <sup>30</sup>P. Flajolet and R. Sedgewick, *Analytic Combinatorics* (Cambridge University Press, Cambridge, 2008).
- <sup>31</sup>J. Riordan, *Combinatorial Identities* (Wiley, New York, 1968).
- <sup>32</sup>We have made extensive use of MAPLE in this paper.
- <sup>33</sup>A. V. Turbiner, *Funct. Anal. Appl.* **29**, 291 (1995).
- <sup>34</sup>L. Hellstroem and S. D. Silvestrov, *Commuting Elements in  $q$ -Deformed Heisenberg Algebras* (World Scientific, Singapore, 2000).