

# The VLT-FLAMES Tarantula Survey

C. J. Evans<sup>1</sup>, N. Bastian, Y. Beletsky, I. Brott, M. Cantiello,  
J. S. Clark, P. A. Crowther, A. de Koter, S. de Mink, P. L. Dufton,  
P. Dunstall, M. Gieles, G. Gräfener, V. Hénault-Brunet, A. Herrero,  
I. D. Howarth, N. Langer, D. J. Lennon, J. Maíz Apellániz,  
N. Markova, F. Najarro, J. Puls, H. Sana, S. Simón-Díaz,  
S. J. Smartt, V. E. Stroud, W. D. Taylor, C. Trundle,  
J. Th. van Loon, J. S. Vink and N. R. Walborn

<sup>1</sup>UK ATC, Royal Observatory Edinburgh, Blackford Hill, Edinburgh, EH9 3HJ, UK  
email@ [chris.evans@stfc.ac.uk](mailto:chris.evans@stfc.ac.uk)

**Abstract.** The Tarantula Survey is an ambitious ESO Large Programme that has obtained multi-epoch spectroscopy of over 1,000 massive stars in the 30 Doradus region of the Large Magellanic Cloud. Here we introduce the scientific motivations of the survey and give an overview of the observational sample. Ultimately, quantitative analysis of every star, paying particular attention to the effects of rotational mixing and binarity, will be used to address fundamental questions in both stellar and cluster evolution.

**Keywords.** stars: early-type – stars: fundamental parameters – binaries: spectroscopic – open clusters and associations: individual: 30 Doradus

---

## 1. Introduction

The Tarantula Nebula (30 Doradus, NGC 2070) in the Large Magellanic Cloud (LMC) is the brightest and most massive H II region in the Local Group. It is a beautiful and very intricate region, far removed from a ‘simple single-stellar-population’. Indeed, Walborn & Blades (1997) identified at least five distinct populations (cf. Walborn, 2009):

- The central ‘Carina Phase’ concentration, rich in early O-type stars and including the dense cluster R136.
- A younger, likely triggered, ‘Orion Phase’ to the north and west of R136.
- A ‘Sco OB1 Phase’ of early-type supergiants throughout the central field.
- An older ‘h &  $\chi$  Persei Phase’ in Hodge 301, containing cooler, more evolved, supergiants, to the northwest of the centre.
- A separate ‘Sco OB1 Phase’ surrounding the luminous blue variable R143.

With its rich stellar populations, 30 Dor is the ideal laboratory in which to investigate a number of important outstanding questions regarding the physics, evolution, binary fraction, and chemical enrichment of the most massive stars. Building on the successes of the VLT-FLAMES Survey of Massive Stars (Evans et al., 2005), here we introduce a new multi-epoch spectral survey of over 1,000 massive stars in the 30 Dor region.

In the broader context, 30 Dor is at the northern end of a large column of molecular gas which extends south for over 2,000 pc (Cohen et al., 1988, Fukui et al., 2008). N-body models examining the recent edge-on motion of the LMC through the halo of the Milky Way suggest significant star formation in the eastern part of the LMC, as manifested by 30 Dor, due to ram pressure (Mastropietro, Burkert & Moore, 2009). With the reservoir of gas to the south, the region seems destined to become an even more spectacular star-formation complex over the next few million years.

## 2. Multiplicity in Massive Stars

The effects of binarity/multiplicity on the formation and subsequent evolution of high-mass stars is a vibrant area of research. Indeed, one of the key ingredients missing from current theories of both star formation and cluster evolution is a robust binary fraction of massive stars, and the distribution of the mass ratios in these systems. Some motivation in this direction was provided by Zinnecker & Yorke (2007):

‘The future of spectroscopic massive binary research lies in the near-IR and in multi-epoch radial velocity surveys of embedded massive stars’

These words were primarily concerned with the earliest stages of star formation but they coincide with growing interest in multi-epoch spectroscopic studies in open clusters, aimed at identification and characterisation of their binary populations (Table 1). The most pertinent of these is the study of 50 early-type stars in 30 Dor by Bosch, Terlevich & Terlevich (2009). From Gemini spectroscopy at seven epochs they found a binary fraction of  $\geq 50\%$ , noting that the data were not inconsistent with it being 100%. Recent multi-epoch AO-corrected SINFONI observations found (tentative) evidence for a short-period companion in only one of the six central WR stars at the core of R136 (Schnurr et al., 2009), but it is clear that there is a very rich binary population in 30 Dor.

**Table 1.** Selected multi-epoch spectroscopic surveys in open clusters.

Cluster	Binary fraction	Reference
IC 1805	$\geq 0.20$	De Becker et al. (2006)
NGC 6231	$\geq 0.63$	Sana et al. (2008)
NGC 6611	$\geq 0.44$	Sana, Gosset & Evans (2009)
NGC 2244	$\geq 0.17$	Mahy et al. (2009)
30 Dor	$\geq 0.50$	Bosch, Terlevich & Terlevich (2009)

One of the serendipitous aspects of the FLAMES Survey of Massive Stars was the large number of spectroscopic binaries discovered (Table 2). The time sampling of the service-mode observations did a reasonable (but not thorough) job of binary detection, with lower limits to the binary fraction of  $\sim 30\%$  in three of the target clusters. Adopting the same methods as Sana, Gosset & Evans (2009), we have calculated the detection probabilities for short, intermediate and long period binaries for each cluster field. The aggregated detection probabilities (for systems with periods of two days to ten years) are given in the final column of Table 2. The similarity in the detection probabilities suggests that the lower fraction found in NGC 330 is genuinely different to the others. While it is unfair to compare the NGC 330 observations with the rich, younger cluster fields of NGC 346 and N11, NGC 2004 is its LMC cousin; this difference in the binary fraction is intriguing and the subject of ongoing work.

**Table 2.** Spectroscopic binaries from Evans et al. (2006).

Cluster	Galaxy	# O+Early B	# Binary	Binary fraction	Detection prob. [2d-10yr]
NGC 346	SMC	103	27	$\geq 26\%$	0.66
NGC 330	SMC	104	4	$\geq 4\%$	0.71
NGC 2004	LMC	105	24	$\geq 23\%$	0.64
N11	LMC	120	43	$\geq 36\%$	0.64

The relationship of the binary fraction with density and the spatial extent of a cluster is still unclear (e.g. Mahy et al., 2009), while the binary fraction in OB associations is often similar to clusters, but with fewer short-period systems (Zinnecker & Yorke, 2007).

### 3. The Tarantula Survey

The new survey comprises 160 hrs of VLT-FLAMES spectroscopy in the 30 Dor region (PI: Evans). Most of the observations (142 hrs) have now been completed, with the remainder scheduled for the coming semester.

One of the prime drivers for this survey was the issue of binarity, shaping the multi-epoch observational strategy. It is clear that identification of binaries, and the mass ratios in those systems, is an important empirical result for N-body models of star and cluster formation/evolution. Moreover, in many clusters, e.g. NGC 6231 (Sana et al., 2008), the majority of O-type stars are members of a binary system. Thus, to gain a true understanding of the upper H-R diagram, the effects of binarity need to be fully included in theoretical models of stellar evolution (e.g. de Mink et al., 2009). Although we have focussed on this aspect for this symposium, the genesis of the survey arose from a much broader range of other scientific motivations, including:

- The role of stellar rotation in the chemical enrichment and evolution of massive stars. Hunter et al. (2008) have revealed new challenges for theory in B-type stars; we seek to investigate these effects in the more dominant, massive O-type stars.
- Determination of the rotational velocity distribution in 30 Dor. Are there sufficient high-mass, rapidly-rotating stars to provide a channel for long-duration  $\gamma$ -ray bursts (cf. Yoon, Langer & Norman, 2006)?
- Armed with precise radial velocities and identification of binaries, do we see kinematic evidence of mass segregation and/or infant mortality in and around R136?
- A more holistic objective of a near-complete census of the closest ‘proto-starburst’, with applications in the context of population synthesis methods and interpretation of spectra of unresolved massive stars clusters at Mpc distances.


### 4. GIRAFFE Observations

The primary dataset comprises spectroscopy of 1,000 stars using the GIRAFFE spectrograph, which is fed by 132 MEDUSA fibres available for science (or sky) observations across a 25' field (Pasquini et al., 2002). Targets were selected from unpublished imaging with the Wide-Field Imager (WFI) on the ESO/MPG 2.2-m telescope, and from Brian Skiff’s reworking of the Selman et al. (1999) photometric catalogue in the central 90". To obtain a representative sample of the upper part of the HR diagram, including evolved luminous stars, no colour cut was applied to potential targets but a faint cut-off ( $V < 17$ ) was enforced to ensure sufficient signal-to-noise for each star.

Nine MEDUSA configurations were observed, each of which was observed at three wavelength settings (see Table 3). This yields full coverage of the classical blue-optical region used for spectroscopic classification and analysis, combined with higher resolution spectroscopy of the H $\alpha$  region to enable determination of the stellar wind intensity. From inspection of initial reductions, the minimum signal-to-noise in the stacked spectra for the faintest stars is  $\sim 50$ , i.e. the spectra are suitable for quantitative analysis as well as radial velocity monitoring.

**Table 3.** Summary of FLAMES-GIRAFFE observations.

GIRAFFE setting	$\lambda$ -coverage ( $\text{\AA}$ )	$R$	Exposures
LR02	3980-4535	6,500	6 $\times$ (2 $\times$ 1815s)
LR03	4505-5050	7,500	3 $\times$ (2 $\times$ 1815s)
HR15N	6470-6790	17,000	2 $\times$ (2 $\times$ 2265s)



**Figure 1.**  $14' \times 14'$  V-band WFI image showing the FLAMES-GIRAFFE targets in and around 30 Dor (north to the top, east to the left).

The distribution of the majority of the MEDUSA targets is shown in Figure 1. The survey samples the full extent of 30 Dor and outwards into the ‘field’ population and other nearby OB associations to fully exploit the FLAMES field-of-view and spare fibres, thus bolstering the observational sample.

#### 4.1. Preliminary Classification


Work is now progressing in earnest with the final science reductions of the GIRAFFE spectra. To characterise the spectral content of the survey, we extracted the reduced spectra from one pair of LR02 observations for each MEDUSA configuration. In advance of the full reductions, it was not possible to classify  $\sim 150$  stars from just one observation (although most are likely B-type stars) but from visual inspection of the spectra the sample contains:

- In excess of 300 O-type stars and  $\sim 20$  Wolf-Rayet/‘slash’ stars. This is a hugely significant improvement in terms of sampling the upper HR diagram, e.g., compared to the analysis of 28 O-type stars in the LMC by Mokiem et al. (2007). Each star will be studied for binary companions, and then analysed to obtain physical and stellar wind parameters, including the first large-scale study of nitrogen enrichment in O-type stars.

- Over 400 B-type spectra, which will be used to establish the baseline chemical abundances in 30 Dor and, with such a large sample in one field, will be used to revisit the role of rotationally-induced mixing on surface nitrogen enrichment (cf. Hunter et al., 2008).
- $\sim 150$  cooler stars with spectral types of A, F and later. Some will be foreground objects to be discarded, but the majority will be evolved, luminous stars which will be used to investigate the short lifetimes of these evolutionary phases via population synthesis models.

#### 4.2. Binary Detection Probabilities

Using the methods from Sana, Gosset & Evans (2009) we have calculated the detection probabilities for binaries from the actual time sampling of the nine observed MEDUSA configurations. In these calculations we assume a  $\Delta v_r$  threshold of  $20 \text{ km s}^{-1}$ , requiring a radial velocity precision of  $\sim 5 \text{ km s}^{-1}$  (which should be achievable at the resolving power of the new spectroscopy for all but the fastest rotating stars). The detection probabilities, as a function of orbital period, for the first MEDUSA configuration are shown by the black line in Figure 2. We are relatively complete up to periods of a few 10s of days, with a steep fall-off beyond 100 days. The inclusion of one additional epoch in the coming observing season significantly helps with the detection of both intermediate and long period binaries (red/grey line). By quantifying our detection probabilities using such simulations, we will be able to put firm limits on the observed binary fraction.



**Figure 2.** Detection probability of binary companions for one of the MEDUSA configurations. The black line shows the detection results for the five epochs already executed; the red (grey) line illustrates the increased probability obtained from a sixth epoch, scheduled for observation in the coming semester, i.e. separated by approximately one year from the other epochs.

## 5. Supplementary Data

### 5.1. ARGUS & UVES Observations of R136

R136 is too dense for effective use of the MEDUSA fibres, so a  $15''$  exclusion radius around the core was employed in the fibre allocations. To investigate the dynamics and binarity of stars in and around R136, as part of the Large Programme we have observed five pointings with the ARGUS integral field unit (IFU) which delivers a  $12'' \times 7''$  field-of-

view. Each pointing has been observed with the LR02 GIRAFFE setting (which delivers a resolving power of  $\sim 10,000$  in the IFU mode) at five epochs.

In parallel to the ARGUS observations, we used the fibre-feed to the red arm of UVES to observe 25 stars that were not included in the MEDUSA configurations. The  $\lambda 5200$  standard set-up was used, delivering spectral coverage of  $\sim \lambda\lambda 4200-6200$  at  $R = 47,000$ .

### 5.2. VLT-SINFONI K-band Spectroscopy

The majority of the known WR and extreme O-type emission-line stars in 30 Dor are in the central regions. Near-IR IFU observations with SINFONI (12hrs; PI: Gräferer) will be used to obtain K-band spectroscopy of the central arcminute around R136. The stellar wind lines in the K-band, principally from Brackett  $\gamma$  and He II, are more sensitive than the optical lines at low mass-loss rates, enabling a more precise determination of the physical parameters of the most extreme stars.

### 5.3. Faulkes Photometric Follow-up

In the longer term, the spectroscopy in the central  $\sim 10$  arcminutes will be supplemented with multi-band photometric monitoring from the Faulkes Telescope South, as part of their schools education programme. Faulkes has a  $4.7 \times 4.7$  field-of-view, so the main body of 30 Dor will be mapped with several pointings, delivering multi-epoch photometry that will, for example, assist with the analysis of identified binary systems.

## 6. Summary

We have an exceptional and unique data resource available to us to investigate the massive-star population in 30 Dor, now the hard work begins!

**Acknowledgements:** Based on observations from ESO programme 182.D-0222. We are indebted to Brian Skiff for his careful reworking of the Selman et al. astrometry.

## References

- Bosch, G., Terlevich, E. & Terlevich, R., 2009, *AJ*, 137, 3437  
 Cohen, R. S. et al., 1988, *ApJ*, 331, L95  
 De Becker, M., Rauw, G., Manfroid, J. & Eenens, P., 2006, *A&A*, 456, 1121  
 de Mink, S. E. et al., 2009, *A&A*, 497, 243  
 Evans, C. J. et al., 2005, *A&A*, 437, 467  
 Evans, C. J., Lennon, D. J., Smartt, S. J. & Trundle, C. 2006, *A&A*, 456, 623  
 Fukui, Y. et al., 2008, *ApJS*, 178, 56  
 García, B. & Mermilliod, J. C., 2001, *A&A*, 368, 122  
 Hunter, I. et al., 2008, *ApJ*, 676, L29  
 Mahy, L. et al., 2009, *A&A*, 502, 937  
 Mastroiello, C., Burkert, A. & Moore, B. 2009, *MNRAS*, arXiv:0905.1126  
 Mokiem, M. R. et al., 2007, *A&A*, 465, 1003  
 Pasquini, L. et al., 2002, *Msngr*, 110, 1  
 Sana, H. et al., 2008, *MNRAS*, 386, 447  
 Sana, H., Gosset, E. & Evans, C. J., 2009, *MNRAS*, arXiv:0909.0486  
 Schnurr, O. et al., 2009, *MNRAS*, 397, 2049  
 Selman, F., Melnick, J., Bosch, G. & Terlevich, R., 1999, *A&AS*, 341, 98  
 Walborn, N. R. & Blades, J. C., 1997, *ApJS*, 112, 457  
 Walborn, N. R. 2009, to appear in: Leitherer et al. (eds.), *Hot & Cool: Bridging Gaps in Massive Star Evolution*, ASP Conference Series.  
 Yoon, S.-C., Langer, N. & Norman, C., 2006, *A&A*, 460, 199  
 Zinnecker, H. & Yorke, H. W., 2007, *ARA&A*, 45, 481