

The Open University's repository of research publications and other research outputs

The submillimetre wavelength spectrum of Orion-A

Conference or Workshop Item

How to cite:

White, Glenn J.; Monteiro, Tania; Rainey, Ruth; Richardson, Kevin; Griffin, Matthew and Avery, L. (1987). The submillimetre wavelength spectrum of Orion-A. In: IAU Symposium No 115 - Star Forming Regions, 11-15 Nov 1985, Tokyo, Japan, p. 153.

For guidance on citations see [FAQs](#).

© 1987 International Astronomical Union

Version: Version of Record

Link(s) to article on publisher's website:

http://dx.doi.org/doi:10.1007/978-94-009-4782-5_46

<http://adsabs.harvard.edu/abs/1987IAUS..115..153W>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.




Fig. 3. A schematic representation to show the inhomogeneous or clumpy structure of the bright bar. Dashed lines represent shock fronts.

THE SUBMILLIMETRE WAVELENGTH SPECTRUM OF ORION A

Glenn J. White¹, Tania Monteiro², Ruth Rainey¹, Kevin Richardson¹, Matthew Griffin¹, L. Avery³

¹Queen Mary College, University of London, England.

²University of Newcastle-upon-Tyne, England.

³National Research Council, Ottawa, Canada.

We report on the first submillimetre wavelength spectral scan of the Orion A molecular cloud in the frequency range 342-463 GHz (0.88-0.65 mm) using the Queen Mary College Submillimetre Heterodyne Receiver at UKIRT. Twenty-eight molecular transitions were detected, the majority of these for the first time. The lines include transitions of CO, CS, HCN, HCO⁺, H₂CO, H₂CS, SO, SO₂, CCH, SiO and CH₃OH. Upper limits are reported for a number of lines including CO⁺ and the ground state transition of NH₂. A number of the lines are surprisingly intense, and we will present maps of the relative distributions of HCO⁺, HCN, H₂CO and CCH, which show striking differences in their spatial structures. We will present details of the excitation of a number of the lines based on the results from this survey.