

# Open Research Online

---

The Open University's repository of research publications and other research outputs

## Differing effects of NT-3 and GDNF on dissociated enteric ganglion cells exposed to hydrogen peroxide in vitro

### Journal Item

How to cite:

Korsak, Kris; Silva, Ayona T. and Saffrey, M. Jill (2012). Differing effects of NT-3 and GDNF on dissociated enteric ganglion cells exposed to hydrogen peroxide in vitro. *Neuroscience Letters*, 517(2) pp. 102–106.

For guidance on citations see [FAQs](#).

© 2012 Elsevier Ireland Ltd.

Version: Accepted Manuscript

Link(s) to article on publisher's website:  
<http://dx.doi.org/doi:10.1016/j.neulet.2012.04.036>

---

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

---

[oro.open.ac.uk](http://oro.open.ac.uk)

## Accepted Manuscript

Title: Differing effects of NT-3 and GDNF on dissociated enteric ganglion cells exposed to hydrogen peroxide *in vitro*

Authors: Kris Korsak, Ayona T. Silva, M. Jill Saffrey


PII: S0304-3940(12)00556-3  
DOI: doi:10.1016/j.neulet.2012.04.036  
Reference: NSL 28897

To appear in: *Neuroscience Letters*

Received date: 22-2-2012  
Revised date: 26-3-2012  
Accepted date: 13-4-2012

Please cite this article as: K. Korsak, A.T. Silva, M.J. Saffrey, Differing effects of NT-3 and GDNF on dissociated enteric ganglion cells exposed to hydrogen peroxide *in vitro*, *Neuroscience Letters* (2012), doi:10.1016/j.neulet.2012.04.036

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Differing effects of NT-3 and GDNF on dissociated enteric ganglion cells exposed to hydrogen peroxide *in vitro*

Kris Korsak<sup>a</sup>, Ayona T. Silva<sup>b</sup>, M. Jill Saffrey<sup>a\*</sup>

<sup>a</sup>Department of Life, Health and Chemical Sciences, The Open University, UK

<sup>b</sup>The Royal Veterinary College, University of London, UK

Department of Life, Health and Chemical Sciences, The Open University, Milton Keynes, MK7 6AA, United Kingdom. Tel.: +0044 1908 54167

[j.saffrey@open.ac.uk](mailto:j.saffrey@open.ac.uk)

Highlights

Oxidative stress has been implicated in aging and disease-related damage of enteric neurons

Neurotrophic factors have been reported to have protective effects against oxidative stress

We assessed protective effects of NT-3 and GDNF in dissociated cultures of enteric ganglia

NT-3, but not GDNF protected against hydrogen peroxide-induced toxicity on enteric neurons

**Abstract**

Oxidative stress is widely recognized to contribute to neuronal death during various pathological conditions and aging. In the enteric nervous system (ENS), reactive oxygen species have been implicated in the mechanism of age-associated neuronal loss. The neurotrophic factors neurotrophin 3 (NT-3) and glial cell line-derived neurotrophic factor (GDNF) are important in the

development of enteric neurons and continue to be expressed in the gut throughout life. It has therefore been suggested that they may have a neuroprotective role in the ENS. We investigated the potential of NT-3 and GDNF to prevent death of enteric ganglion cells in dissociated cell culture after exposure to hydrogen peroxide ( $H_2O_2$ ).  $H_2O_2$  treatment resulted in a dose-dependent death of enteric neurons and glial cells, as demonstrated by MTS assay, Bis benzimide and propidium iodide staining and immunolabelling. Cultures treated with NT-3 prior to exposure showed reduced cell death compared to untreated control or GDNF-treated cultures. GDNF treatment did not affect neuronal survival in  $H_2O_2$ -treated cultures. These results suggest that NT-3 is able to enhance the survival of enteric ganglion cells exposed to oxidative stress.

*Keywords:* Neurotrophin-3, Glial cell line derived neurotrophic factor, enteric ganglia, oxidative stress, hydrogen peroxide

## Introduction

The neurons of the enteric nervous system (ENS), located in the wall of the digestive tract, regulate intestinal functions, such as motility and secretion [4,6,13]. Neurodegenerative changes, including neuronal loss, have been described in the ENS during aging [7,27,30], and may contribute to gastrointestinal dysfunction, such as constipation and incontinence, which increase in incidence in the elderly [e.g.3]. The mechanisms underlying age-related enteric neuronal loss, however, are not understood, but there is evidence that reactive oxygen species (ROS) are elevated in myenteric neurons in old rats [37]. It has also been reported that myenteric neuronal loss is reduced in calorically-restricted rats[10,37].

Neurotrophic factors have been reported to protect neurons from oxidative stress [e.g.12,26,37,see 16,25]. It has therefore been suggested that increased survival of myenteric neurons in calorically-restricted animals might be due to the actions of neurotrophic factors present in the gut [37]. Two such factors, which continue to be expressed in the adult gut are neurotrophin 3 (NT-3) and glial cell line-derived neurotrophic factor (GDNF)[5,32]. Treatment of segments of intestinal smooth muscle (muscularis externa, in which myenteric ganglia are embedded) from calorically-restricted rats with NT-3 and GDNF reduced neuronal ROS levels and also enhanced resistance to menadione-induced apoptosis [37].

Here we examined the possible protective effects of NT-3 and GDNF in the ENS further, using a culture model of dissociated myenteric ganglion cells.

Possible protective effects of NT-3 and GDNF were examined under conditions of oxidative stress induced by hydrogen peroxide ( $H_2O_2$ ), which is an established model [14,17,24] causing oxidative damage to cells *in vivo* [29]. Exogenous  $H_2O_2$  readily enters cells [18] and induces apoptosis in many cell types [9].  $H_2O_2$  has also recently been shown to reduce numbers of vulnerable enteric neurons in an organotypic culture system [38]. Here we describe the effects of  $H_2O_2$  treatment, in the presence and absence of NT-3 or GDNF, on dissociated myenteric ganglion cells from rat ileum.

## 2. Materials and Methods

### 2.1 Primary cultures of enteric ganglion cells and factor treatment

Dissociated cultures of isolated myenteric ganglia were used, to allow equivalent access of reagents to individual cells, and facilitate discrimination between individual cells when counting. Segments of myenteric ganglia were separated from muscularis externa of 7-day-old Sprague-Dawley rat ileum after incubation in collagenase type II (1mg/ml in HBSS containing 10 $\mu$ g/ml DNase) at 37°C. Ganglia were dissociated after 15 minutes incubation in trypsin-EDTA (Sigma) and passage through a 25 gauge needle. Cells were seeded in 150 $\mu$ l 199 medium with N1 supplements (199N1) containing 10% fetal calf serum (Sigma) at 2X10<sup>4</sup> cells per 13mm glass coverslip coated with poly-L-lysine. After 1 hour incubation at 37°C, 2.5% CO<sub>2</sub>, 850 $\mu$ l 199N1 medium was added. 16 hours later medium was replaced with serum-free 199N1. Cells were supplemented with desired concentrations of factors 24 hours after plating.

### *2.2 Hydrogen peroxide exposure*

Dilutions of H<sub>2</sub>O<sub>2</sub> (Sigma) were made fresh from 30% stock solution into HBSS for each experiment and was used at 1, 5, 10 and 25μM. Cultures of enteric ganglion cells grown with NT-3 or GDNF (10ng/ml) were exposed to H<sub>2</sub>O<sub>2</sub> and subsequently incubated prior to the assay for 4 hours (for bis-benzimide/propidium iodide staining) or 6 hours (for MTS assay) at 37°C, 2.5% CO<sub>2</sub>.

### *2.3 PGP 9.5 immunolabelling*

Cultures were washed with phosphate buffered saline (PBS, pH7.3) and fixed with 4% paraformaldehyde for 1 hour at room temperature. Fixed cultures were incubated with primary rabbit anti-PGP 9.5 antibody (1:30000 in antibody diluting solution (ABDS:PBS, lysine, 0.1%sodium azide, 0.1%BSA, 1% Triton X-100) for 2 hours at room temperature, washed with PBS and incubated with fluorescein-conjugated goat anti-rabbit antibody (1:100 in ABDS) for one hour at room temperature. Cultures were washed with PBS and mounted on glass slides in Citifluor mountant.

### *2.4 Bis-benzimide/propidium iodide staining*

Culture medium was replaced with 1ml of fresh medium. 10μl bis-benzimide (Hoechst stain, stock 500μg/ml in PBS) was added to each coverslip. After 20 minutes incubation at 37°C 10μl of propidium iodide (PI) was added and cultures were incubated at room temperature for 5 minutes. Subsequently all wells were washed two times with HBSS without phenol red (Sigma), fixed for 1 hour in 4%

glutaraldehyde, washed in PBS and mounted on glass slides in Citifluor mountant.

### *2.5 Quantification of cell numbers*

Cells stained with PGP9.5 or bis-benzimide and PI were counted under 400X magnification using either Zeiss Axiophot or Nikon Eclipse EB800 microscope. Bis-benzimide and PI-stained cells were counted in five random fields of view on each coverslip; PGP 9.5 stained cells were counted in a strip across the diameter of the coverslip. Data were obtained from three experiments and pooled for subsequent statistical analysis.

### *2.6 Viability assay*

Viability of the cells exposed to H<sub>2</sub>O<sub>2</sub> was assayed using CellTiter96<sup>R</sup> kit (Promega), which allows colorimetric estimation of the number of viable cells. Briefly, growth medium of cultures exposed to H<sub>2</sub>O<sub>2</sub> was replaced with HBSS without phenol red (Sigma), and MTS reagent was added to the cultures. After 4 hours incubation at 37°C absorbance was measured at 492nm.

### *2.7 Statistical analysis*

Data were analysed by one-way analysis of variance (ANOVA) followed by Tukey's post-hoc test.

## **3. Results**

### *3.1 Effects of NT-3 and GDNF on cultured enteric ganglion cells*

Preliminary experiments were performed to assess the effects of NT-3 and GDNF in our culture model and to examine the extent of cell death occurring naturally during culture. Cultures containing enteric neurons and glial cells were

grown with 10ng/ml NT-3 or GDNF for 12 and 36 hours. Immunolabelling of cultures containing both neurons and glial cells with the neuronal marker PGP9.5 allows clear distinction between the two cell types, enabling accurate cell counts. No significant change in neuronal number was measured after 12 hours treatment with either NT-3 or GDNF (Figure 1). Glial cell numbers were also similar in control, NT-3- and GDNF-treated cultures after 12 hours (data not shown). After 36 hours incubation, control cultures showed a significant ( $p < 0.01$ ) decline in neuronal numbers. In contrast, cultures grown with NT-3 exhibited reduced neuronal death compared to controls ( $p < 0.01$ ). Interestingly, GDNF treatment stimulated a marked increase ( $p < 0.01$ ) of neuronal numbers, which exceeded those found in cultures treated with GDNF for 12 hours. Glial cell numbers were affected by GDNF in a manner similar to neurons, whereas glial cell numbers in NT-3-treated cultures were not different than those found in controls (data not shown).

These results confirm that both NT-3 and GDNF are able to affect the behaviour of enteric ganglion cells in this culture system. In order to minimize effects of cell death observed in cultures grown for 36 hours on the results of survival assays, further experiments were restricted to the 12 hour time point.

### *3.2 Effects of different concentrations of H<sub>2</sub>O<sub>2</sub> on enteric ganglion cells in culture*

Pilot experiments were performed to determine the response of enteric ganglion cells to different concentrations of H<sub>2</sub>O<sub>2</sub>. Cultures were exposed to 1, 5, 10 or 25  $\mu$ M H<sub>2</sub>O<sub>2</sub> then incubated for 6 hours before assessing cell viability by MTS assay. Effects of H<sub>2</sub>O<sub>2</sub> on the viability of enteric ganglion cells are shown in

Figure 2. While 1 $\mu$ M H<sub>2</sub>O<sub>2</sub> did not significantly affect cell viability, higher concentrations decreased it in a significant dose-dependent manner, with about 50% decrease in viability at 10 $\mu$ M H<sub>2</sub>O<sub>2</sub> compared to control. 25 $\mu$ M H<sub>2</sub>O<sub>2</sub> eliminated almost all cells, therefore for further experiments H<sub>2</sub>O<sub>2</sub> was used at 1, 5 and 10  $\mu$ M.

Analysis of PGP9.5-immunolabelled cultures treated in the same way as those used for MTS assay revealed that neurons and glial cells were differentially affected by H<sub>2</sub>O<sub>2</sub> (Table 1). Neurons were more vulnerable to H<sub>2</sub>O<sub>2</sub> than glial cells.

### *3.3 Effects of pre-treatment with neurotrophic factors on viability of H<sub>2</sub>O<sub>2</sub>-treated enteric ganglion cells*

After optimising culture time and H<sub>2</sub>O<sub>2</sub> concentrations, we investigated whether treatment of enteric neurons with 10ng/ml NT-3 or GDNF prior to H<sub>2</sub>O<sub>2</sub> exposure increased viability of enteric ganglion cells. Cultures grown with NT-3 exhibited a slightly increased survival compared to controls, although the difference was not statistically significant at any H<sub>2</sub>O<sub>2</sub> concentration (Figure 3A); GDNF treatment did not affect cell viability (Figure 3B).

The effects of factors were then examined in more detail, employing bis-benzimide and PI (live/dead) staining; bis benzimide stains nuclei of all cells, while PI is excluded from living cells, therefore allowing accurate measure of total cell numbers and the extent of cell survival. Cell cultures were incubated for 12 hours with 10ng/ml NT-3 or GDNF, then treated with 10 $\mu$ M H<sub>2</sub>O<sub>2</sub> and stained with PI and bis-benzimide 4 hours later. This shorter H<sub>2</sub>O<sub>2</sub> exposure time resulted in

reduced cell losses compared to those seen after 6 hours exposure (Table 1).  $H_2O_2$  treatment of control cultures without trophic factors resulted in a significant increase in PI-positive cells ( $p < 0.02$ , Figure 4A). Cultures grown with NT-3 prior to  $H_2O_2$  exposure exhibited significantly ( $p < 0.01$ ) decreased numbers of PI-stained cells compared to  $H_2O_2$  controls without factor treatment. GDNF treatment also decreased PI positive cells in cultures exposed to  $H_2O_2$ , but the effect was not statistically significant. Cultures not exposed to  $H_2O_2$  showed comparable numbers of PI stained cells in each factor treatment and in controls.

Counts of Hoechst positive cells are summarized in Figure 4B. In the absence of  $H_2O_2$ , NT-3 did not affect total cell numbers, but GDNF increased cell numbers compared to both control ( $p < 0.05$ ) and NT-3-treated cultures ( $p < 0.02$ ). This observation confirms the results of the cell counts performed on PGP9.5 immunolabelled cultures. Addition of  $H_2O_2$  had no effect on cell numbers in untreated or NT-3-treated cultures (Hoescht labels both live and dead cells), but interestingly decreased the total cell numbers in cultures grown with GDNF ( $p = 0.01$ ), reducing cell numbers to levels similar to those in control and NT-3-treated cultures exposed to  $H_2O_2$ .

#### 4. Discussion

In this study, we investigated the effects of NT-3 and GDNF on cultured enteric ganglion cells exposed to  $H_2O_2$ . Our results demonstrate decreased cell death in cultures grown with NT-3 prior to  $H_2O_2$  exposure, as opposed to control cultures, and cultures grown with GDNF. In contrast, GDNF stimulated an

increase in the number of cells present in the cultures, an effect abrogated by H<sub>2</sub>O<sub>2</sub> treatment.

Two previous studies have demonstrated that H<sub>2</sub>O<sub>2</sub> has a toxic effect on enteric neurons *in vitro*. These studies either used an organotypic model of rat myenteric ganglia, in which the ganglia remained intact [1,38] or a mixed preparation of cells from dissociated whole intestine from embryonic mice [1]. The cultures in the latter study were manipulated to remove enteric glia, which were thereby shown to have a protective effect against H<sub>2</sub>O<sub>2</sub>-induced neuronal toxicity [1]. The protective effects of NT-3 in the present study were unlikely to be due to an increased number of enteric glia in NT-3-treated cultures, because glial cell numbers were not significantly different between the NT-3-treated cultures and the control or GDNF-treated cultures. Moreover in previous work, in which dissociated enteric ganglia were grown in the presence of NT-3 for longer periods, glial numbers were not increased [31]. In future studies, however, counts of neurons and glia in H<sub>2</sub>O<sub>2</sub>-treated cultures should be performed.

NT-3, a member of the neurotrophin family, promotes differentiation of enteric neurons and glia from precursor cells during ENS development [5,32]. NT-3 has also been found to protect against menadione-induced apoptosis of myenteric neurons in the muscularis externa from calorically-restricted rats [37]. The present results are thus in keeping with this previous report. Other members of the neurotrophin family, notably BDNF, have been found to be protective against oxidative stress in several different neuronal culture systems [36,16,25].

GDNF is a member of the GDNF family of neurotrophic factors. It has been shown to perform a critical role in the survival, as well as proliferation of ENS precursor cells [5,32] and has previously reported to protect against oxidative stress in isolated intestinal preparations [37] and some other systems [e.g.26], and against hyperglycaemia –induced myenteric neuronal death in culture [2]. Here we found that GDNF did not prevent H<sub>2</sub>O<sub>2</sub>-induced cell death in cultures of enteric ganglion cells, under the conditions employed. We did observe however, that GDNF treatment increased the total number of cells in the cultures, as shown by the results of cell counts of both PGP9.5 and Hoechst stained cultures. Postnatal enteric ganglia contain neural precursors [33,34], so it is not unexpected that GDNF would promote differentiation of these cells *in vitro*. GDNF is known to have a number of actions in the postnatal ENS [28]. Interestingly, the present results suggest that new cells arising due to GDNF treatment may be more vulnerable to oxidative damage than differentiated neurons and glia present in the cultures. This possibility is suggested by the reduction of total cell numbers measured in GDNF-treated cultures exposed to H<sub>2</sub>O<sub>2</sub>, that reached levels found in non-factor treated controls. One possible explanation of this finding is differing sensitivities of mature and dividing cells to DNA-damaging agents, such as oxidative stress; immature cells undergoing apoptosis rather than activating DNA repair and cell survival programs [20,24].

Our findings raise the question of the mechanisms involved in the protective effect of NT-3 on enteric ganglion cells. One possible explanation is that NT-3 might influence protein levels or activity of antioxidant enzymes [23].

Evidence supporting this possibility includes observations that nerve growth factor stimulates activity of glutathione peroxidase [11] and BDNF exerts protective effects on auditory neurons via increased levels of glutathione [15]. Mattson et al. [23] demonstrated that neurotrophic factors increased antioxidant enzyme activity in cultured hippocampal neurons. Neurotrophic factor treatment could also lead to reduced ROS generation; NT-3 treatment decreased generation of free radicals by myenteric neurons in muscularis externa preparations [37]. Finally, it is known that neurotrophic factors activate signalling pathways that promote expression of other survival-promoting proteins [e.g.29].

In conclusion, these results provide further support for the suggestion that neurotrophic factors, particularly NT-3, may have protective roles in the ENS. Such an effect could have important implications not just for aging, but also disease states such as diabetes, which affect the ENS and gastrointestinal functions [e.g.8]. In this context, it is important to note that oxidative stress has been implicated in enteric diabetic neuropathy [8,19,38] and that recent evidence has shown a reduction in the levels of neurotrophic factors in the diabetic gut [21].

**Acknowledgements:**

This work was supported by BBSRC SAGE Initiative grant 108/SAG10013. KK was an OU funded student. We thank Tina Wardhaugh and OU BRU staff for technical assistance.

## References

1. **H.Abdo, P.Derkinderen, P. Gomes, J. Chevalier, P. Aubert, D. Masson, J.-P. Galmiche, P.Vanden Berghe, M. Neunlist, B. Lardeux**, Enteric glial cells protect neurons from oxidative stress in part *via* reduced glutathione *FASEB J.* 24(2010):1082-94
2. **M. Anitha, C. Gondha, Sutliff R, Parsadonian A, Mwangi S, Sitaraman SV, Srinivasan S**. GDNF rescues hyperglycemia-induced diabetic enteric neuropathy through activation of the PI3K/Akt pathway. *The Journal of Clinical Investigation* 116(2006):344-356
3. **K.Bitar, B. Greenwood-Van Meerveld, R.Saad, J. Wiley**, (2011). Aging and gastrointestinal neuromuscular function: insights from within and outside the gut. *Neurogastroenterol. Motil.* 23(2011) 490-501
4. **J.C.Bornstein, M.Costa, J.R.Grider**, Enteric motor and interneuronal circuits controlling motility. *Neurogastroenterol. Motil.*16(2004) 34–38.
5. **G.B.T. von Boyen, M. Reinshagen, M. Steinkamp, G. Adler, J. Kirsch**, Enteric nervous plasticity and development: dependence on neurotrophic factors. *J. Gastroenterol.* 37(2002) 583–588
6. **S.J.H. Brookes**, Classes of enteric nerve cells in the guinea-pig small intestine. *The Anatomical Record*, 262(2001) 58–70.
7. **M. Camilleri, T. Cowen, T.Koch**, Enteric neurodegeneration in ageing.

Neurogastroenterology and Motility 20(2008) 418–429.

**8. B. Chandrasekharan, M. Anitha, R. Blatt, N. Shahnava, D. Kooby, C.**

**Staley, S. Mwangi, D. P. Jones, S.V. Sitaraman, S. Srinivasan**

Colonic motor dysfunction in human diabetes is associated with enteric neuronal loss and increased oxidative stress. *Neurogastroenterol.Motil.* ;23(2011):131-139

**9. M V Clement, A Ponton, S Perivaiz,** Apoptosis mediated by hydrogen peroxide is mediated by decreased superoxide anion concentration and reduction of intracellular milieu. *FEBS Lett.* 440(1998), 13-18

**10. T.Cowen, R.J.R.Johnson, V. Soubeyre, R. M. Santer,** Restricted diet rescues rat enteric motor neurones from age related cell death. *Gut* 47(2000) 653-660

**11. C Guegan, I C Picot, E Chevalier, A Nicole, B Sola,** Reduction of ischemic damage in NGF transgenic mice: correlation with enhancement of antioxidant enzyme activities *Neurobiol. Disease*, 6(1999) 180-189

**12. L. L. Dugan, D. J. Creedon, E. M. Johnson, Jr, D. M. Holtzman,** Rapid suppression of free radical formation by nerve growth factor involves the mitogen-activated protein kinase pathway. *PNAS* 94(1997) 4086–4091

**13. J. B Furness,** Types of neurons in the enteric nervous system, *J. Aut. Nerv. Sys.*, 81(2000) 87-96

**14. K Fuson, R Mark, J Panetta, P May,** Characterisation of LY231617 protection against hydrogen peroxide toxicity *J. Neurochem.*, 72(1999) 1154-1160

- 15. R Gabaizadeh, H Staecker, W Liu, T R Van De Water,** BDNF protection of auditory neurons from cisplatin involves changes in intracellular levels of both reactive oxygen species and glutathione. *Mol. Brain Res.* 50(1997) 71–78
- 16. G.E. Hardingham, S.A. Lipton,** Regulation of Neuronal Oxidative and Nitrosative Stress by Endogenous Protective Pathways and Disease Processes. *Antioxidants and Redox Signalling* 14(2011) 1421-1424
- 17. K Hoyt, A Gallagher, T Hastings, I Reynolds,** Characterisation of hydrogen peroxide toxicity in cultured rat forebrain neurons. *Neurochem. Res.*, 22(1997) 333-340
- 18. J.A. Imlay** Cellular defenses against superoxide and hydrogen peroxide. *Annu Rev Biochem.* 77(2008) 755–776
- 19. P. Kashyap, G. Farrugia,** Oxidative stress: key player in gastrointestinal complications of diabetes. *Neurogastroenterol Motil* 23(2011) 111–114
- 20. C L Limoli, E Giedzinski, R Rola, S Otsuka, T D Palmer, J R Fike,** Radiation response of neural precursor cells: linking cellular sensitivity to cell cycle checkpoints, apoptosis and oxidative stress, *Radiat Res.*, 161(2004), 17-27
- 21. W. Liu, W. Yue, R. Wu,** Effects of diabetes on expression of glial fibrillary acidic protein and neurotrophins in rat colon. *Auton Neurosci.* 154(2010):79-83
- 22. M.P. Mattson,** Glutamate and Neurotrophic Factors in Neuronal Plasticity and Disease. *Ann N Y Acad Sci.* 1144(2008) 97–112
- 23. M.P. Mattson, M.A. Lovell, K. Furukawa, W.R. Markesbery,** Neurotrophic factors attenuate glutamate-induced accumulation of peroxides, elevation of

- intracellular Ca<sup>2+</sup> concentration, and neurotoxicity and increase antioxidant enzyme activities in hippocampal neurons. *J Neurochem.* 65(1995) 1740-51.
- 24. R Mischel, Y Kim, R Sheldon, D Ferriero**, Hydrogen peroxide is selectively toxic to immature murine neurons *in vitro*. *Neurosci. Lett.*, 231(1997), 17-20
- 25. T. Numakawa, T. Matsumoto, Y. Numakawa, M. Richards, S. Yamawaki, H. Kunugi**, Protective Action of Neurotrophic Factors and Estrogen against Oxidative Stress-Mediated Neurodegeneration. *J Toxicol.* 2011(2011) 405194
- 26. I.G. Onyango, J. B. Tuttle, J. P. Bennett Jr.**, Brain-derived growth factor and glial cell line-derived growth factor use distinct intracellular signaling pathways to protect PD cybrids from H<sub>2</sub>O<sub>2</sub>-induced neuronal death. *Neurobiol Dis.* 20(2005)141-54
- 27. R.Phillips, T.Powley**, Innervation of the gastrointestinal tract: patterns of aging. *Autonomic Neuroscience*, 136(2007), 1–19.
- 28. D. M. Rodrigues, A. Y. Li, D. G. Nair, M. G. Blennerhassett**, Glial cell line-derived neurotrophic factor is a key neurotrophin in the postnatal enteric nervous system. *Neurogastroenterol Motil* 23(2011) e44–e56
- 29.P. S. Sastry, K. S. Rao** Apoptosis and the Nervous System. *J.Neurochem* 74(2000) 1-20
- 30. M.J.Saffrey, Ageing of the enteric nervous system.** *Mech Ageing Dev.* 125(2004):899-906
- 31. M.J.Saffrey, T.Wardhaugh, T.Walker, J.Daisley, A. T. Silva**, Trophic actions of neurotrophin-3 on postnatal rat myenteric neurons *in vitro*, *Neurosci. Letters*, 278(2000), 133-136

- 32. M.J.Saffrey**, Enteric Nervous System:Neurotrophic Factors. In:Ed. Greg Lemke. Developmental Neurobiology. Elsevier. 2009 pp 628-634
- 33. K-H. Schafer, C. Van Ginneken, S. Copray**, Plasticity and Neural Stem Cells in the Enteric Nervous System. The Anatomical Record 292(2009)1940–1952
- 34. A.T.Silva, T.Wardhaugh, N.F.Dolatshad, S.Jones, M.J.Saffrey**, Neural progenitors from isolated postnatal rat myenteric ganglia: expansion as neurospheres and differentiation in vitro.Brain Res 1218(2008) 47-53
- 35. S.D.Skaper, M.Floreani, A.Negro,L.Facci, P.Giusti**, Neurotrophins rescue cerebellar granule neurons from oxidative stress mediated apoptotic death: selective involvement of phosphatidylinositol 3-kinase and the mitogen activated protein kinase pathway, J. Neurochem. 70(1998) 1859-1868
- 36. S.D. Skaper**,. The Biology of Neurotrophins, Signalling Pathways, and Functional Peptide Mimetics of Neurotrophins and their Receptors CNS and Neurological Disorders - Drug Targets, 7(2008) 46-62
- 37. C. Thrasivoulou, V. Soubeyre, H.Ridha, D.Giuliani, C.Giaroni,G.Michael, M.J.Saffrey, T.Cowen**, Reactive oxygen species, dietary restriction and neurotrophic factors in age-related loss of myenteric neurons. Aging Cell, 5(2006), 247–257.
- 38. E. Voulaki, H.R.Shotton, J.Lincoln** Selective responses of myenteric neurons to oxidative stress and diabetic stimuli. Neurogastroenterol Motil 23(2011) 964–e411

- 32. M.J.Saffrey**, Enteric Nervous System:Neurotrophic Factors. In:Ed. Greg Lemke. Developmental Neurobiology. Elsevier. 2009 pp 628-634
- 33. K-H. Schafer, C. Van Ginneken, S. Copray**, Plasticity and Neural Stem Cells in the Enteric Nervous System. The Anatomical Record 292(2009)1940–1952
- 34. A.T.Silva, T.Wardhaugh, N.F.Dolatshad, S.Jones, M.J.Saffrey**, Neural progenitors from isolated postnatal rat myenteric ganglia: expansion as neurospheres and differentiation in vitro.Brain Res 1218(2008) 47-53
- 35. S.D.Skaper, M.Fioreani, A.Negro,L.Facci, P.Giusti**, Neurotrophins rescue cerebellar granule neurons from oxidative stress mediated apoptotic death: selective involvement of phosphatidylinositol 3-kinase and the mitogen activated protein kinase pathway, J. Neurochem. 70(1998) 1859-1868
- 36. S.D. Skaper**,. The Biology of Neurotrophins, Signalling Pathways, and Functional Peptide Mimetics of Neurotrophins and their Receptors CNS and Neurological Disorders - Drug Targets, 7(2008) 46-62
- 37. C. Thrasivoulou, V. Soubeyre, H.Ridha, D.Giuliani, C.Giaroni,G.Michael, M.J.Saffrey, T.Cowen**, Reactive oxygen species, dietary restriction and neurotrophic factors in age-related loss of myenteric neurons. Aging Cell, 5(2006), 247–257.
- 38. E. Voulaki, H.R.Shotton, J.Lincoln** Selective responses of myenteric neurons to oxidative stress and diabetic stimuli. Neurogastroenterol Motil 23(2011) 964–e411

**Figure Legends:**

**Figure 1:** Mean cell numbers in cultures after 12 and 36 hours incubation with NT-3 or GDNF (10ng/ml). Data from 3 separate experiments; 3 replicas per experiment. Error bars represent  $\pm$ S.E.M. \*\*\*  $p<0.001$ ; \*\* $p<0.01$

**Figure 2:** Survival of cultured enteric ganglion cells exposed to different concentrations of  $H_2O_2$ , measured using MTS assay. Results are expressed as percentage of untreated controls. Data are means of 3 separate experiments; 4 replicas per experiment. Error bars represent  $\pm$ S.E.M. \*\*\* $p<0.001$


**Figure 3:** Survival of enteric ganglion cells treated with 10ng/ml NT-3 (**A**) or GDNF (**B**) prior to  $H_2O_2$  exposure. Means from 3 separate experiments, expressed as a percentage of untreated controls; 4 replicas per experiment. Error bars represent  $\pm$ S.E.M.

**Figure 4:** Total number of PI (**A**) and Hoechst (**B**) positive cells in NT-3 and GDNF (10ng/ml) treated cultures after exposure to  $H_2O_2$ , compared to controls. Graphs show mean number of stained cells counted in 3 separate experiments; 3 replicas per experiment. Error bars represent  $\pm$ S.E.M.

\*\*\* $p\leq 0.01$ ; \*\* $p<0.02$ ; \* $p\leq 0.05$

Table 1: Effects of H<sub>2</sub>O<sub>2</sub> on neuronal and glial cell numbers

<b>Treatment</b>	<b>Number of neurons (± SEM)</b>	<b>Number of glial cells (± SEM)</b>	<b>N/GC ratio</b>
<b>Control</b>	440 ± 9.3	912 ± 14.6	0.48
<b>5 μM H<sub>2</sub>O<sub>2</sub></b>	129 ± 13.9	404 ± 33.6	0.32
<b>10 μM H<sub>2</sub>O<sub>2</sub></b>	15 ± 1	88 ± 5.1	0.17
<b>25 μM H<sub>2</sub>O<sub>2</sub></b>	1	15 ± 2.6	0.06



Preprint

