

Open Research Online

The Open University's repository of research publications
and other research outputs

Implantation of multiply charged carbon ions in water ice

Journal Item

How to cite:

Lv, X. Y.; de Barros, A. L. F.; Boduch, P.; Bordalo, V.; da Silveira, E. F.; Domaracka, A.; Fulvio, D.; Hunniford, C. A.; Langlinay, T.; Mason, N. J.; McCullough, R. W.; Palumbo, M. E.; Pilling, S.; Rothard, H. and Strazzulla, G. (2012). Implantation of multiply charged carbon ions in water ice. *Astronomy & Astrophysics*, 546, article no. A81.

For guidance on citations see [FAQs](#).

© 2012 ESO

Version: Version of Record

Link(s) to article on publisher's website:

<http://dx.doi.org/doi:10.1051/0004-6361/201219886>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

Implantation of Multiply Charged Carbon Ions in Water Ice

X. Y. Lv^{1,2*}, A. L. F. de Barros³, P. Boduch¹, V. Bordalo⁴, E. F. da Silveira⁴, A. Domaracka¹, D. Fulvio⁵, C. A. Hunniford⁶, T. Langlinay¹, N. J. Mason⁷, R. W. McCullough⁶, M. E. Palumbo⁸, S. Pilling⁹, H. Rothard¹, and G. Strazzulla⁸

¹ Centre de Recherche sur les Ions, les Matériaux et la Photonique (CEA/CNRS/ENSICAEN/UCBN), CIMAP - CIRIL - GANIL, Boulevard Henri Becquerel, BP 5133, F-14070 Caen Cedex 05, France

² School of Nuclear Science and Technology, Lanzhou University, Lanzhou 730000, China
e-mail: lv@ganil.fr

³ Departamento de Disciplinas Básicas e Gerais, CEFET-RJ, Av. Maracanã 229, 20271-110 Rio de Janeiro, RJ, Brazil

⁴ Departamento de Física, Pontifícia Universidade Católica do Rio de Janeiro, Rua Marquês de São Vicente 225, 22451-900, Rio de Janeiro, RJ, Brazil

⁵ Laboratory for Atomic and Surface Physics, University of Virginia, 351 McCormick Road, P.O. Box 400238, Charlottesville, VA 22904-4238, USA

⁶ Centre for Plasma Physics, School of Mathematics and Physics, Queen's University, Belfast, Belfast BT7 1NN, Northern Ireland

⁷ Department of Physics and Astronomy, The Open University, Walton Hall, Milton Keynes, UK

⁸ INAF Osservatorio Astrofisico di Catania, Catania, Italy

⁹ IP&D/UNIVAP, Av. Shishima Hifumi, 2911, São Jose dos Campos, SP, Brazil

Preprint online version: June 25, 2012

ABSTRACT

Context. Several objects in the Solar System like Europa, Ganymede and Callisto have frozen surface (main component: H₂O). The associated thickness is bigger than the penetration depth of the relevant projectile ions. Additionally, other species such as H₂O₂, SO₂ and CO₂ have been detected on these surface. The formation mechanisms of these molecules are still under discussion.

Aims. We present new experimental results on the implantation of ¹³C^{q+} (q = 2, 3) ions at an energy of 30 keV in water ice at low temperatures (15 and 80 K). Experiments with multiply-charged ions at energies of tens of keV are particularly relevant to simulating the complexity of the irradiation environment to which the surfaces of the icy moons in the outer solar system are exposed.

Methods. The experiments were performed at the low energy ion beam facility ARIBE of GANIL in Caen (France). 30 keV ¹³C^{q+} (q = 2, 3) ions have been used to bombard solid H₂O surface which were frozen at 15K and 80K. Fourier Transform Infrared Spectrometer (FTIR) was used to analyze the sample in the 5000 - 600 cm⁻¹ (2-16.7 μm) region with a spectral resolution of 1 cm⁻¹.

Results. The results of our experiments indicate that implantation produces ¹³CO₂ with yields in the range of 0.32-0.57 molecules ion⁻¹. This yield seems to be independent of the temperature of the ices in the range studied. We have estimated the time scale necessary to accumulate by implantation of magnetospheric carbon ions the observed quantity of carbon dioxide on the surface of Europa, a Jovian moon. This time scale is of the order of 1.0-1.3×10⁴ yrs which is higher than that evaluated for carbon dioxide production by other relevant processes.

Conclusions. We conclude that although a relevant quantity of CO₂ can be formed by carbon ion implantation, this is not the dominant formation mechanism at Europa.

Key words. planets and satellites: surfaces – methods: laboratory – techniques: spectroscopic

1. Introduction

Ion implantation is a very common technique for the doping (i.e. introduction of impurities) of different materials. It is widely used in the semiconductor industries for e.g. modifying the electrical properties (such as the threshold voltage) of transistors. Dopant ions are produced, accelerated, selected by the mass-to-charge ratio, and directed to the target. The ions penetrate into the target, collide with the host atoms, lose energy, and remain implanted at some depth within the solid. The penetration depth depends on the mass and energy of the incoming ion and on the properties of the target. Experiments are performed with energies from several hundred to several million electron volts, resulting in ion implantation depths from <10 nm to >10 μm.

Implantation experiments are also relevant in very different scenarios as those of reactive ions (H, O, C, N, S) (Baratta

et al. 2012) irradiating ices in several astrophysical environments. Implanted ions deposit energy in the icy target (for a detailed description of the phenomenology of interaction between fast ions and target species see Johnson 1990). In addition only reactive ions (carbon in this paper) have a chance to form species that include the implanted projectile with a maximum yield of one molecule per incoming ion.

There are several objects in the Solar System where ices on planetary surfaces, comets, etc., are much thicker than the penetration depth of the relevant ion populations. As an example, earth based and space observations indicate that on the surface of Europa, Ganymede and Callisto water ice is the dominant species along with hydrated materials and minor amounts of some volatile species such as H₂O₂, SO₂ and CO₂ (Carlson et al. 1997, 1999; Carlson 2001; Noll et al. 1995, 1997). The formation mechanism of those molecules is still an open question. A possible way is via exogenic processes such as implantation of

* corresponding author

carbon or sulfur ions of the Jovian magnetosphere, as suggested since several years on the basis of numerous experimental results (for a review see Strazzulla 2011).

In this paper we present new experimental results on the implantation of $^{13}\text{C}^{q+}$ ($q = 2, 3$) ions at an energy of 30 keV in water ice at low temperatures (15 and 80 K). Experiments with multiply charged ions at energies of tens of keV are particularly relevant to simulating the complexity of the bombardment environment to which the surfaces of the icy moons in the outer Solar System are exposed.

The present results confirm and extend those obtained by bombardment of frozen layers of pure H_2O ice with 10-30 keV singly charged (Bibring & Rocard 1984; Strazzulla et al. 2003) and singly/doubly charged 2-4 keV (Dawes et al. 2007; Hunniford et al. 2009) carbon ions. The results of our experiments indicate that implantation produces $^{13}\text{CO}_2$ with yields in the range of 0.32-0.57 molecules ion $^{-1}$.

2. Experiments

The experiments were performed at the low energy ion beam facility ARIBE of GANIL in Caen (France). $^{13}\text{C}^{q+}$ ($q = 2, 3$) ion beams ($\sim 10^{12}$ ions cm^{-2} s^{-1}) were produced in a 14.5 GHz electron cyclotron resonance (ECR) ion source. After selection of mass and charge by a dipole magnet, the beams were guided toward the experimental set-up (shown in Figure 1). A sweeping device made of two parallel plates assured a uniform bombardment of the target. The beam passed through an insulated collimator. A part of the beam was stopped by the collimator and the corresponding collimator current was measured. A Faraday-Cup was inserted into the beam about 12 cm upstream of the target. From the ratio of the current measured on the collimator and in the Faraday-Cup, the beam flux (number of projectiles per square centimeter and second) on the target can be determined. The accumulated charge on the collimator was measured by means of current integrators. From this accumulated charge and the ratio of the current on the collimator and the current in the Faraday-Cup, the projectile fluence (number of projectiles accumulated per square centimeter) could be determined. The ratio was checked several times before, during and after bombardment to monitor for any variation of the flux.

The ice layers were prepared by condensing water in the gas phase on a CsI window at 15 K and 80 K. About 1 cm^3 pure H_2O which was produced by a Milli-Q Integral Water Purification System was stored in a cuvette. The H_2O was frozen to solid phase, then while pumping the cuvette for a few minutes, the H_2O was warm up to liquid phase. This procedure was repeated 2 or 3 times in order to obtain of H_2O vapor of high purity. H_2O vapor was introduced in a pre-chamber, A fine valve, which allowed controlling the deposition rate, was used to transmit the gas into the high vacuum chamber and onto the cold CsI substrate. The pressure in the high vacuum chamber was below 10^{-7} mbar during the measurements. The CsI window was installed in the center of the chamber on a cold finger connected to a closed cycle helium cryostat. The temperature of the substrate was controlled by a carbon resistance and a compound linear thermal sensor (CLTS) situated on the holder, providing a precision of 0.1 K. The cold head with the CsI window could be rotated from 0° to 180° and fixed in three positions allowing bombardment (0°), FTIR analysis (90°) and deposition (180°). A Nicolet Magna 550 Fourier Transform Infrared Spectrometer (FTIR) was used to analyze the sample in the 5000 - 600 cm^{-1} (2-16.7 μm) region with a spectral resolution of 1 cm^{-1} . The

spectra were always corrected by a background recorded before deposition.

The thickness of the sample (estimated from their column density assuming an ice density of 1 g cm^{-3} , details of calculation can be found in Pilling et al. 2011) was between 0.7 μm and 1.8 μm (Table 1). The range (or penetration depth) of 30 keV carbon ions in water ice is of the order of 178 ± 50 nm (Ziegler et al. 2008). Thus, the ices that we deposited were sufficiently thick to assure complete stopping and implantation of all projectiles.

3. Results

Figure 2 shows spectra measured before and after bombardment of H_2O ices deposited at 15 K and 80 K, respectively, with 30 keV $^{13}\text{C}^{q+}$ ($q = 2, 3$) ions. Up to two different absorption bands are observed. The band at 2346 cm^{-1} belongs to the ν_3 vibration mode of $^{12}\text{CO}_2$ (Sandford & Allamandola 1990; Palumbo et al. 1998; Jamieson et al. 2006; Seperuelo Duarte et al. 2010; Pilling et al. 2010, 2011) and is observed before bombardment. Figure 2 clearly shows that the area of 2346 cm^{-1} are not proportional to the thickness of the ices, thus the impurities in the original water can be excluded. It is believed to mainly stem from residual gas in the pre-chamber and the bombardment chamber but not from CO_2 impurities in the original H_2O ice, since it not only depends on the pressure in the chamber and pre-chamber but also on the elapsed time between deposition and measurement after the background recording. The band at 2277 cm^{-1} belongs to the ν_3 vibration mode of $^{13}\text{CO}_2$ (Gerakines et al. 1995; Strazzulla et al. 2005; Seperuelo Duarte et al. 2010). The intensity of this band is increasing during the bombardment, indicating that the corresponding new formed $^{13}\text{CO}_2$ molecules are clearly correlated with the implantation of $^{13}\text{C}^{q+}$.

We calculated the column density of $^{12}\text{CO}_2$ and $^{13}\text{CO}_2$ by assuming a band strength of 7.6×10^{-17} cm molecule^{-1} and 7.8×10^{-17} cm molecule^{-1} , respectively (Yamada & Person 1964; Gerakines et al. 1995; Jamieson et al. 2006). This was accomplished by using the equation 1:

$$N = \frac{2.3 \int \tau(\nu) d\nu}{A} \quad (1)$$

where N represents the column density (in molecules cm^{-2}) of $^{12}\text{CO}_2$ and $^{13}\text{CO}_2$, A represents the band strength (in cm molecule^{-1}), $\int \tau(\nu) d\nu$ represents the area (in cm^{-1}) of a given measured band on absorbance scale. On the one hand, $^{13}\text{CO}_2$ is produced in the target by implantation of $^{13}\text{C}^{q+}$ ions. On the other hand, $^{12}\text{CO}_2$ from the residual gas could also condense on the substrate. This has been taken into account by assuming a natural isotopic ratio of ^{12}C and ^{13}C in the atmosphere of about 90 to determine the amount of $^{13}\text{CO}_2$ associated with the implantation process.

Figure 3 shows the column density of produced $^{13}\text{CO}_2$ as a function of the fluence of implanted 30 keV $^{13}\text{C}^{3+}$ and $^{13}\text{C}^{2+}$ ions into water ices at 15 K and 80 K. Also included are experimental data obtained in Catania (Italy), for 30 keV $^{13}\text{C}^{+}$ implantation in water ice at slightly different temperatures of 16 K and 77 K (Strazzulla et al. 2003), for comparison. $^{13}\text{CO}_2$ production yields are shown in the fourth column of Table 1. Within error bars, which stem from the uncertainties of the determination of the column density on one hand, and the uncertainty of the fluence measurement on the other hand, the results obtained in Catania for $^{13}\text{C}^{+}$ and in Caen for $^{13}\text{C}^{2+}$ and $^{13}\text{C}^{3+}$ are in good agreement. All of these results can be summarized as follows:




Fig. 1. Schematic diagram of our experimental apparatus.

Table 1. Summary of implantation yields in molecules ion⁻¹.

Ions	Energy /keV	Temperature /K	Thickness / μm	Implantation yield /molecules ion ⁻¹	References
¹³ C ³⁺	30	15	1.8	0.32 ± 0.07	This work
¹³ C ²⁺	30	15	0.8	0.57 ± 0.13	This work
¹³ C ³⁺	30	80	0.8	0.40 ± 0.09	This work
¹³ C ²⁺	30	80	0.7	0.50 ± 0.11	This work
¹³ C ⁺	30	16	0.6	0.47 ± 0.10	Strazzulla et al. (2003)
¹³ C ⁺	30	77	0.6	0.42 ± 0.09	Strazzulla et al. (2003)




Fig. 2. Spectra of water ices bombarded with 30 keV ¹³C³⁺ and ¹³C²⁺ ions at 15 K and 80 K, respectively. Dashed lines represent spectra before bombardment, solid lines represent spectra after bombardment (about 10¹⁶ ions cm⁻²).

- ¹³CO₂ was produced after implantation of 30 keV ¹³C^{q+} ions with different charge states into H₂O ices deposited at different temperatures.
- No effects related to the temperature of the ices have been observed.
- No effects related to the charge states of the incident ions have been found.

The main difference between samples deposited at 15 K and 80 K is that the sample deposited at lower temperature has a higher porosity than that deposited at higher temperature (Baragiola 2003; Palumbo 2006). This has been confirmed by the observation, in our spectra, of the O-H dangling bonds (db) bands that peak at about 3720 and 3695 cm⁻¹ in water ice deposited at 15 K and at 3691 cm⁻¹ for 80 K ice (Palumbo & Strazzulla 2003; Palumbo 2005). Pilling et al. (2011) have shown that for H₂O:CO₂ (1:1) ices, the sample deposited at 80 K has the OH db (3690-3633 cm⁻¹) area about 10-50 times smaller than the sample deposited at 13 K. Our result that no temperature effects have been observed shows that the porosity does not play an important role in the formation of carbon dioxide during 30 keV ¹³C^{q+} interaction with pure water ices.

The interaction of slow ions impacting on solid surface has been studied extensively (Arnau et al. 1997). Ions approaching the surface will capture electrons from the target surface to the excited states of the ions (the number of captured electrons depending on the charge states of the incoming ions). These electron capture processes take place only at the surface. As soon as the projectiles enter into the bulk, they will rapidly reach the effective charge state (typically within one single monolayer) which should be the same for all of the implanted ¹³C²⁺ and ¹³C³⁺ ions at the same velocity (Herrmann et al. 1994). The memory of the incoming charge state is lost. Indeed, our results show that the values of the ¹³CO₂ production yields do not vary with the projectile charge state. The mean value is 0.45 molecules of ¹³CO₂ formed per incoming projectile ion.

Dawes et al. (2007) reported results on 4 keV ¹³C⁺ and ¹³C²⁺ implantation into water ice deposited at 30 K and 90 K




Fig. 3. Column density of $^{13}\text{CO}_2$ produced after implantation of 30 keV $^{13}\text{C}^{q+}$ ($q = 1, 2, 3$) ions at 15 K and 80 K. The experimental data for $^{13}\text{C}^+$ implanted at 16 K and 77 K in water ice were obtained in Catania (Strazzulla et al. 2003). The solid lines represent linear least square fits, the corresponding production yields are shown in Table 1.




Fig. 4. Comparison of $^{13}\text{CO}_2$ the production yields between our results and the results of Dawes et al. (2007) Data from singly and doubly charged projectiles are presented in left and right figures, respectively.

(shown in Figure 4). For $^{13}\text{C}^+$ ions, the column density of synthesized $^{13}\text{CO}_2$ showed an asymptotic increase, reaching saturation at higher fluences. For $^{13}\text{C}^{2+}$ ions, the evolution of the column density of synthesized $^{13}\text{CO}_2$ with fluence can be described as the sum of two asymptotically increasing functions (see also Hunniford et al. 2009). Viewing the result as a whole (shown in Figure 4 where also some of our results are reported to facilitate the comparison), the reported yields of $^{13}\text{CO}_2$ by Dawes et al. (2007) are much smaller than those observed in the experiments at both Catania (Strazzulla et al. 2003) and Caen (this work) (shown in Table 1).

The electronic stopping power of 30 keV $^{13}\text{C}^{q+}$ ($S_e \approx 85 \text{ keV } \mu\text{m}^{-1}$) dominates over the nuclear stopping power ($S_n \approx 62 \text{ keV } \mu\text{m}^{-1}$). For 4 keV $^{13}\text{C}^{q+}$ ions, the nuclear stopping power ($S_n \approx 110 \text{ keV } \mu\text{m}^{-1}$) is much larger than the electronic stopping power ($S_e \approx 30 \text{ keV } \mu\text{m}^{-1}$) that we calculated by using the avail-

able software SRIM-2008 (Ziegler et al. 2008). Comparing our results to those of Dawes et al. (2007) one could be led to the conclusion that electronic stopping power plays an important role in driving chemical reactions between implanted particles and neighbor radicals. According to a SRIM 2008 calculation (Ziegler et al. 2008), the sputtering yields were found to be 0.15 and 0.33 molecules ion^{-1} for 30 keV $^{13}\text{C}^{q+}$ and 4 keV $^{13}\text{C}^{q+}$ ions respectively. This means that at the end of the bombardment, only less than 10 monolayers H_2O molecules have been sputtered by 4 keV carbon ions, while the number of molecules sputtered by 30 keV ions can be neglected. This is not sufficient to explain the differences of $^{13}\text{CO}_2$ production yield between 30 keV and 4 keV $^{13}\text{C}^{q+}$ ions. The amount of ions associated with the back scattering effect is less than 1% for both projectiles, therefore this effect can also be neglected. The penetration depth were found to be of the order of 178 ± 50 and 25 ± 9 nm for 30 keV $^{13}\text{C}^{q+}$ and 4 keV $^{13}\text{C}^{q+}$ ions, respectively (Ziegler et al. 2008). This means that 30 keV $^{13}\text{C}^{q+}$ ions interact with a greater number of H_2O molecules than 4 keV $^{13}\text{C}^{q+}$ ions. If the implantation yields are compared to the number of H_2O molecules exposed to the ions, the ratio between implantation yields of 30 keV $^{13}\text{C}^{q+}$ and 4 keV $^{13}\text{C}^{q+}$ ions should be around 7 to 1, but this is not that case in our experimental results. Also, a larger longitudinal and lateral straggling is more important for 30 keV $^{13}\text{C}^{q+}$ ions. Since the sputtering effect and the amount of back scattered ions are negligible, only the range of the scattering region could explain the difference of our present results with those of Dawes et al. (2007). This difference, however remains an open question. Complementary experiments are needed to understand the basic physical processes occurring during implantation.

4. Discussion and conclusion

The surfaces of the icy moons in the Jovian system are subjected to intense bombardment by electrons, protons and multiply charged ions (Johnson et al. 2004; Bagenal et al. 2007). It is usual to refer to ions and electrons at or below 10 keV as "plasma" and above as "energetic particles". The plasma nearly corotates with Jupiter and then it flows preferentially onto the hemisphere trailing the satellite's motion. The less abundant but higher energy particles bombard the satellite in more complex ways exhibiting different spatial distributions depending on their energy, mass, charge, and the electric and magnetic fields near each moon (see Johnson et al. 2004, and references therein).

The temperatures of the water ice samples investigated in this paper are adequate to simulate those measured on many of the icy moons in the external solar system. As examples the temperatures of the Europa's surface are measured to be between 50 and 125 K; on Enceladus they are between 30 and 145 K and between 50 and 100 K on Rhea (McFadden et al. 2007; Spencer et al. 2006).

With this in mind it is clear that measuring in the laboratory parameters such as the implantation yields of relevant ions (C in the present case) in a wide range of charge states and energies are of great interest. Nowadays the progress in the field of space observations yields spatial resolutions high enough that the amount of a given chemical species can be mapped on the surface and related to exogenic effects such as magnetospheric charged particle bombardment, ultraviolet photolysis, impacts. As an example analysis of data relative to Europa have shown that sulfuric acid hydrate abundance is independent of underlying geologic unit, but is very well correlated with the flux of bombarding sulfur ions (Dalton et al. 2011). Interestingly a first set of experiments have demonstrated that sulfur ion implanta-

tion produces hydrated sulfuric acid with a high yield (Strazzulla et al. 2007).

At present CO₂ has been observed on the surfaces of the galilean moons and other objects in the outer solar system (for a review see Dalton et al. 2010). Its abundance and its superficial distribution is not known. McCord et al. (1997, 1998) first noticed that CO₂, because of the absence of vibro-rotational spectral signatures, should be in a condensed phase, possibly dispersed in another material. The same authors evaluated that a CO₂ column density as low as 3×10^{17} molecules cm⁻² is required to produce the absorption observed at Europa McCord et al. (1997, 1998).

Different possibilities exist to explain its presence at the surface. Some are endogenic: it could be degassed from internal sources (Moore et al. 1999). Other mechanisms are exogenic such as those that have to do with energetic processing of the surface by ions, electrons and photons and/or with carbon implantation. Concerning the energetic processing by ions, a large amount of experiments have shown that icy mixtures of H₂O with a plethora of carbon-bearing molecules (e.g. CO, CH₄, CH₃OH etc) all produce CO₂ although with different efficiencies (Palumbo et al. 1998; Ioppolo et al. 2009; Pilling et al. in press). The same is also true for photons and electrons (Watanabe et al. 2007; Baratta et al. 2002; Gerakines et al. 1996; Wada et al. 2006; Bennett et al. 2011). It is also been demonstrated that CO₂ can be formed at the interface between water ice and different kinds of solid carbon substrates (Mennella et al. 2004; Gomis & Strazzulla 2005; Raut et al. 2012) and it has been suggested that this could produce enough carbon dioxide to explain the observed amount.

Concerning the possibility that carbon implantation plays an important role, the knowledge of the flux of carbon ions impinging on the surfaces of the moons is needed. Those fluxes are, however, not known with great precision. A rough estimation of the average flux of carbon ions has been used by Strazzulla et al. (2003) In fact the fluxes of sulphur (keV-MeV) ions at the surfaces of the galilean moons have been reported (it values $\sim 9 \times 10^6$ S ions cm⁻² s⁻¹ at Europa (Cooper et al. 2001)) and although the abundance of carbon ions has not been quantified, the number ratio C/S has been reported. As an example, it amounts to about 0.2 near by Europa in the 1 MeV nucl⁻¹ range (Hamilton et al. 1981; Cohen et al. 2001). Assuming C/S = 0.2 at Europa, and using the experimental values ranging between 0.4 and 0.5 CO₂ molecules (impinging-ion)⁻¹ as production yield (see table 1; here we use the results obtained at 80 K, a temperature more appropriate to the Jovian moons) we can evaluate the time necessary to produce a column density of 3×10^{17} molecules cm⁻². This time scale results to be, at Europa, on the order of $1.0\text{-}1.3 \times 10^4$ yrs. This time has to be compared with those of other processes. It turns out that the estimated time scale for the production of carbon dioxide by ion bombardment of water on top of carbonaceous materials ranges between 50 and 2×10^3 yrs (Gomis & Strazzulla 2005). We conclude that although a relevant quantity of CO₂ can be formed by carbon ion implantation, this is not the dominant formation mechanism.

It is trivial to say that more observational data on carbon ion flux distribution at the surfaces of icy moons are needed as well as their relationship with the local CO₂ abundance.

Acknowledgements. This research has been partially supported by the European COST Action CM 0805: The Chemical Cosmos: Understanding Chemistry in Astronomical Environments. We acknowledge CAPES, COFECUB, the Basse-Normandie region for financial support. We thank S. Guillous, T. Been, F. Noury and J. M. Ramillon for their technical assistance.

References

- Arnaud, A., Aumayr, F., Echenique, P. M., et al. 1997, *Surf. Sci. Rep.*, 27, 113
 Bagenal, F., Dowling, T. E., & McKinnon, W. B. 2007, *Jupiter: the planet, satellites and magnetosphere*, Vol. 1 (Cambridge Univ Pr)
 Baragiola, R. A. 2003, *Planet. Space. Sci.*, 51, 953
 Baratta, G. A., Fulvio, D., Garozzo, M., et al. 2012, *Memorie della Societa Astronomica Italiana Supplementi*, 20, 94
 Baratta, G. A., Leto, G., & Palumbo, M. E. 2002, *A&A*, 384, 343
 Bennett, C., Hama, T., Kim, Y., Kawasaki, M., & Kaiser, R. 2011, *ApJ*, 727, 27
 Bibring, J. P. & Rocard, F. 1984, *Adv. Space. Res.*, 4, 103
 Carlson, R. W. 2001, in *Division for Planetary Sciences Meeting Abstracts*, Vol. 33, 1125
 Carlson, R. W., Anderson, M. S., Johnson, R. E., et al. 1999, *Science*, 283, 2062
 Carlson, R. W., Smythe, W. D., Lopes-Gautier, R. M. C., et al. 1997, *Geophys. Res. Lett.*, 24, 2479
 Cohen, C. M. S., Stone, E. C., & Selesnick, R. S. 2001, *J. Geophys. Res.*, 106, 29871
 Cooper, J. F., Johnson, R. E., Mauk, B. H., Garrett, H. B., & Gehrels, N. 2001, *Icarus*, 149, 133
 Dalton, J. B., Cruikshank, D. P., Stephan, K., et al. 2010, *Space. Sci. Rev.*, 153, 113
 Dalton, J. B., Shirley, J. H., Cassidy, T., Paranicas, C. J., & Kamp, L. W. 2011, in *EPSC-DPS Joint Meeting*, 649
 Dawes, A., Hunniford, A., Holtom, P. D., et al. 2007, *Phys. Chem. Chem. Phys.*, 9, 2886
 Gerakines, P., Schutte, W., & Ehrenfreund, P. 1996, *A&A*, 312, 289
 Gerakines, P. A., Schutte, W. A., Greenberg, J. M., & van Dishoeck, E. F. 1995, *A&A*, 296, 810
 Gomis, O. & Strazzulla, G. 2005, *Icarus*, 177, 570
 Hamilton, D. C., Gloeckler, G., Krimigis, S. M., & Lanzerotti, L. J. 1981, *J. Geophys. Res.*, 86, 8301
 Herrmann, R., Cocke, C. L., Ullrich, J., et al. 1994, *Phys. Rev. A*, 50, 1435
 Hunniford, C. A., Dawes, A., Fulvio, D., et al. 2009, in *Journal of Physics: Conference Series*, Vol. 163, IOP Publishing, 012078
 Ioppolo, S., Palumbo, M. E., Baratta, G. A., & Mennella, V. 2009, *A&A*, 493, 1017
 Jamieson, C. S., Mebel, A. M., & Kaiser, R. I. 2006, *Astrophys. J. Suppl. S.*, 163, 184
 Johnson, R. E. 1990, *Energetic charged-particle interactions with atmospheres and surfaces*, Vol. 19 (Springer-Verlag Heidelberg)
 Johnson, R. E., Carlson, R. W., Cooper, J. F., et al. 2004, *Jupiter: The Planet, Satellites and Magnetosphere*, 485
 McCord, T. B., Carlson, R. W., Smythe, W. D., et al. 1997, *Science*, 278, 271
 McCord, T. B., Hansen, G. B., Clark, R. N., et al. 1998, *J. Geophys. Res.*, 103, 8603
 McFadden, L. A. A., Weissman, P. R., & Johnson, T. V. 2007, *Encyclopedia of the solar system* (Academic)
 Mennella, V., Palumbo, M. E., & Baratta, G. A. 2004, *ApJ*, 615, 1073
 Moore, J. M., Asphaug, E., Morrison, D., et al. 1999, *Icarus*, 140, 294
 Noll, K. S., Johnson, R. E., McGrath, M. A., & Caldwell, J. J. 1997, *Geophys. Res. Lett.*, 24, 1139
 Noll, K. S., Weaver, H. A., & Gonnella, A. M. 1995, *J. Geophys. Res.*, 100, 19
 Palumbo, M. E. 2005, in *Journal of Physics: Conference Series*, Vol. 6, IOP Publishing, 211
 Palumbo, M. E. 2006, *A&A*, 453, 903
 Palumbo, M. E., Baratta, G. A., Brucato, J. R., et al. 1998, *A&A*, 334, 247
 Palumbo, M. E. & Strazzulla, G. 2003, *Can. J. Phys.*, 1, 217
 Pilling, S., Andrade, D. P. P., da Silveira, E. F., et al. in press, *Mon. Not. R. Astron. Soc.*
 Pilling, S., Duarte, E. S., Domaracka, A., et al. 2011, *Phys. Chem. Chem. Phys.*, 13, 15755
 Pilling, S., Seperuelo Duarte, E., Domaracka, A., et al. 2010, *A&A*, 523, A77
 Raut, U., Fulvio, D., Loeffler, M. J., & Baragiola, R. A. 2012, *The Astrophysical Journal*, 752, 159
 Sandford, S. A. & Allamandola, L. J. 1990, *Icarus*, 87, 188
 Seperuelo Duarte, E., Domaracka, A., Boduch, P., et al. 2010, *A&A*, 512, A71
 Spencer, J. R., Pearl, J. C., Segura, M., et al. 2006, *Science*, 311, 1401
 Strazzulla, G. 2011, *Nucl. Instrum. Meth. B*, 269, 842
 Strazzulla, G., Baratta, G. A., Leto, G., & Gomis, O. 2007, *Icarus*, 192, 623
 Strazzulla, G., Leto, G., Gomis, O., & Satorre, M. A. 2003, *Icarus*, 164, 163
 Strazzulla, G., Leto, G., Spinella, F., & Gomis, O. 2005, *Astrobiology*, 5, 612
 Wada, A., Mochizuki, N., & Hiraoka, K. 2006, *ApJ*, 644, 300
 Watanabe, N., Mouri, O., Nagaoka, A., et al. 2007, *ApJ*, 668, 1001
 Yamada, H. & Person, W. B. 1964, *J. Chem. Phys.*, 41, 2478
 Ziegler, J. F., Biersack, J. P., & Ziegler, M. D. 2008, *Stopping and range of ions in matter* (SRIM Co.)