

# Open Research Online

---

The Open University's repository of research publications and other research outputs

## The organisation and functions of local $\text{Ca}^{2+}$ signals

### Journal Item

How to cite:

Bootman, Martin D.; Lipp, Peter and Berridge, Michael J. (2001). The organisation and functions of local  $\text{Ca}^{2+}$  signals. *Journal of cell science*, 114 pp. 2213–2222.

For guidance on citations see [FAQs](#).

© 2001 The Company of Biologists Ltd

Version: Version of Record

Link(s) to article on publisher's website:  
<http://jcs.biologists.org/content/114/12/2213.long>

---

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's [data policy](#) on reuse of materials please consult the policies page.

---

[oro.open.ac.uk](http://oro.open.ac.uk)

# The organisation and functions of local Ca<sup>2+</sup> signals

Martin D. Bootman\*, Peter Lipp and Michael J. Berridge

Laboratory of Molecular Signalling, The Babraham Institute, Babraham, Cambridge, CB2 4AT, UK

\*Author for correspondence (e-mail: martin.bootman@bbsrc.ac.uk)

Journal of Cell Science 114, 2213-2222 © The Company of Biologists Ltd

## Summary

Calcium (Ca<sup>2+</sup>) is a ubiquitous intracellular messenger, controlling a diverse range of cellular processes, such as gene transcription, muscle contraction and cell proliferation. The ability of a simple ion such as Ca<sup>2+</sup> to play a pivotal role in cell biology results from the facility that cells have to shape Ca<sup>2+</sup> signals in space, time and amplitude. To generate and interpret the variety of observed Ca<sup>2+</sup> signals, different cell types employ components selected from a Ca<sup>2+</sup> signalling 'toolkit', which comprises an array of homeostatic and sensory

mechanisms. By mixing and matching components from the toolkit, cells can obtain Ca<sup>2+</sup> signals that suit their physiology. Recent studies have demonstrated the importance of local Ca<sup>2+</sup> signals in defining the specificity of the interaction of Ca<sup>2+</sup> with its targets. Furthermore, local Ca<sup>2+</sup> signals are the triggers and building blocks for larger global signals that propagate throughout cells.

Key words: Calcium, Signalling, Channel

## Introduction

Over the past decade, there has been an increasing awareness of local interactions between Ca<sup>2+</sup> and its target molecules. The term 'local' has been variously used to describe microdomains of Ca<sup>2+</sup> (dimensions ~10-100 nm) at the mouth of channels or larger subcellular Ca<sup>2+</sup> signals spanning several micrometers. All of these signals are local in the sense that they are spatially restricted rises in Ca<sup>2+</sup> concentration within a cell. However, the sources and targets of the Ca<sup>2+</sup> signals are usually very different.

Most cells utilise Ca<sup>2+</sup> influx from the extracellular space and Ca<sup>2+</sup> release from intracellular stores to generate intracellular signals. When activated, both Ca<sup>2+</sup> entry and Ca<sup>2+</sup> release channels can give rise to brief pulses of Ca<sup>2+</sup> that form a small plume around the mouth of the channel before diffusing into the cytoplasm (reviewed by Neher, 1998). Such signals can remain localised and activate effectors within the immediate vicinity of the channels, they can recruit effectors (for example, see Maasch et al., 2000) or they can be summated to yield global increases that propagate throughout and between cells. Local Ca<sup>2+</sup> signals therefore serve to activate specific targets before diffusion or regenerative mechanisms spread the Ca<sup>2+</sup> across a cell.

## Non-excitable cells

### Ca<sup>2+</sup> puffs

A diverse array of local Ca<sup>2+</sup> signals have been visualised in electrically non-excitable cells (Bootman, 1996). In addition, spatially restricted Ca<sup>2+</sup> signals below the resolution of visible imaging techniques have been inferred from the activity of Ca<sup>2+</sup>-dependent processes. One type of local Ca<sup>2+</sup> signal that appears to operate in many, if not all, electrically non-excitable cells is 'Ca<sup>2+</sup> puffs' (Fig. 1). These elementary events have amplitudes typically ranging from ~50-600 nM, a spatial spread of ~6 µm and a total duration of ~1 second. These Ca<sup>2+</sup>


puff dimensions indicate that there are <100 such sites in small somatic cells. Such events were first observed in *Xenopus* oocytes (for example, see Yao et al., 1995) but have subsequently been observed in HeLa cells (Bootman et al., 1997a), PC12 cells (Reber and Schindelholtz, 1996; Koizumi et al., 1999) and endothelial cells (Hüser and Blatter, 1997). The non-stereotypic nature of Ca<sup>2+</sup> puffs indicates that they probably arise from sites containing variable numbers of inositol 1,4,5-trisphosphate receptors (Ins(1,4,5)P<sub>3</sub>Rs; Sun et al., 1998; Thomas et al., 1998). The temporally and spatially coordinated recruitment of Ca<sup>2+</sup> puffs is responsible for the generation of repetitive Ca<sup>2+</sup> waves and oscillations observed during hormonal stimulation (Bootman et al., 1997b; Marchant et al., 1999; reviewed by Berridge, 1997). Although different non-excitable cell types give rise to distinct global Ca<sup>2+</sup> spikes, similar Ca<sup>2+</sup> puffs can be recorded from various cell lines expressing different combinations of the three Ins(1,4,5)P<sub>3</sub>R isoforms (Tovey et al., 2000). This suggests that cell-specific recruitment of a generic elementary signal underlies different global signals.

In HeLa cells (Thomas et al., 2000) and *Xenopus* oocytes (Marchant and Parker, 2001), Ca<sup>2+</sup> puff sites that have a higher sensitivity to Ins(1,4,5)P<sub>3</sub> consistently trigger Ca<sup>2+</sup> waves (Fig. 1). What gives these pacemaking Ca<sup>2+</sup> puff sites their enhanced sensitivity is unclear. In the case of somatic cells, the pacemaker sites tend to be distributed in a perinuclear region (Lipp et al., 1997), which raises the possibility that they can send Ca<sup>2+</sup> signals specifically into the nucleus.

### Apical Ca<sup>2+</sup> spikes in pancreatic acinar cells

Another well-known local Ca<sup>2+</sup> signal occurs in the apical region of secretory cells such as pancreatic acinar cells (reviewed by Petersen et al., 1999). In common with the pacemaker sites described above, the Ins(1,4,5)P<sub>3</sub>Rs that underlie the apical Ca<sup>2+</sup> spikes are distinguished by a heightened sensitivity to Ins(1,4,5)P<sub>3</sub> (Fogarty et al., 2000).

**Fig. 1.**  $\text{Ca}^{2+}$  puffs in a HeLa cell. The black and red traces in A show the onset of a  $\text{Ca}^{2+}$  wave in a single histamine-stimulated HeLa cell. The  $\text{Ca}^{2+}$  puffs are visible prior to the  $\text{Ca}^{2+}$  wave. The spatial profile of a  $\text{Ca}^{2+}$  puff is indicated by the surface plot in B.  $\text{Ca}^{2+}$  concentration is coded by the height and colour of the surface. The black and red traces in A were obtained by averaging the  $\text{Ca}^{2+}$  concentration over the regions marked by the correspondingly coloured circles on the inset cell image. Modified figure reproduced, with permission, from Thomas et al., 2000.



Like  $\text{Ca}^{2+}$  puffs, such apical  $\text{Ca}^{2+}$  spikes probably arise from the coordinated release of  $\text{Ca}^{2+}$  from multiple  $\text{Ca}^{2+}$ -release channels (Kidd et al., 1999). However, their slow rise time ( $>1$  second; Kidd et al., 1999) and large spatial spread ( $\sim 10 \mu\text{m}$ ; Kidd et al., 1999) distinguishes them from the  $\text{Ca}^{2+}$  puffs in HeLa cells and *Xenopus* oocytes. Recent evidence indicates that the apical spikes arise from a stimulus-dependent hierarchical activation of different types of  $\text{Ca}^{2+}$ -release channel (Cancela et al., 1999; Cancela et al., 2000).

When cells are stimulated to low levels, the  $\text{Ca}^{2+}$  spikes stay restricted to the apical pole of the acinar cells, where they can activate ion channels and trigger limited secretion. Greater stimulation causes the  $\text{Ca}^{2+}$  spikes to trigger  $\text{Ca}^{2+}$  waves that propagate towards the basal pole. The restriction of the  $\text{Ca}^{2+}$  signal to the apical pole appears to be due in part to a 'firewall' of mitochondria that buffer  $\text{Ca}^{2+}$  as it diffuses from the apical pole and prevent the activation of ryanodine receptors (RyRs) in the basal pole (Tinel et al., 2000; Straub et al., 2000).

#### Local signalling via $\text{Ca}^{2+}$ entry

In addition to local  $\text{Ca}^{2+}$  signals arising through release from internal stores,  $\text{Ca}^{2+}$  entry can give rise to local signals with specific functions in electrically non-excitable cells. Cooper and colleagues have shown that both  $\text{Ca}^{2+}$ -activated and  $\text{Ca}^{2+}$ -inhibited isozymes of adenylyl cyclase are more sensitive to influx through store-operated  $\text{Ca}^{2+}$  channels (SOCC) than to  $\text{Ca}^{2+}$  release (Fagan et al., 2000a; Fagan et al., 2000b).  $\text{Ca}^{2+}$  influx through voltage-operated  $\text{Ca}^{2+}$  channels (VOCCs; Fagan et al., 2000b) or a novel arachidonic acid-activated  $\text{Ca}^{2+}$  channel (Shuttleworth and Thompson, 1999) does not modulate adenylyl cyclase activity to the same degree as SOCC entry. These data point to a close localisation of SOCC channels and adenylyl cyclases. One way in which this coupling might be achieved is through the co-localisation of SOCCs and target enzymes in specific cellular structures such as caveolae. These 'flask-shaped' invaginations of the plasma membrane have been proposed to contain  $\text{Ca}^{2+}$ -entry channels and  $\text{Ins}(1,4,5)\text{P}_3\text{Rs}$  (reviewed by Isshiki and Anderson, 1999). In addition to regulating adenylyl cyclase, localisation of  $\text{Ca}^{2+}$  entry to caveolae could underlie the specific requirement for  $\text{Ca}^{2+}$  entry for activation of endothelial nitric oxide synthase (Paltauf-Doburzynska et al., 1998; Lin et al., 2000).

#### Mitochondria sense local $\text{Ca}^{2+}$ signals

Sequestration of  $\text{Ca}^{2+}$  signals by mitochondria (by both electrically excitable and non-excitable cells) serves at least two essential purposes: buffering of cytosolic  $\text{Ca}^{2+}$  loads and consequent activation of citric acid enzymes (for example, see


Robb-Gaspers et al., 1998; reviewed by Duchon, 1999). Mitochondria sequester  $\text{Ca}^{2+}$  through a low-affinity, high-speed uniporter powered by the mitochondrial membrane potential. The in vitro affinity of the uniporter for  $\text{Ca}^{2+}$  is  $\sim 10 \mu\text{M}$ , which is lower than the global  $\text{Ca}^{2+}$  concentrations observed in vivo during physiological responses. However, mitochondria have been shown to sense and modulate both  $\text{Ca}^{2+}$  entry and  $\text{Ca}^{2+}$  release (for example, see Hoth et al., 1997; Sheppard et al., 1997; Landolfi et al., 1998; Hajnoczky et al., 1999; Collins et al., 2000a). To explain the paradox that mitochondria accumulate  $\text{Ca}^{2+}$  during physiological responses despite the insensitivity of the  $\text{Ca}^{2+}$  uniporter, it has been suggested that mitochondria are located in close proximity to sites of either  $\text{Ca}^{2+}$  release or  $\text{Ca}^{2+}$  entry (for example, see Rizzuto et al., 1998; Csordás et al., 1999). In these locations, mitochondria can respond to  $\text{Ca}^{2+}$  changes within the microdomains around the mouth of  $\text{Ca}^{2+}$  channels.

It is therefore generally thought that microdomains of cytosolic  $\text{Ca}^{2+}$  may be the most efficient way of increasing mitochondrial matrix  $\text{Ca}^{2+}$  concentration. However, several studies have shown that modest cytosolic  $\text{Ca}^{2+}$  increases that do not form microdomains can also substantially increase mitochondrial  $\text{Ca}^{2+}$  levels (for example, see Collins et al., 2000b; Colegrove et al., 2000; Collins et al., 2001). Furthermore, mitochondrial  $\text{Ca}^{2+}$  uptake shows a steep dependence on the cytosolic  $\text{Ca}^{2+}$  concentration and might therefore facilitate its own sequestration (for example, see Colegrove et al., 2000; Collins et al., 2001; reviewed by Gunter et al., 1994).

#### Excitable cells

##### Cardiac and skeletal muscle

In heart and skeletal muscle, release of  $\text{Ca}^{2+}$  from the sarcoplasmic reticulum (SR) by RyRs is the key event linking membrane depolarisation and mechanical activity during



**Fig. 2.**  $\text{Ca}^{2+}$  sparks underlie global  $\text{Ca}^{2+}$  signals in cardiac myocytes. The figure shows a sequence of images of the same ventricular cardiomyocyte taken at 33 millisecond intervals. The left-hand image shows the cell with a low resting  $\text{Ca}^{2+}$  concentration prior to stimulation. The right-hand image shows that 66 milliseconds after electrical stimulation (marked with 'lightening zap' symbol) the cell shows a homogenous  $\text{Ca}^{2+}$  increase. On the rising phase (middle image), the  $\text{Ca}^{2+}$  increase can be seen as many focal  $\text{Ca}^{2+}$  increases ( $\text{Ca}^{2+}$  sparks).

excitation-contraction coupling. RyRs occur in clusters that give rise to localised  $\text{Ca}^{2+}$ -release events denoted ' $\text{Ca}^{2+}$  sparks'. These events are analogous to the  $\text{Ca}^{2+}$  puffs described above, although they are usually faster in onset and decline, and have a more restricted spread ( $\sim 1\text{--}3\ \mu\text{m}$ ). Spatio-temporal recruitment of  $\text{Ca}^{2+}$  sparks underlies the global  $\text{Ca}^{2+}$  signals that subsequently activate myocyte contraction (Fig. 2; reviewed by Cannell and Soeller, 1998; Niggli, 1999). The failure to recruit  $\text{Ca}^{2+}$  sparks appropriately can lead to defective excitation-contraction coupling in cardiac cells (Gomez et al., 1997).

In addition to forming the global  $\text{Ca}^{2+}$  transient underlying contraction,  $\text{Ca}^{2+}$  sparks can also cause depolarisation of cardiac cells and thereby enhance or corrupt the rhythm of the heart. An example of this is in primary pacemaker cells, which maintain the normal heartbeat. The electrical activity of cardiac pacemaking cells in the sino-atrial node depends on the interplay of several different sarcolemmal ion channels. In addition,  $\text{Ca}^{2+}$  release from the SR can also affect the frequency of generating action potentials (Rigg and Terrar, 1996). Hüser and colleagues observed that, following recovery from a previous action potential, there is a gradual increase in cytosolic  $\text{Ca}^{2+}$  concentration until the next depolarisation (Hüser et al., 2000). Imaging this slow  $\text{Ca}^{2+}$  rise revealed that it is due to the summation of infrequent subsarcolemmal  $\text{Ca}^{2+}$  sparks. Essentially, low-voltage-activated T-type  $\text{Ca}^{2+}$  channels provide a trigger  $\text{Ca}^{2+}$ -influx current that evokes  $\text{Ca}^{2+}$  spark activity. In turn, the progressive increase in cytoplasmic  $\text{Ca}^{2+}$  caused by the  $\text{Ca}^{2+}$  sparks promotes electrogenic forward-mode  $\text{Na}^{+}/\text{Ca}^{2+}$  exchange, and the resultant inward current drives the cell towards the threshold for depolarisation (Fig. 3).

A similar situation occurs in atrial myocytes stimulated with hormones that activate phospholipase C (PLC). Incubation of electrically-paced atrial myocytes with hormones such as endothelin 1 or phenylephrine causes the appearance of spontaneous subsarcolemmal  $\text{Ca}^{2+}$  sparks (Lipp et al., 1999a; Lipp et al., 1999b), which are probably due to the activation of  $\text{Ins}(1,4,5)\text{P}_3\text{Rs}$  that co-localise with RyRs in these cells (Lipp et al., 2000). If the  $\text{Ca}^{2+}$  spark frequency is sufficient, a spontaneous action potential can be evoked.

These two examples of  $\text{Ca}^{2+}$  sparks driving myocyte depolarisation are entirely analogous. In both situations, a

small trigger  $\text{Ca}^{2+}$  current (either from T-type  $\text{Ca}^{2+}$  channels or  $\text{Ins}(1,4,5)\text{P}_3\text{Rs}$ ) activates  $\text{Ca}^{2+}$  sparks that lead to enhancement of  $\text{Na}^{+}/\text{Ca}^{2+}$  exchange (Fig. 3). Because of the strategic firing of subsarcolemmal  $\text{Ca}^{2+}$  spark sites, only a few such events may be necessary to drive a cell to the threshold for depolarisation (Hüser et al., 2000). The ability of a few  $\text{Ca}^{2+}$  sparks to enhance cardiac automaticity – that is, to increase the frequency of spontaneous action potentials – has potentially serious implications for the generation of cardiac arrhythmias and sudden heart failure.

The generation of spontaneous action potentials, and therefore a global  $\text{Ca}^{2+}$  response, by the  $\text{Ca}^{2+}$  sparks is analogous to the situation whereby  $\text{Ca}^{2+}$  puffs trigger regenerative  $\text{Ca}^{2+}$  waves in non-excitable cells (see above). Although the mechanisms involved are distinct, in both cases the infrequent firing of a limited number of elementary  $\text{Ca}^{2+}$ -release sites leads to larger regenerative  $\text{Ca}^{2+}$  signals.

### Smooth muscle – STOCs, STICs and STOICs

Another cell type in which local  $\text{Ca}^{2+}$  signals have a clear role is smooth muscle. Paradoxically,  $\text{Ca}^{2+}$  can activate smooth muscle contraction, but it can also inhibit contraction by causing hyperpolarisation of the sarcolemma (reviewed by Jaggard et al., 2000). The different effects of  $\text{Ca}^{2+}$  in smooth muscle are essentially determined by the spatial properties of the  $\text{Ca}^{2+}$  signal; global responses induce vasoconstriction by activating  $\text{Ca}^{2+}$ /calmodulin-dependent enzymes, whereas subsarcolemmal  $\text{Ca}^{2+}$  sparks promote vasorelaxation by activating  $\text{Ca}^{2+}$ -dependent plasma membrane ion channels (Nelson et al., 1995).

It is well established that the  $\text{Ca}^{2+}$  sparks in smooth muscle can activate  $\text{K}^{+}$  and  $\text{Cl}^{-}$  conductances, and thus give rise to brief currents known as STOCs (Spontaneous Transient Outward Currents;  $\text{K}^{+}$  current), STICs (Spontaneous Transient Inward Current;  $\text{Cl}^{-}$  current) and STOICs (mixed  $\text{K}^{+}$  and  $\text{Cl}^{-}$  currents). STOCs have been measured in a wide variety of smooth muscle cell types and serve to hyperpolarise the cell membrane by  $\sim 20$  mV. They primarily arise through activation of large-conductance  $\text{Ca}^{2+}$ -activated  $\text{K}^{+}$  channels (BK channels). These  $\text{Ca}^{2+}$ -activated channels have a low sensitivity to cytosolic  $\text{Ca}^{2+}$ , requiring concentrations of  $>1\ \mu\text{M}$  for significant activity (although their actual sensitivity also depends on membrane

potential). It has been proposed that the BK channels are closely apposed to  $\text{Ca}^{2+}$  spark sites and sense rapid step-like  $\text{Ca}^{2+}$  changes during RyR activation (Fig. 4; ZhuGe et al., 2000). Walsh and co-workers have estimated that STOCs reflect the activation of a cluster of ~15 BK channels on the sarcolemma, although the actual BK-channel:RyR stoichiometry may be variable (ZhuGe et al., 2000).

The global  $\text{Ca}^{2+}$  signals that activate smooth muscle cell contraction are largely due to activation of L-type VOCCs. Unlike cardiac muscle, in which  $\text{Ca}^{2+}$  influx also occurs through L-type VOCCs, there is little amplification of the entry signal by  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release (CICR) in most smooth muscle cell types (Jaggar et al., 2000), which means that few subplasmalemmal  $\text{Ca}^{2+}$  spark sites are recruited. Local and global  $\text{Ca}^{2+}$  signalling in smooth muscle cells is therefore segregated in terms of both source and function.

The dramatic effect of a few  $\text{Ca}^{2+}$  sparks on the automaticity of atrial cardiomyocytes (see above) is mimicked in smooth muscle cells, in which the membrane potential can be regulated by STOCs occurring at low frequencies (~1 Hz; Nelson et al., 1995). Pharmacological inhibition of either BK channels or RyRs leads to cellular depolarisation and vasoconstriction. The modulation of vascular tone by vasoactive messengers might be largely due to changes in  $\text{Ca}^{2+}$  spark frequency (Jaggar et al., 2000). In some smooth muscle cell types, only one or a few 'frequent-discharge sites' are responsible for generating the  $\text{Ca}^{2+}$  sparks that activate STOCs (Gordienko et al., 2001). In addition to showing repetitive  $\text{Ca}^{2+}$  sparks, these sites also activate  $\text{Ca}^{2+}$  waves and may be analogous to the pacemaking  $\text{Ca}^{2+}$  puff sites in non-excitable cells (see above).


The BK channels are closely apposed to  $\text{Ca}^{2+}$  spark sites to ensure that they can sense the narrow plume of  $\text{Ca}^{2+}$  around the mouth of the RyRs. In addition, the  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channels express subunits that 'tune' their  $\text{Ca}^{2+}$  sensitivity and enable them to respond to  $\text{Ca}^{2+}$  sparks. BK channels comprise channel-forming  $\alpha$  and auxiliary  $\beta$  subunits. Murine smooth muscle cells lacking the  $\beta$  subunit exhibit reduced sensitivity of BK channels such that they barely respond to the on-going  $\text{Ca}^{2+}$  sparks (Brenner et al., 2000; Pluger et al., 2000). In addition, the  $\beta$ -subunit-deficient animals display an elevated mean arterial blood pressure and cardiac hypertrophy.

$\text{Ca}^{2+}$ -activated  $\text{Cl}^-$  channels are not as widely expressed in smooth muscle types, compared with BK channels. The STICs produced by the  $\text{Ca}^{2+}$ -activated  $\text{Cl}^-$  channels have an essentially opposite effect to that of STOCs in that they serve to depolarise smooth muscle cells (Fig. 4; Jaggar et al., 2000). The STIC channels are approximately one order of magnitude more sensitive to cytosolic  $\text{Ca}^{2+}$  than are STOC channels. This means that their activity more closely follows the decay of the  $\text{Ca}^{2+}$  spark than a STOC. In cells expressing both  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  and  $\text{Cl}^-$  channels, mixed events denoted 'STOICs' have been recorded (ZhuGe et al., 1998). The ambient membrane potential reciprocally controls the flux through the  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  and  $\text{Cl}^-$  channels. This suggests that such STOICs might function to stabilise membrane potential (ZhuGe et al., 1998).

Most smooth muscle cells express both RyRs and  $\text{Ins}(1,4,5)\text{P}_3\text{Rs}$ . The relative contributions of these


two channels to  $\text{Ca}^{2+}$  release is not entirely clear. In ureteric smooth muscle, the elementary events are ' $\text{Ca}^{2+}$  puffs' derived solely from  $\text{Ins}(1,4,5)\text{P}_3\text{Rs}$  (Boittin et al., 2000). In contrast, the  $\text{Ins}(1,4,5)\text{P}_3\text{Rs}$  in pulmonary artery cells do not appear to form clusters that can produce significant elementary events (Janiak et al., 2001). Instead,  $\text{Ins}(1,4,5)\text{P}_3\text{Rs}$  appear to reside within a cluster of RyRs, so that activation of  $\text{Ins}(1,4,5)\text{P}_3\text{Rs}$  gives rise to an event whose characteristics are more similar to those of  $\text{Ca}^{2+}$  sparks (Boittin et al., 1998). A similar crosstalk between  $\text{Ins}(1,4,5)\text{P}_3\text{Rs}$  and RyRs to give 'mixed' elementary events has also been described in neuronal (Koizumi et al., 1999) and cardiac (Lipp et al., 2000) cells.

The  $\text{Ca}^{2+}$  puffs observed during purinergic stimulation of colonic myocytes stimulate STOCs, which suggests that  $\text{Ins}(1,4,5)\text{P}_3\text{Rs}$  and RyRs are functionally equivalent in generating local subplasmalemmal  $\text{Ca}^{2+}$  signals that regulate smooth muscle tone (Bayguinov et al., 2000). However, in contrast to purinergic stimulation of colonic myocytes, purinergic stimulation of cerebral artery smooth muscle cells causes vasoconstriction. In both colonic myocytes and cerebral artery smooth muscle cells, the purinergic agonists are coupled via  $\text{P}_2\text{Y}$  receptors to PLC. The difference between the responses appears to be the relative actions of  $\text{Ins}(1,4,5)\text{P}_3$  and protein kinase C (PKC), the downstream effectors of PLC. In the colonic myocytes,  $\text{Ins}(1,4,5)\text{P}_3$  dominates and increases the frequency of  $\text{Ca}^{2+}$  puffs and associated STOCs, thus causing relaxation (Bayguinov et al., 2000). In cerebral artery smooth muscle cells, PKC-dependent desensitisation of RyRs causes a drop in the frequencies of  $\text{Ca}^{2+}$  sparks and STOCs, leading to contraction (Jaggar and Nelson, 2000). The effect of PLC on smooth muscle activity is therefore cell type dependent. Even more confusing are the observations that within the same smooth muscle cell type different PLC-linked agonists can cause contraction or relaxation because of their differential regulation of STOCs (Bayguinov et al., 2001).



**Fig. 3.** Triggering  $\text{Ca}^{2+}$  sparks by T-type VOCCs or  $\text{Ins}(1,4,5)\text{P}_3\text{Rs}$ . This figure illustrates the activation of spontaneous  $\text{Ca}^{2+}$  sparks by T-type VOCCs in primary cardiac pacemaker cells and  $\text{Ins}(1,4,5)\text{P}_3\text{Rs}$  in atrial cells. RyRs and  $\text{Ins}(1,4,5)\text{P}_3\text{Rs}$  are coloured orange and blue, respectively. The  $\text{Na}^+/\text{Ca}^{2+}$  exchanger is green.

**Fig. 4.** Activation of STOCs and STICs by  $\text{Ca}^{2+}$  sparks in smooth muscle. The  $\text{Ca}^{2+}$  spark originating from RyRs can activate BK channels,  $\text{Cl}^-$  channels or both (producing a STOIC).  $\text{Ca}^{2+}$  entry via L-type VOCCs does not recruit such  $\text{Ca}^{2+}$  spark sites but leads to global  $\text{Ca}^{2+}$  increases that trigger contraction.



## Neurons

### Local $\text{Ca}^{2+}$ signalling in dendritic spines

Subcellular  $\text{Ca}^{2+}$  signalling is the forte of neuronal cells. Because of their often intricate morphologies and the selective distribution of  $\text{Ca}^{2+}$  channels, neurons produce a dazzling array of local  $\text{Ca}^{2+}$  signals. Synaptic input can produce a hierarchy of  $\text{Ca}^{2+}$  signals. These can range from single spines or portions of the dendritic tree, up to responses that invade the soma (reviewed by Denk et al., 1996).

$\text{Ca}^{2+}$  signals within spines are akin to elementary events in other cell types in that they are spatially constrained, short-lived responses that can trigger further propagation of a  $\text{Ca}^{2+}$  wave, depending on the underlying excitability of the cell (reviewed by Berridge, 1998). Spine morphology and number can be dramatically affected by  $\text{Ca}^{2+}$  signals that occur within their tiny volumes (reviewed by Segal, 2001).

$\text{Ca}^{2+}$  signalling within spines underlies changes in synaptic plasticity such as long-term potentiation (LTP) and long-term depression (LTD). Paradoxically,  $\text{Ca}^{2+}$  is implicated in both stimulating and depressing the transmission of nervous signals. It seems that subtle changes in the amplitude, spatial or temporal arrangement of  $\text{Ca}^{2+}$  signals account for the different actions of  $\text{Ca}^{2+}$  in synaptic plasticity (reviewed by Yuste et al., 2000). In particular, the need for coincident signals can dictate the resulting change in synaptic coupling. LTD in cerebellar Purkinje fibres, for example, requires  $\text{Ca}^{2+}$  release from  $\text{Ins}(1,4,5)\text{P}_3\text{R}$  within the spines in conjunction with a large  $\text{Ca}^{2+}$  signal emanating from the activation of VOCCs. Knockout mice that lack type 1  $\text{Ins}(1,4,5)\text{P}_3\text{Rs}$  in their Purkinje cells do not display LTD in response to a standard stimulation protocol (Inoue et al., 1998). Furthermore, Purkinje neurons lacking the motor protein myosin Va, which express  $\text{Ins}(1,4,5)\text{P}_3\text{Rs}$  in dendritic shafts but not in spines, also fail to show LTD (Miyata et al., 2000). Although the  $\text{Ins}(1,4,5)\text{P}_3\text{Rs}$  in the dendrites are  $<1\ \mu\text{m}$  distant from those in the spines, they are not functionally equivalent. A complex interplay between local  $\text{Ca}^{2+}$  signals appears to underlie the polarity and spread of synaptic modification in CA1 hippocampal neurons (Nishiyama et al., 2000). A relay of  $\text{Ca}^{2+}$  between  $\text{Ca}^{2+}$ -influx channels, RyRs and  $\text{Ins}(1,4,5)\text{P}_3\text{Rs}$  within the same microdomain determines whether LTD or LTP is observed following synaptic stimulation (but see Fujii et al., 2000).

The production of  $\text{Ins}(1,4,5)\text{P}_3$  in spines is due to activation of metabotropic glutamate receptors, which are generally located on the periphery of these structures (reviewed by Blackstone and Sheng, 1999). The glutamate receptors are physically linked with  $\text{Ins}(1,4,5)\text{P}_3\text{Rs}$  through members of the Homer protein family (Tu et al., 1998). By virtue of their coiled-coil domains, these proteins can multimerise and link other proteins bearing a proline-rich motif (found in


$\text{Ins}(1,4,5)\text{P}_3\text{Rs}$ , RyRs and other signalling proteins). One member of the Homer family, Homer 1a, does not possess a coiled-coil domain, and can disrupt the coupling between glutamate receptors and  $\text{Ins}(1,4,5)\text{P}_3\text{Rs}$ , which results in reduced  $\text{Ca}^{2+}$ -release signals (Tu et al., 1998). Interestingly, *Homer 1a* is an immediate early gene that is rapidly transcribed after synaptic activity. The induction of this protein therefore provides an activity-dependent mechanism for regulating local  $\text{Ca}^{2+}$  release.

### Local $\text{Ca}^{2+}$ signals underlying repolarisation and hyperpolarisation of neurons

As in smooth muscle cells, neurons express  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channels that repolarise or hyperpolarise the plasma membrane following an action potential. Events analogous to STOCs have been recorded in neuronal cell types. These spontaneous miniature outward currents (SMOCs) are probably also generated by  $\text{Ca}^{2+}$  sparks firing in close proximity to the plasma membrane (Berridge, 1998). Elementary  $\text{Ca}^{2+}$  signals analogous to  $\text{Ca}^{2+}$  sparks and  $\text{Ca}^{2+}$  puffs have also been observed in the neurites of PC12 cells and hippocampal neurons (Koizumi et al., 1999).

A significant function of the  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channels in neuronal cells is to promote repolarisation of the cell after an action potential and to regulate the frequency of firing action potentials. Recent studies have illustrated complex relationships between VOCCs,  $\text{Ca}^{2+}$ -release channels and  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channels in different types of neurons (for a review on the subcellular distribution of VOCCs in neurons, see Caterall, 1998). In the soma of hippocampal pyramidal neurons, N-type VOCCs give a coincident activation of BK channels (Marrion and Tavalin, 1998). The communication between N-type VOCCs and BK channels is not inhibited by the rapid  $\text{Ca}^{2+}$  buffer BAPTA, which indicates that these two types of channel are probably within 30 nm of each other. In contrast, L-type VOCCs are associated with small-conductance  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channels (SK channels) and are estimated to be  $\sim 50\text{--}150\ \text{nm}$  apart (Marrion and Tavalin, 1998). The close coupling of BK channels with N-type VOCCs allows them to be rapidly activated and cause cellular repolarisation. The spatial linkage between SK channels and L-type VOCCs is

**Fig. 5.** Modulation of the after hyperpolarisation (AHP) by  $\text{Ca}^{2+}$  release. The figure illustrates the consequences of single or multiple action potentials on the activation of BK and SK channels in bullfrog sympathetic neurons. See text for details. Modified figure reproduced, with permission, from Akita and Kuba, 2000.



more intriguing. The SK channels are responsible for the 'slow after hyperpolarisation' (sAHP) – a period in which the cell remains hyperpolarised and action potential firing is inhibited. Although the close proximity of L-type VOCCs and SK channels allows communication between them, the AHP develops relatively slowly, and usually when the cell membrane has a negative potential (Berridge, 1998). In this case, the L-type  $\text{Ca}^{2+}$  channels must change their behaviour to be activated at negative membrane potentials, or the SK channels need to rely on local  $\text{Ca}^{2+}$  release from intracellular stores to generate the sAHP.

The involvement of intracellular  $\text{Ca}^{2+}$  release in the generation of AHPs seems clear (Berridge, 1998), but there can be an intricate interplay in the microdomains connecting the  $\text{Ca}^{2+}$  channels and  $\text{K}^{+}$  channels that modulate the membrane potential. In bullfrog sympathetic neurons, N-type VOCCs open during a single action potential and lead to the subsequent activation of BK channels that are responsible for repolarisation (Akita and Kuba, 2000). However, it appears that the  $\text{Ca}^{2+}$  signal from the VOCCs to the BK channels is relayed via RyRs, since inhibitors of CICR depress the rate of repolarisation and the extent of the subsequent AHP (Fig. 5). The situation is more complex when multiple action potentials are evoked, in that the AHP becomes less dependent on BK channels, and instead SK channels are recruited following RyR activation (Akita and Kuba, 2000). Akita and Kuba estimated that BK channels, RyRs and N-type VOCCs are within a ~100 nm domain and that the SK channels are outside this region. The switch from BK to SK channels following multiple action potentials probably reflects the progressive diffusion of  $\text{Ca}^{2+}$  from the N-type VOCCs and RyRs beyond the 100 nm domain (for a detailed discussion see Marchant and Parker, 2000). The SK channels can respond to a more diffuse  $\text{Ca}^{2+}$  signal, since they have a high sensitivity to  $\text{Ca}^{2+}$ , compared with BK channels (Hirschberg et al., 1999).

The regulation of  $\text{Ca}^{2+}$ -sensitive ion channels by local  $\text{Ca}^{2+}$ -release signals in neurons and smooth muscle depends on the close apposition of  $\text{Ca}^{2+}$ -release channels to their targets on the plasma membrane. In these cell types, projections of the ER/SR bring the  $\text{Ca}^{2+}$ -release channels within 10–20 nm of the cell boundary (Berridge, 1998; Jaggar et al., 2000). These structures are akin to the diadic junctions of cardiac muscle, in which RyRs are brought to within ~15 nm of sarcolemma (Cannell and Soeller, 1998). However, a significant difference between cardiac and neuronal/smooth-muscle cells is the degree of regenerativity of  $\text{Ca}^{2+}$  signals following activation of VOCCs. In cardiac myocytes, a large fraction of global  $\text{Ca}^{2+}$  signals is due to  $\text{Ca}^{2+}$  release (i.e. recruitment of  $\text{Ca}^{2+}$  sparks).

By contrast, in neuronal and smooth muscle cells, the degree of regenerative  $\text{Ca}^{2+}$  release is variable, and when it does occur it can lag behind the  $\text{Ca}^{2+}$  influx by hundreds of milliseconds (Berridge, 1998; Jaggar et al., 2000). These data suggest that the coupling between L-type VOCCs and  $\text{Ca}^{2+}$ -release channels in cardiac cells is much tighter than in neuronal and smooth muscle cells. Alternatively, loose coupling might introduce plasticity into the response. In both neuronal and smooth muscle cells, the  $\text{Ca}^{2+}$  stores can retain a 'memory' of previous stimulation, and  $\text{Ca}^{2+}$  release can be modulated by a variety of different factors (for example, see Andreassen and Lambert, 1995; Berridge, 1998; Jaggar et al., 2000).

The effect of T-type VOCCs on excitation-contraction coupling in guinea pig cardiac myocytes functionally resembles the loose coupling observed between VOCCs and  $\text{Ca}^{2+}$ -release channels in neurons and smooth muscle. For the same  $\text{Ca}^{2+}$  current, T-type VOCCs activate much less CICR than does L-type VOCCs (Sipido et al., 1998), which suggests that they are not localised within the diadic junctions. However, blockade of the T-type VOCCs does cause a progressive decrease of the amplitude of action-potential-induced  $\text{Ca}^{2+}$  transients (Sipido et al., 1998). It is plausible that these channels are expressed outside the diadic junctions and are not needed for excitation-contraction coupling but function to refill depleted  $\text{Ca}^{2+}$  stores.

#### Local $\text{Ca}^{2+}$ signalling and neurotransmitter release

Local  $\text{Ca}^{2+}$  signals control the release of neurotransmitter in the active zones of pre-synaptic nerve terminals and neuroendocrine cells (Neher, 1998). In rat cortical synaptosomes, release of small synaptic vesicles (SSVs) containing amino acids (glutamate and GABA) and large dense-core vesicles (LDCVs) containing neuropeptide (cholecystokinin) can be activated by N-, P- and Q-type VOCCs. However, these channels are not equivalent in their efficacy of releasing SSVs and LDCVs: P-type VOCCs preferentially release SSVs, whereas LDCVs are more sensitive to  $\text{Ca}^{2+}$  influx through Q-type channels (Leenders et al., 1999). Such selectivity between VOCCs and exocytotic

vesicles is reminiscent of the linkage between VOCCs and BK/SK channels described earlier. The SSVs are released from the active zones of pre-synaptic nerve terminals, where they can be pre-docked and ready for fusion. In these locations, they are ideally placed to sense the high-amplitude transient Ca<sup>2+</sup> microdomains upon activation of VOCCs (for example, see DiGregorio et al., 1999). LDCVs, in contrast, are more distant from the active zone, and their exocytosis requires prolonged depolarisation (Leenders et al., 1999). The significance of the relative distances of these vesicle types from the active zone is that, for the same Ca<sup>2+</sup> signal, there will be a different secretory output. A single action potential, for example, will release SSVs but have little effect on LDCVs.

Ca<sup>2+</sup> release from intracellular stores can also trigger exocytosis of neurotransmitters from pre-synaptic junctions (Emptage et al., 2001). Release of Ca<sup>2+</sup> from stores in presynaptic boutons contributes to the enhancement of neurotransmitter release with repetitive stimulation (Emptage et al., 2001). In addition, spontaneous Ca<sup>2+</sup>-release events, akin to Ca<sup>2+</sup> sparks and puffs, were recorded in synaptic boutons and appear to be responsible for unevoked synaptic signals. The continual release of neurotransmitter by such spontaneous events could cause long-term changes in strength of synaptic communication (Emptage et al., 2001).

Since the profile of a local Ca<sup>2+</sup> signal falls dramatically with distance from the source (Neher, 1998), activation of exocytosis by release of Ca<sup>2+</sup> from internal stores critically relies on the close apposition of elements of the ER/SR and the secretory machinery (reviewed by Tse and Tse, 1999).

### Local Ca<sup>2+</sup> signalling controls neuronal growth

Many neuronal cells display spontaneous Ca<sup>2+</sup> signals necessary for growth, migration and differentiation (for example, see Komuro and Rakic, 1996; Owens and Kriegstein, 1998; Gomez and Spitzer, 2000; reviewed by Spitzer et al., 2000). In particular, it is well established that Ca<sup>2+</sup> regulates the motility and directionality of neuronal growth cones. However, recent studies have shown that it is not always easy to predict the consequence of a rise in Ca<sup>2+</sup> levels on growth cone behaviour. Large rises can cause growth cone collapse, whereas more modest signals can either slow or promote neurite outgrowth (Spitzer et al., 2000). Using localised release of caged Ca<sup>2+</sup>, Zheng showed that growth cones turn into or away from a focal Ca<sup>2+</sup> source (Zheng, 2000). Local gradients of Ca<sup>2+</sup> might also underlie the turning responses of growth cones to netrin 1, a molecule involved in axonal pathfinding in vivo (Hong et al., 2000). Interestingly, such local gradients must be set up by both Ca<sup>2+</sup> influx and Ca<sup>2+</sup> release, since abolition of turning responses occurs only if both of these Ca<sup>2+</sup> sources are removed. Furthermore, whether a Ca<sup>2+</sup> increase evoked attraction or repulsion depended on the basal level of Ca<sup>2+</sup> (Zheng, 2000) and the activity of protein kinase A (Hong et al., 2000). The ability of such local Ca<sup>2+</sup> signals to activate growth cone turning may be due to restricted rearrangement of the cytoskeleton on one side of a growth cone, perhaps by activation of the Ca<sup>2+</sup>-dependent phosphatase calcineurin (Lautermilch and Spitzer, 2000).

Just how the local Ca<sup>2+</sup> signals are generated by growth cones is unclear. One suggestion is that they are due to activation of VOCCs at resting membrane potential by the action of arachidonic acid (Archer et al., 1999). Exogenous

application of arachidonic acid can stimulate axon outgrowth that is blocked by L-type VOCC channel antagonists. Interestingly, in dorsal root ganglia, the Ca<sup>2+</sup> signal resulting from arachidonic-acid-mediated VOCC activation cannot be visualised by imaging, but is blocked by BAPTA, which suggests that it is localised to a diameter of tens of nanometres (Archer et al., 1999).

The finger-like filopodia that project from growth cones can also relay local Ca<sup>2+</sup> signals to regulate the direction of neurite outgrowth. Spitzer and colleagues have shown that the filopodia of *Xenopus* spinal neurons can discriminate between different types of substrate by generating local Ca<sup>2+</sup> signals of varying frequency that are relayed back to the growth cone (Gomez et al., 2001). The neurons turn away from substrates that give higher frequencies of Ca<sup>2+</sup> transients. The impression is that the filopodia act as scouts that sense the environment in front of the growth cone and direct the axon to preferential substrates.

### Ca<sup>2+</sup> effectors mediate local and global Ca<sup>2+</sup> signals

The plasticity of Ca<sup>2+</sup> as an intracellular messenger is almost matched by the versatility of some of its effectors. The ubiquitous Ca<sup>2+</sup>-binding protein calmodulin (CaM), for example, mediates many of the effects of Ca<sup>2+</sup>, and it can similarly work over different temporal and spatial scales (for example, see Török et al., 1998; reviewed by Chin and Means, 2000; Toutenhoofd and Strehler, 2000). Although a large portion of intracellular CaM is tethered, it can redistribute around cells in a Ca<sup>2+</sup>-dependent manner. In particular, CaM has been shown to move by facilitated diffusion into the nucleus following activation of post-synaptic Ca<sup>2+</sup> channels (Deisseroth et al., 1998) or global Ca<sup>2+</sup> waves (Craske et al., 1999). Furthermore, the extent of CaM translocation is determined by the nature of the cellular Ca<sup>2+</sup> signal (Teruel et al., 2000). The example of nuclear CaM translocation in neurons demonstrates how local Ca<sup>2+</sup> signals can have more distant longer-term consequences.

CaM has two pairs of 'EF hand' motifs that can bind one Ca<sup>2+</sup> ion each. The binding of Ca<sup>2+</sup> to the different EF hands may mediate distinct effects. In the case of L- and P/Q-type VOCCs, for example, CaM can mediate both facilitation and inhibition of the Ca<sup>2+</sup> current by binding to a single site on the VOCC (Zuhlke et al., 1999; DeMaria et al., 2001). Although this implicates CaM as an essential component in short-term regulation of Ca<sup>2+</sup> channels, it seems paradoxical that CaM can have opposing effects at a single binding site on its target. Interestingly, in the case of P/Q-type VOCCs, it appears that binding of Ca<sup>2+</sup> to the N-terminal EF hand pair is responsible for inactivation, whereas the C-terminal EF hand pair underlies facilitation (DeMaria et al., 2001). The empirical observation that facilitation precedes inactivation (DeMaria et al., 2001) suggests that the two pairs of EF hands impart their effects in a temporally regulated manner, perhaps by causing different parts of CaM to interact with the Ca<sup>2+</sup> channel. Since the different EF hands display distinct affinities for Ca<sup>2+</sup>, it is plausible they could sense different components of a Ca<sup>2+</sup> signal (i.e. local versus global). Therefore, just as with Ca<sup>2+</sup> itself, CaM can have distinct functions over distances of nanometres up to whole cells.

In addition to its Ca<sup>2+</sup>-regulated activities, CaM is known to bind to Ca<sup>2+</sup> entry and Ca<sup>2+</sup> release channels in its Ca<sup>2+</sup>-free



form (apoCaM) (DeMaria et al., 2001; Rodney et al., 2001). Such binding of apoCaM may be a means of tethering a Ca<sup>2+</sup> sensor for rapid detection of signalling events (DeMaria et al., 2001), or apoCaM may have regulatory effects distinct from those of Ca<sup>2+</sup>-bound CaM (for example, see Cardy and Taylor, 1998).

## Conclusion

The many functions of local Ca<sup>2+</sup> signals illustrate the versatility of Ca<sup>2+</sup> as an intracellular messenger. No doubt, future studies will provide more examples of Ca<sup>2+</sup> acting at the microscopic and nanoscopic levels within cells. The panoply of local Ca<sup>2+</sup> signals and functions arises through the utilisation of a 'Ca<sup>2+</sup> signalling toolkit' (Berridge et al., 1998; Berridge et al., 2000), whereby different cells express particular combinations of proteins that determine the nature and output of the Ca<sup>2+</sup> signal. In addition, it is becoming apparent that many of the proteins involved in producing and sensing Ca<sup>2+</sup> signals are organised into multiprotein complexes, which provide further means of shaping local events. These complexes include receptors that produce Ins(1,4,5)P<sub>3</sub> (see Muallem and Wilkie, 1999), intracellular Ca<sup>2+</sup>-release channels (see Mackrill, 1999) and Ca<sup>2+</sup>-entry channels (see Blackstone and Sheng, 1999). Furthermore, many effectors of Ca<sup>2+</sup> signals are scaffolded to their Ca<sup>2+</sup> source (for example, see Isshiki and Anderson, 1999). Although local Ca<sup>2+</sup> signals can have immediate effects within the vicinity of the channels from which they originate, the effects of such signals on cardiac pacemaking, growth cone directionality, neuronal membrane potential (see above) and gene transcription (reviewed by Bito et al., 1997) illustrate that brief local Ca<sup>2+</sup> signals can have long-term consequences.

## References

- Akita, T. and Kuba, K. (2000). Functional triads consisting of ryanodine receptors, Ca<sup>2+</sup> channels, and Ca<sup>2+</sup>-activated K<sup>+</sup> channels in bullfrog sympathetic neurons – plastic modulation of action potential. *J. Gen. Physiol.* **116**, 697-720.
- Andreasen, M. and Lambert, J. D. C. (1995). The excitability of CA1 pyramidal dendrites is modulated by a local Ca<sup>2+</sup>-dependent K<sup>+</sup>-conductance. *Brain Res.* **698**, 193-203.
- Archer, F. R., Doherty, P., Collins, D. and Bolsover, S. R. (1999). CAMs and FGF cause a local submembrane calcium signal promoting axon outgrowth without a rise in bulk calcium concentration. *Eur. J. Neurosci.* **11**, 3565-3573.
- Bayguinov, O., Hagen, B., Bonev, A. D., Nelson, M. T. and Sanders, K. M. (2000). Intracellular calcium events activated by ATP in murine colonic myocytes. *Am. J. Physiol.* **279**, C126-C135.
- Bayguinov, O., Hagen, B. and Sanders, K. M. (2001). Muscarinic stimulation increases basal Ca<sup>2+</sup> and inhibits spontaneous Ca<sup>2+</sup> transients in murine colonic myocytes. *Am. J. Physiol.* **280**, C689-C700.
- Berridge, M. J. (1997). Elementary and global aspects of calcium signalling. *J. Physiol.* **499**, 291-306.
- Berridge, M. J. (1998). Neuronal calcium signalling. *Neuron* **21**, 13-26.
- Berridge, M. J., Bootman, M. D. and Lipp, P. (1998). Calcium – a life and death signal. *Nature* **395**, 645-648.
- Berridge, M. J., Lipp, P. and Bootman, M. D. (2000). The versatility and universality of calcium signalling. *Nat. Rev. Mol. Cell Biol.* **1**, 11-21.
- Bito, H., Deisseroth, K. and Tsien, R. W. (1997). Ca<sup>2+</sup>-dependent regulation in neuronal gene expression. *Curr. Opin. Neurobiol.* **7**, 419-429.
- Blackstone, C. and Sheng, M. (1999). Protein targeting and calcium signaling microdomains in neuronal cells. *Cell Calcium* **26**, 181-192.
- Boittin, F.-X., Coussin, F., Macrez, N., Mironneau, C. and Mironneau, J. (1998). Inositol 1,4,5-trisphosphate- and ryanodine-sensitive Ca<sup>2+</sup> release channel dependent Ca<sup>2+</sup> signalling in rat portal vein myocytes. *Cell Calcium* **23**, 303-311.
- Boittin, F.-X., Coussin, F., Morel, J.-L., Halet, G., Macrez, N. and Mironneau, J. (2000). Ca<sup>2+</sup> signals mediated by Ins(1,4,5)P<sub>3</sub>-gated channels in rat ureteric myocytes. *Biochem. J.* **349**, 323-332.
- Bootman, M. D. (1996). Hormone-evoked subcellular Ca<sup>2+</sup> signals in HeLa cells. *Cell Calcium* **20**, 97-104.
- Bootman, M. D., Niggli, E., Berridge, M. J. and Lipp, P. (1997a). Imaging the hierarchical calcium signalling system in HeLa cells. *J. Physiol.* **499**, 307-314.
- Bootman, M., Berridge, M. J. and Lipp, P. (1997b). Cooking with calcium; the recipes for composing global signals from elementary events. *Cell* **91**, 367-373.
- Brenner, R., Perez, G. J., Bonev, A. D., Eckman, D. M., Kosek, J. C., Wiler, S. W., Patterson, A. J., Nelson, M. T. and Aldrich, R. W. (2000). Vasoregulation by the  $\beta$ 1 subunit of the Ca<sup>2+</sup>-activated potassium channel. *Nature* **407**, 870-876.
- Cancela, J. M., Churchill, G. C. and Galione, A. (1999). Coordination of agonist-induced Ca<sup>2+</sup>-signalling patterns by NAADP in pancreatic acinar cells. *Nature* **398**, 74-76.
- Cancela, J. M., Gerasimenko, O. V., Gerasimenko, J. V., Tepikin, A. V. and Petersen, O. H. (2000). Two different but converging messenger pathways to intracellular Ca<sup>2+</sup> release: the roles of nicotinic acid adenine dinucleotide phosphate, cyclic ADP-ribose and inositol trisphosphate. *EMBO J.* **19**, 2549-2557.
- Cannell, M. B. and Soeller, C. (1998). Sparks of interest in cardiac excitation-contraction coupling. *Trends Pharmacol. Sci.* **19**, 16-20.
- Cardy, T. J. A. and Taylor, C. W. (1998). A novel role for calmodulin: Ca<sup>2+</sup>-independent inhibition of type-1 inositol trisphosphate receptors. *Biochem. J.* **334**, 447-455.
- Catterall, W. A. (1998). Structure and function of neuronal Ca<sup>2+</sup> channels and their role in neurotransmitter release. *Cell Calcium* **24**, 307-323.
- Chin, D. and Means, A. R. (2000). Calmodulin: a prototypical calcium sensor. *Trends Cell Biol.* **10**, 322-328.
- Colegrove, S. L., Albrecht, M. A. and Friel, D. D. (2000). Quantitative analysis of mitochondrial Ca<sup>2+</sup> uptake and release in sympathetic neurons. *J. Gen. Physiol.* **115**, 371-388.
- Collins, T. J., Lipp, P., Berridge, M. J., Li, W. and Bootman, M. D. (2000a). Inositol 1,4,5-trisphosphate-induced Ca<sup>2+</sup> release is inhibited by mitochondrial depolarisation. *Biochem. J.* **347**, 593-600.
- Collins, T. J., Bootman, M. D., Lipp, P. and Berridge, M. J. (2000b). Mitochondrial sequestration of cytosolic calcium is dependent upon the source and temporal characteristics of the cytosolic calcium signal in HeLa cells. *J. Physiol.* **527**, 75P-76P.
- Collins, T. J., Lipp, P., Berridge, M. J. and Bootman, M. D. (2001). Mitochondrial Ca<sup>2+</sup> uptake depends on the spatial and temporal profile of cytosolic Ca<sup>2+</sup> signals. *J. Biol. Chem.* (in press).
- Craske, M., Takeo, T., Gerasimenko, O., Vaillant, C., Török, K., Petersen, O. H., Tepikin, A. V. (1999). Hormone-induced secretory and nuclear translocation of calmodulin: oscillations of calmodulin concentration with the nucleus as an integrator. *Proc. Natl. Acad. Sci. USA* **96**, 4426-4431.
- Csordás, G., Thomas, A. P. and Hajnóczky, G. (1999). Quasi-synaptic calcium signal transmission between endoplasmic reticulum and mitochondria. *EMBO J.* **18**, 96-108.
- Deisseroth, K., Heist, E. K. and Tsien, R. W. (1998). Translocation of calmodulin to the nucleus supports CREB phosphorylation in hippocampal neurons. *Nature* **392**, 198-202.
- DeMaria, C. D., Soong, T. W., Alseikhan, B. A., Alvania, R. A. and Yue, D. T. (2001). Calmodulin bifurcates the local Ca<sup>2+</sup> signal that modulates P/Q-type Ca channels. *Nature* **411**, 484-489.
- Denk, W., Yuste, R., Svoboda, K. and Tank, D. W. (1996). Imaging Ca<sup>2+</sup> dynamics in dendritic spines. *Curr. Opin. Neurobiol.* **6**, 372-378.
- DiGregorio, D. A., Peskoff, A. and Vergara, J. L. (1999). Measurement of action potential-induced presynaptic calcium domains at a cultured neuromuscular junction. *J. Neurosci.* **19**, 7846-7859.
- Duchen, M. R. (1999). Contributions of mitochondria to animal physiology: from homeostatic sensor to calcium signalling and cell death. *J. Physiol.* **516**, 1-17.
- Emptage, N. J., Reid, C. A. and Fine, A. (2001). Calcium stores in hippocampal synaptic boutons mediate short-term plasticity, store-operated Ca<sup>2+</sup> entry, and spontaneous transmitter release. *Neuron* **29**, 197-208.
- Fagan, K. A., Smith, K. E. and Cooper, D. M. F. (2000a). Regulation of the Ca<sup>2+</sup>-inhibitable adenylyl cyclase type VI by capacitative Ca<sup>2+</sup> entry

- requires localization in cholesterol-rich domains. *J. Biol. Chem.* **275**, 26530-26537.
- Fagan, K. A., Graf, R. A., Tolman, S., Schaak, J. and Cooper, D. M. F.** (2000b). Regulation of Ca<sup>2+</sup>-sensitive adenylyl cyclase in an excitable cell. *J. Biol. Chem.* **275**, 40187-40194.
- Fogarty, K. E., Kidd, J. F., Tuft, D. A. and Thorn, P.** (2000). Mechanisms underlying InsP<sub>3</sub>-evoked Ca<sup>2+</sup> signals in mouse pancreatic acinar cells. *J. Physiol.* **526**, 515-526.
- Fujii, S., Matsumoto, M., Igarashi, K., Kato, H. and Mikoshiba K.** (2000). Synaptic plasticity in hippocampal CA1 neurons of mice lacking type 1 inositol-1,4,5-trisphosphate receptors. *Learn. Mem.* **7**, 312-320.
- Gomez, T. M. and Spitzer, N. C.** (2000). Regulation of growth cone behaviour by calcium: new dynamics to earlier perspectives. *J. Neurobiol.* **44**, 174-183.
- Gomez, A. M., Valdivia, H. H., Cheng, H., Lederer, M. R., Santana, L. F., Cannell, M. B., McCune, S. A., Altschuld, R. A. and Lederer, W. J.** (1997). Defective excitation-contraction coupling in experimental cardiac hypertrophy and heart failure. *Science* **276**, 800-806.
- Gomez, T. M., Robles, E., Poo, M.-M. and Spitzer, N. C.** (2001). Filopodal calcium transients promote substrate-dependent growth cone turning. *Science* **291**, 1983-1987.
- Gordienko, D. V., Greenwood, I. A. and Bolton, T. B.** (2001). Direct visualisation of sarcoplasmic reticulum regions discharging Ca<sup>2+</sup> sparks in vascular myocytes. *Cell Calcium* **29**, 13-28.
- Gunter, T. E., Gunter, K. K., Sheu, S.-S. and Gavin, C. E.** (1994). Mitochondrial calcium transport: physiological and pathological relevance. *Am. J. Physiol.* **267**, C313-C399.
- Hajnoczky, G., Hager, R. and Thomas, A. P.** (1999). Mitochondria suppress local feedback activation of inositol 1,4,5- trisphosphate receptors by Ca<sup>2+</sup>. *J. Biol. Chem.* **274**, 14157-14162.
- Hirschberg, B., Maylie, J., Adelman, J. P. and Marrion, N. V.** (1999). Gating properties of single SK channels in hippocampal CA1 pyramidal neurons. *Biophys. J.* **77**, 1905-1913.
- Hoth, M., Fanger, C. M. and Lewis, R. S.** (1997). Mitochondrial regulation of store-operated calcium signaling in T lymphocytes. *J. Cell Biol.* **137**, 633-648.
- Hong, K., Nishiyama, M., Henley, J., Tessier-Lavigne, M. and Poo, M.-M.** (2000). Calcium signalling in the guidance of nerve growth by netrin-1. *Nature* **403**, 93-98.
- Hüser, J. and Blatter, L. A.** (1997). Elementary events of agonist-induced Ca<sup>2+</sup> release in vascular endothelial cells. *Am. J. Physiol.* **42**, C1775-C1782.
- Hüser, J., Blatter, L. A. and Lipsius, S. L.** (2000). Intracellular Ca<sup>2+</sup> release contributes to automaticity in cat atrial pacemaker cells. *J. Physiol.* **524**, 415-422.
- Inoue, T., Kato, K., Kohda, K. and Mikoshiba, K.** (1998). Type 1 inositol 1,4,5-trisphosphate receptor is required for induction of long-term depression in cerebellar Purkinje neurons. *J. Neurosci.* **18**, 5366-5373.
- Isshiki, M. and Anderson, R. G. W.** (1999). Calcium signal transduction from caveolae. *Cell Calcium* **26**, 201-208.
- Jaggari, J. H. and Nelson, M. T.** (2000). Differential regulation of Ca<sup>2+</sup> sparks and Ca<sup>2+</sup> waves by UTP in rat cerebral artery smooth muscle cells. *Am. J. Physiol.* **279**, C1528-C1539.
- Jaggari, J. H., Porter, V. A., Lederer, W. J. and Nelson, M. T.** (2000). Ca<sup>2+</sup> sparks in smooth muscle. *Am. J. Physiol.* **278**, C235-C256.
- Janiak, R., Wilson, S. M., Montague, S. and Hume, J. R.** (2001). Heterogeneity of calcium stores and elementary release events in canine pulmonary arterial smooth muscle cells. *Am. J. Physiol.* **280**, C22-C33.
- Kidd, J. F., Fogarty, K. E., Tuft, R. A. and Thorn, P.** (1999). The role of Ca<sup>2+</sup> feedback in shaping InsP<sub>3</sub>-evoked Ca<sup>2+</sup> signals in mouse pancreatic acinar cells. *J. Physiol.* **520**, 187-201.
- Koizumi, S., Bootman, M. D., Bobanovic, L. K., Schell, M. J., Berridge, M. J. and Lipp, P.** (1999). Characterisation of elementary Ca<sup>2+</sup> release signals in NGF-differentiated PC12 cells and hippocampal neurones. *Neuron* **22**, 125-137.
- Komuro, H. and Rakic, P.** (1996). Intracellular Ca<sup>2+</sup> fluctuations modulate the rate of neuronal migration. *Neuron* **17**, 275-285.
- Landolfi, B., Curci, S., Debellis, L., Pozzan, T. and Hofer, A. M.** (1998). Ca<sup>2+</sup> homeostasis in the agonist-sensitive internal store: functional interactions between mitochondria and the ER measured in situ in intact cells. *J. Cell Biol.* **142**, 1235-1243.
- Lautermilch, N. J. and Spitzer, N. C.** (2000). Regulation of calcineurin by growth cone calcium waves controls neurite extension. *J. Neurosci.* **20**, 315-325.
- Leenders, A. G. M., Scholten, G., Wiegant, V. M., Lopes Da Silva, F. and Ghijsen, W. E. J. M.** (1999). Activity-dependent neurotransmitter release kinetics: correlation with changes in morphological distributions of small and large vesicles in central nerve terminals. *Eur. J. Neurosci.* **11**, 4269-4277.
- Lin, S., Fagan, K. A., Li, K. X., Shaul, P. W., Cooper, D. M. F. and Rodman, D. M.** (2000). Sustained endothelial nitric-oxide synthase activation requires capacitative Ca<sup>2+</sup> entry. *J. Biol. Chem.* **275**, 17979-17985.
- Lipp, P., Thomas, D., Berridge, M. J. and Bootman, M. D.** (1997). Nuclear calcium signalling by individual cytoplasmic calcium puffs. *EMBO J.* **16**, 7166-7173.
- Lipp, P., Laine, M., Berridge, M. J. and Bootman, M. D.** (1999a). Enhanced elementary Ca<sup>2+</sup> release caused by InsP<sub>3</sub> mobilising agonists in rat heart. *Pflügers Arch.* **437**, R150.
- Lipp, P., Laine, M., Burrell, K., Berridge, M. J. and Bootman, M. D.** (1999b). Elementary Ca<sup>2+</sup> signals underlie cardiac arrhythmia caused by endothelin. *Pflügers Arch.* **437**, R151.
- Lipp, P., Laine, M., Tovey, S. C., Burrell, K. M., Berridge, M. J., Li, W. and Bootman, M. D.** (2000). Functional InsP<sub>3</sub> receptors that may modulate excitation-contraction coupling in the heart. *Curr. Biol.* **10**, 939-942.
- Maasch, C., Wagner, S., Lindschau, C., Alexander, G., Buchner, K., Gollasch, M., Luft, F. C. and Haller, H.** (2000). Protein kinase C $\alpha$  targeting is regulated by temporal and spatial changes in intracellular free calcium concentration [Ca<sup>2+</sup>]<sub>i</sub>. *FASEB J.* **14**, 1653-1663.
- Mackrill, J. J.** (1999). Protein-protein interactions in intracellular Ca<sup>2+</sup>-release channel function. *Biochem. J.* **337**, 345-361.
- Marchant, J. S. and Parker, I.** (2000). Functional interactions in Ca<sup>2+</sup> signaling over different time and distance scales – Commentary. *J. Gen. Physiol.* **116**, 691-695.
- Marchant, J. S. and Parker, I.** (2001). Role of elementary Ca<sup>2+</sup> puffs in generating repetitive Ca<sup>2+</sup> oscillations. *EMBO J.* **20**, 65-76.
- Marchant, J. S., Callamaras, N. and Parker, I.** (1999). Initiation of IP<sub>3</sub>-mediated Ca<sup>2+</sup> waves in *Xenopus* oocytes. *EMBO J.* **18**, 5285-5299.
- Marrion, N. V. and Tavalin, S. J.** (1998). Selective activation of Ca<sup>2+</sup>-activated K<sup>+</sup> channels by co-localized Ca<sup>2+</sup> channels in hippocampal neurons. *Nature* **395**, 900-905.
- Miyata, M., Finch, E. A., Khiroug, L., Hashimoto, K., Hayasaka, S., Oda, S.-I., Inouye, M., Takagishi, Y., Augustine, G. J. and Kano, M.** (2000). Local calcium release in dendritic spines required or long-term synaptic depression. *Neuron* **28**, 233-244.
- Muallem, S. and Wilkie, T. M.** (1999). G protein-dependent Ca<sup>2+</sup> signaling complexes in polarized cells. *Cell Calcium* **26**, 173-180.
- Neher, E.** (1998). Vesicle pools and Ca<sup>2+</sup> microdomains: new tools for understanding their roles in neurotransmitter release. *Neuron* **20**, 389-399.
- Nelson, M. T., Cheng, H., Rubart, M., Santana, L. F., Bonev, A. D., Knot, H. J. and Lederer, W. J.** (1995). Relaxation of arterial smooth muscle by Ca<sup>2+</sup> sparks. *Science* **270**, 633-637.
- Niggli, E.** (1999). Localized intracellular calcium signaling in muscle: calcium sparks and calcium quarks *Annu. Rev. Physiol.* **61**, 311-335.
- Nishiyama, M., Hong, K., Mikoshiba, K., Poo, M.-M. and Kato, K.** (2000). Calcium stores regulate the polarity and input specificity of synaptic modification. *Nature* **408**, 584-588.
- Owens, D. F. and Kriegstein, A. R.** (1998). Patterns of Ca<sup>2+</sup> fluctuation in precursor cells of the neocortical ventricular zone. *J. Neurosci.* **18**, 5374-5388.
- Paltauf-Doburzynska, J., Posch, K., Paltauf, G. and Graier, W. F.** (1998). Stealth ryanodine-sensitive Ca<sup>2+</sup> release contributes to activity of capacitative Ca<sup>2+</sup> entry and nitric oxide synthase in bovine endothelial cells. *J. Physiol.* **513**, 369-379.
- Petersen, O. H., Burdakov, D. and Tepikin, A. Y.** (1999). Polarity in intracellular calcium signaling. *BioEssays* **21**, 851-860.
- Pluger S., Faulhaber J., Furstenua, M., Lohn, M., Waldschutz, R., Gollasch, M., Haller, H., Luft, F. C., Ehmke, H. and Pongs, O.** (2000). Mice with disrupted BK channel beta 1 subunit gene feature abnormal Ca<sup>2+</sup> spark/STOC coupling and elevated blood pressure. *Circ. Res.* **87**, E53-E60.
- Reber, B. F. X. and Schindelholz, A.** (1996). Detection of a trigger zone of bradykinin-induced fast calcium waves in PC12 neurites. *Pflügers Arch.* **432**, 893-903.
- Rigg, L. and Terrar, D. A.** (1996). Possible role of Ca<sup>2+</sup> release from sarcoplasmic reticulum in pacemaking in guinea-pig sino-atrial. *Exp. Physiol.* **81**, 877-880.
- Rizzuto, R., Pinton, P., Carrington, W., Fay, F. S., Fogarty, K. E., Lifshitz, R. A. and Pozzan, T.** (1998). Close contacts with the endoplasmic reticulum as determinants of mitochondrial Ca<sup>2+</sup> responses. *Science* **280**, 1763-1766.
- Robb-Gaspers, L. D., Burnett, P., Rutter, G. A., Denton, R. M., Rizzuto,**

- R. and Thomas, A. P.** (1998). Integrating cytosolic calcium signals into mitochondrial metabolic responses. *EMBO J.* **17**, 4987-5000.
- Rodney, G. G., Moore, C. P., Williams, B. Y., Zhang, J. Z., Krol, J., Pedersen, S. E. and Hamilton, S. L.** (2001). Calcium binding to calmodulin leads to an N-terminal shift in its binding site on the ryanodine receptor. *J. Biol. Chem.* **276**, 2069-2074.
- Segal, M.** (2001). Rapid plasticity of dendritic spine: hints to possible functions? *Prog. Neurobiol.* **63**, 61-70.
- Sheppard, C. A., Simpson, P. B., Sharp, A. H., Nucifora, F. C., Ross, C. A., Lange, G. D. and Russell, J. T.** (1997). Comparison of type 2 inositol 1,4,5-trisphosphate receptor distribution and subcellular Ca<sup>2+</sup> release sites that support Ca<sup>2+</sup> waves in cultured astrocytes. *J. Neurochem.* **68**, 2317-2327.
- Shuttleworth, T. J. and Thompson, J. L.** (1999). Discriminating between capacitative and arachidonate-activated Ca<sup>2+</sup> entry pathways in HEK293 cells. *J. Biol. Chem.* **274**, 31174-31178.
- Sipido, K. R., Carmeliet, E. and Van de Werf, F.** (1998). T-type Ca<sup>2+</sup> current as a trigger for Ca<sup>2+</sup> release from the sarcoplasmic reticulum in guinea-pig ventricular myocytes. *J. Physiol.* **508**, 439-451.
- Spitzer, N. C., Lautermilch, N. J., Smith, R. D. and Gomez, T. M.** (2000). Coding of neuronal differentiation by calcium transients. *BioEssays* **22**, 811-817.
- Straub, S. V., Giovannucci, D. R. and Yule, D. I.** (2000). Calcium wave propagation in pancreatic acinar cells. *J. Gen. Physiol.* **116**, 547-559.
- Sun, X. P., Callamaras, N., Marchant, J. S. and Parker, I.** (1998). A continuum of InsP<sub>3</sub>-mediated elementary Ca<sup>2+</sup> signalling events in *Xenopus* oocytes. *J. Physiol.* **509**, 67-80.
- Teruel, M. N., Chen, W., Persechini, A. and Meyer, T.** (2000). Differential codes for free Ca<sup>2+</sup>-calmodulin signals in nucleus and cytosol. *Curr. Biol.* **10**, 86-94.
- Thomas, D., Lipp, P., Berridge, M. J. and Bootman, M. D.** (1998). Hormone-stimulated calcium puffs in non-excitabile cells are not stereotypic, but reflect activation of different size channel clusters and variable recruitment of channels within a cluster. *J. Biol. Chem.* **273**, 27130-27136.
- Thomas, D., Lipp, P., Tovey, S. C., Berridge, M. J., Li, W. H., Tsien, R. Y. and Bootman, M. D.** (2000). Microscopic properties of elementary Ca<sup>2+</sup> release sites in non-excitabile cells. *Curr. Biol.* **10**, 8-15.
- Tinel, H., Cancela, J. M., Mogami, H., Gerasimenko, J. V., Gerasimenko, O. V., Tepikin, A. V. and Petersen, O. H.** (2000). Active mitochondria surrounding the pancreatic acinar granule region prevent spreading of inositol trisphosphate-evoked local cytosolic Ca<sup>2+</sup> signals. *EMBO J.* **18**, 4999-5008.
- Torok, K., Wilding, M., Groigno, L., Patel, R. and Whitaker, M.** (1998). Imaging the spatial dynamics of calmodulin activation during mitosis. *Curr. Biol.* **8**, 692-699.
- Toutenhoofd, S. L. and Strehler, E. E.** (2000). The calmodulin multigene family as a unique case of genetic redundancy: multiple levels of regulation to provide spatial and temporal control of calmodulin pools? *Cell Calcium* **28**, 83-96.
- Tovey, S. C., Thomas, D., Lipp, P., Berridge, M. J. and Bootman, M. D.** (2000). Functional redundancy of inositol 1,4,5-trisphosphate receptor isoforms at the level of elementary Ca<sup>2+</sup> release. *J. Physiol.* **527**, 80P.
- Tse, F. W. and Tse, A.** (1999). Regulation of exocytosis via release of Ca<sup>2+</sup> from intracellular stores. *BioEssays* **21**, 861-865.
- Tu, J. C., Xiao, B., Yuan, J. P., Lanahan, A. A., Leoffert, K., Li, M., Linden, D. J. and Worley, P. F.** (1998). Homer binds a novel proline-rich motif and links group 1 metabotropic glutamate receptors with IP<sub>3</sub> receptors. *Neuron* **21**, 717-726.
- Yao, Y., Choi, J. and Parker, I.** (1995). Quantal puffs of intracellular Ca<sup>2+</sup> evoked by inositol trisphosphate in *Xenopus* oocytes. *J. Physiol.* **482**, 533-553.
- Yuste, R., Majewska, A. and Holthoff, K.** (2000). From form to function: calcium compartmentation in dendritic spines. *Nat. Neurosci.* **3**, 653-659.
- Zheng, J. Q.** (2000). Turning of nerve growth cones induced by localized increases in intracellular calcium ions. *Nature* **403**, 89-93.
- ZhuGe, R., Sims, S. M., Tuft, R. A., Fogarty, K. E. and Walsh, J. V.** (1998). Ca<sup>2+</sup> sparks activate K<sup>+</sup> and Cl<sup>-</sup> channels, resulting in spontaneous currents in guinea-pig tracheal myocytes. *J. Physiol.* **513**, 711-718.
- ZhuGe, R., Fogarty, K. E., Tuft, R. A., Lifshitz, L. M., Sayar, K. and Walsh, J. V.** (2000). Dynamics of signaling between Ca<sup>2+</sup> sparks and Ca<sup>2+</sup>-activated K<sup>+</sup> channels studied with a novel image-based method for direct intracellular measurement of ryanodine receptor Ca<sup>2+</sup> current. *J. Gen. Physiol.* **116**, 845-864.
- Zuhlke, R. D., Pitt, G. S., Deisseroth, K., Tsien, R. W. and Reuter, H.** (1999). Calmodulin supports both inactivation and facilitation of L-type calcium channels. *Nature* **399**, 159-162.