



Open Research Online

The Open University's repository of research publications
and other research outputs

Super-rotating jets in a re-analysis of the martian atmosphere

Conference or Workshop Item

How to cite:

Lewis, S. R.; Read, P. L.; Ruan, T. and Montabone, L. (2012). Super-rotating jets in a re-analysis of the martian atmosphere. In: European Planetary Science Congress, 23-28 Sep 2012, Madrid.

For guidance on citations see [FAQs](#).

© 2012 The Authors

Version: Version of Record

Link(s) to article on publisher's website:

<http://meetingorganizer.copernicus.org/EPSC2012/EPSC2012-570-1.pdf>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

Super-rotating jets in a re-analysis of the martian atmosphere

S. R. Lewis (1), P. L. Read (2), T. Ruan (2), and L. Montabone (2,3)

(1) Department of Physical Sciences, The Open University, Walton Hall, Milton Keynes MK7 6AA, UK

(s.r.lewis@open.ac.uk), (2) Atmospheric, Oceanic and Planetary Physics, University of Oxford, Oxford, OX1 3PU, UK, (3) Université Paris VI, Laboratoire de Météorologie Dynamique - Tour 45-55, case courrier 99 – 4, place Jussieu, Paris, France.

Abstract

Strong westerly, prograde jets have been identified in the martian atmosphere between about 10–20 km altitude throughout much of the year in a Mars Global Circulation Model (MGCM) study [2]. The development of data assimilation techniques for Mars [3, 5] now permits the analysis of super-rotation in less highly idealized cases using an atmospheric reanalysis, as would be done for the Earth. This paper reviews recent atmospheric reanalyses, in order to validate previous modeling results, to quantify jet amplitudes and to diagnose possible mechanisms supplying angular momentum to the jets.

1. Introduction

Super-rotation is observed in each of the four substantial atmospheres possessed by solid bodies in the solar system. The slowly rotating planet, Venus, and moon, Titan, both have atmospheres that, at least at some altitudes, rotate much more quickly than does the solid surface underneath. The more rapidly rotating planets, Mars and Earth, exhibit less spectacular global super-rotation, but both can possess prograde jets near the equator which rotate more rapidly than does the equatorial surface.

In each case the detailed mechanism, or combination of mechanisms, which produces the super-rotating jets might vary, but all require longitudinally asymmetric motions, waves or eddies, to transport angular momentum up-gradient into the jets [1, 6]. A model simulation or a re-analysis of observations by data assimilation provides an ideal data set which may be used to quantify the amount of atmospheric super-rotation and to diagnose the relative importance of each eddy forcing mechanism.




Figure 1: Zonal wind (contours) and local super-rotation index, s , (colours) at northern hemisphere autumn equinox ($L_S = 180^\circ$) in the UK MGCM [2]. Only positive values of s are shaded to highlight regions of super-rotation. The dust optical depth is varied from moderate to levels approaching planet-encircling dust storms, such as the $L_S = 185^\circ$ event in 2001 ($\tau_{610\text{Pa}} = 0.6, 1.2$ and 3.0 from top to bottom).

2. Model studies

Idealized Mars model studies [2, 10] have already demonstrated the formation of super-rotating jets under Mars-like conditions. Figure 1 illustrates zonal-mean wind at northern hemisphere equinox from [2] under three different, uniform dust loadings. The local super-rotation index, s , which is shown as a colour-scale in Fig. 1, is a measure of the excess zonal-mean angular momentum about the rotation axis, normalized by the angular momentum of air at rest at the equator [6]. If an atmosphere starts from a rest state, as was the case here, $s > 0$ is not possible through purely zonal-mean processes, such as transport by the mean meridional circulation [1]. The super-rotation in Fig. 1 increases with dust loading and may be attributed to the atmospheric sun-synchronous thermal tides [2], although an almost exactly equal global super-rotation will occur, with jets peaking elsewhere in the model if the thermal tides are artificially suppressed.

3. Data assimilation

The use of data assimilation for Mars [3, 5] now means that the atmospheric super-rotation can be diagnosed from an MGCM reanalysis spanning almost six martian years covering the period for which thermal and dust observations are available from the Thermal Emission Spectrometer on Mars Global Surveyor [7] and the Mars Climate Sounder on Mars Reconnaissance Orbiter [4]. A super-rotating equatorial jet is frequently found in the re-analysis.

4. Summary and conclusions

Local and global super-rotation require the transfer of angular momentum up-gradient with respect to the longitudinal-mean gradient of angular momentum. This transfer must be accomplished by eddy processes, which might include thermal tides, large-scale Kelvin or Rossby waves or small-scale inertio-gravity waves or even turbulent diffusive processes. On Earth, strong tropical convection will mix angular momentum and may suppress the development of clear, local super-rotating jets in the troposphere. Mars also has a deep convective boundary layer that varies strongly with surface height [8, 9] and its impact on the equatorial jet in the re-analysis will be examined.

Acknowledgements

The authors thank the UK Space Agency, the UK Science and Technology Facilities Council and ESA for funding.

References

- [1] Hide, R.: Dynamics of the atmospheres of the major planets with an appendix on the viscous boundary layer at the rigid bounding surface of an electrically-conducting rotating fluid in the presence of a magnetic field, *J. Atmos. Sci.*, Vol. 26, 841–853, 1969.
- [2] Lewis, S. R., and Read, P. L.: Equatorial jets in the dusty martian atmosphere, *J. Geophys. Res.*, Vol. 108 (E4), 5034, pp. 1–15, 2003.
- [3] Lewis, S. R., Read, P. L., Conrath, B. J., Pearl, J. C., and Smith, M. D.: Assimilation of Thermal Emission Spectrometer atmospheric data during the Mars Global Surveyor aerobraking period, *Icarus*, Vol. 192 (2), pp. 327–347, 2007.
- [4] McCleese, D. J., et al.: The structure and dynamics of the martian lower and middle atmosphere as observed by the Mars Climate Sounder: 1. Seasonal variations in zonal mean temperature, dust and water ice aerosols”, *J. Geophys. Res.*, Vol. 115, E12016, pp. 1–16, 2010.
- [5] Montabone, L., Lewis, S. R., Read, P. L., Hinson, D. P., Validation of Martian meteorological data assimilation for MGS/TES using radio occultation measurements, *Icarus* Vol. 185 (1), pp. 113–132, 2006.
- [6] Read, P. L.: Super-rotation and diffusion of axial angular momentum: 1. ‘Speed limits’ for axisymmetric flow in a rotating cylindrical fluid annulus, *Quart. J. Roy. Meteorol. Soc.*, Vol. 112, 231–251, 1986.
- [7] Smith, M. D.: Interannual variability in TES atmospheric observations of Mars during 1999–2003, *Icarus*, Vol. 167 (1), pp. 148–165, 2004.
- [8] Spiga, A., Forget, F., Lewis, S. R., and Hinson, D. P.: Structure and dynamics of the convective boundary layer on Mars as inferred from large-eddy simulations and remote-sensing measurements, *Quart. J. Roy. Meteorol. Soc.*, Vol. 136, pp. 414–428, 2010.
- [9] Spiga, A. and Lewis, S. R.: Martian mesoscale and microscale wind variability of relevance for dust lifting, *Mars*, Vol. 5, pp. 146–158, 2010.
- [10] Wilson, R. J., and Hamilton, K.: Comprehensive model simulation of thermal tides in the martian atmosphere, *J. Atmos. Sci.*, Vol. 53 (9), 1290–1326, 1996.