

Open Research Online

The Open University's repository of research publications and other research outputs

Weathering as controlled by shock processes

Conference or Workshop Item

How to cite:

Steer, Elisabeth; Schwenzer, Susanne; Wright, Ian and Grady, Monica (2015). Weathering as controlled by shock processes. In: 4th UK in Aurora Community Meeting, 15 May 2015, London.

For guidance on citations see [FAQs](#).

© 2015 The Authors

Version: Version of Record

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's [data policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

Weathering as Controlled by Shock Processes

E. D. Steer¹, S. P. Schwenzer¹, I. P. Wright¹, M. M. Grady¹ – ¹Department of Physical Sciences, The Open University, Milton Keynes, Buckinghamshire, UK. MK7 6AA.

Shock processes affect all bodies in the solar system and the effect on rocky targets is profound on both the scale of hundreds of km and on the microscopic scale, as evidenced by mineral textures [1, 2]. We examine the micro-environments of weathering in L6 chondrites, both in terms of textural and chemical alteration; and we relate them to shock created features. Weathering, when mediated by fluid, is sped up or slowed down by the presence of fractures [3]. The most profound difference seen in more shocked samples is the formation of compositionally heterogeneous, polycrystalline sulphides (similar to those found in ALH 84001 by [4]). The shocked sulphides contain micro-fractures, which allow enhanced retention of fluid throughout the crystal. The sulphides also have significant compositional heterogeneity which makes the minerals more vulnerable to weathering [4]. This in turn modifies the altering fluid to a more acidic form. The acid then attacks surrounding silicates. The effect of the acidic fluid is seen through the trace elements of the affected olivine and pyroxene, and a decrease in Mn, Fe and Mg can be seen relative to Si adjacent to weathering sulphides. The estimated pressures needed to create polycrystalline sulphides is 35-60 GPa [5], well within the limits of martian crater formation [6], and therefore is likely to play a part in fluid based crater alteration processes.

[1] Schwenzer, S. P. et al. (2012) *Planetary and Space Science*, 70: 84-95. [2] Rubin, A. E. (2003) *Geochimica et Cosmochimica Acta*, 67: 2695-2709. [3] Barber, D. J. & Scott, E. R. D. (2006) *Meteoritics and Planetary Science*, 41: 643-662. [4] Kwong, Y-T.J. (1993) MEND/NEDEM report 1.32.1. [5] Bennett, M. E., McSween, H. (1996) *Meteoritics and Planetary Science*, 31: 255-264. [6] Abramov, O. et al. (2012) *Icarus*, 218: 906-916.