



Open Research Online

Citation

Bachmair, Sophie; Stahl, Kerstin; Collins, Kevin; Hannaford, Jamie; Acreman, Mike; Svoboda, Mark; Knutson, Cody; Helm Smith, Kelly; Wall, Nicole; Fuchs, Brian; Crossman, Neville and Overton, Ian C. (2016). Drought indicators revisited: the need for a wider consideration of environment and society. *WIREs Water*, 3(4) pp. 516–536.

URL

<https://oro.open.ac.uk/46672/>

License

(CC-BY-NC-ND 4.0)Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Policy

This document has been downloaded from Open Research Online, The Open University's repository of research publications. This version is being made available in accordance with Open Research Online policies available from [Open Research Online \(ORO\) Policies](#)

Versions

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding



Article type: Overview

Article title: Drought indicators revisited: the need for a wider consideration of environment and society

Authors:

Full name and affiliation; email address if corresponding author; any conflicts of interest

| |
|---|
| Sophie Bachmair* [Hydrology, Faculty of Environment and Natural Resources, University of Freiburg, Germany; sophie.bachmair@hydrology.uni-freiburg.de] |
| Kerstin Stahl [Hydrology, Faculty of Environment and Natural Resources, University of Freiburg, Germany] |
| Kevin Collins [Open University, Milton Keynes, United Kingdom] |
| Jamie Hannaford [Centre for Ecology & Hydrology, Wallingford, United Kingdom] |
| Mike Acreman [Centre for Ecology & Hydrology, Wallingford, United Kingdom] |
| Mark Svoboda [National Drought Mitigation Center, University of Nebraska-Lincoln, USA] |
| Cody Knutson [National Drought Mitigation Center, University of Nebraska-Lincoln, USA] |
| Kelly Helm Smith [National Drought Mitigation Center, University of Nebraska-Lincoln, USA] |
| Nicole Wall [National Drought Mitigation Center, University of Nebraska-Lincoln, USA] |
| Brian Fuchs [National Drought Mitigation Center, University of Nebraska-Lincoln, USA] |
| Neville D. Crossman [CSIRO Land and Water, Adelaide, SA, Australia] |
| Ian C. Overton [CSIRO Land and Water, Adelaide, SA, Australia] |

Abstract

[Drought indicators are proliferating, but with little consideration of which are most meaningful for describing drought impacts. A number of recent reviews compare different drought indicators, but

none assess which indicators are actually used in the many operational drought monitoring and early warning efforts, why they were selected, or whether they have been 'ground-truthed', i.e., compared with information representing local drought conditions and/or impacts. Also lacking is a comprehensive assessment of the state of monitoring of drought impacts. To help fill this gap, we combine a review of drought indicators and impacts with a survey of 33 providers of operational drought monitoring and early warning systems from global to regional scales. Despite considerable variety in the indicators used operationally, certain patterns emerge. Both the literature review and the survey reveal that impact monitoring does exist but has rarely been systematized. Efforts to test drought indicators have mostly focused on agricultural drought. Our review points to a current trend towards the design and use of composite indicators, but with limited evaluation of the links between indicators and drought impacts. Overall, we find that much progress has been made both in research and practice on drought indicators, but monitoring and early warning systems are not yet strongly linked with the assessment of wider impacts on the environment and society. To understand drought impacts fully requires a better framing of drought as a coupled dynamic between the environment and society.]

Introduction

[Drought poses a threat to water and food security, to every water-use sector, and thus to livelihoods, in virtually every climate zone. In recent years, a number of major droughts have revealed the vulnerability of even wealthy societies to drought and caused conflicts among water users. Drought can perturb the environment, at least temporarily, with a reduction in ecosystem condition and resilience and a loss of ecosystem services. With climate projections suggesting that droughts will intensify in many regions¹ the magnitude of drought and associated impacts is likely to increase. At the same time, drought is an elusive phenomenon that differs substantially from other natural hazards, making its management a challenging task.² First, it is a slowly developing hazard without a distinct onset and end; second, it is not precisely and universally defined^{3,4}; third, drought is multifaceted, affecting different parts of the hydrological cycle, ecosystems and sectors of society; and fourth, impacts of drought are often non-structural and difficult to quantify or monetize.^{5,6}

While little can be done to prevent low precipitation, drought monitoring and early warning can lessen societal vulnerability by providing more lead-time for responding to drought, planning responses and avoiding a potential crisis situation. A drought early warning system (DEWS)^{7,8} consists of monitoring and early warning components. According to the World Meteorological Organization (WMO) a drought monitoring system should track, assess and report climate and water supply trends and current conditions (e.g., rainfall, reservoirs, impacts, etc.).⁹ An early warning system is the "set of capacities needed to generate and disseminate timely and meaningful warning information to enable individuals, communities and organizations threatened by a hazard to prepare and to act appropriately and in sufficient time to reduce the possibility of harm or loss".^{10(p12)} DEWS are therefore not limited to monitoring and forecasting but may also include further components – for example, capacity to disseminate information via communication networks, or assessment of response and management options alongside the measuring of the current water situation. The design of a DEWS is complicated by the lack of a universal drought definition (see Belal et al.¹¹ for a selection of drought definitions). Generally, two types of drought definitions can be distinguished: conceptual and operational.³ Conceptual definitions aim to explain the general idea. For instance, drought is a "deficiency of precipitation from expected or 'normal' that, when extended over a

season or longer period of time, is insufficient to meet demands".^{12(pA-1)} Operational definitions, in contrast, are targeted towards determining drought onset, severity, and termination for practical applications (e.g., declaration of drought or activation of drought plans for response and mitigation measures).

Operational definitions are typically based on the use of drought indicators. Identifying and selecting appropriate drought indicators for DEWS is complicated by multiple ways of thinking about droughts. Wilhite and Glantz³ differentiate four types of drought – meteorological, agricultural, hydrological, and socio-economic – to describe the propagation of precipitation deficits through the hydrological cycle and its impacts. Over time, several sub-types have been proposed. For example, groundwater drought is treated as an additional category, but it has also been subsumed under hydrological drought (see Wilhite and Glantz³ or Mishra and Singh¹³ on drought types). Another proposed type is ecological drought, although there is a dearth of information on this type despite some ecological metrics being implicit in economic indicators. A vast range of drought indicators has been proposed (>100 according to Lloyd-Hughes⁴) or are used in operational DEWS for monitoring different types of drought in various regions and for different purposes. In addition, novel indicators are frequently published. A recent trend, aimed at capturing a more integrated picture of the drought hazard, has been the design of composite indicators.¹⁴ Such indicators include multiple types of data.

There is little consensus on which indicators are most meaningful for the measurement of drought impacts on society and the environment. A drought impact is “an observable loss or change that occurred at a specific place and time because of drought”.¹⁵ There is a multitude of possible drought impacts (see Wilhite and Glantz³ or Stahl et al.¹⁶ for a comprehensive list). Common classifications differentiate between economic, ecological, and social impacts, or between direct (also termed primary) and indirect (or secondary) impacts, i.e. direct biophysical impacts, and consequences of these impacts.^{3,5,6} A few examples are reduced crop yield, forest dieback, increased mortality of aquatic and terrestrial species, water supply shortages, reduction of hydropower production, impaired navigability of streams, or impacts on human health. Society needs information about when and where drought conditions (expressed by some indicator) translate into impacts. Such information can help people prepare and react proactively, e.g., by developing management and response strategies to mitigate impacts.

Despite several existing reviews of drought indicators (Table 1), what is missing is an integrated review that bridges the gap between scientific developments and current practices in operational DEWS, including a link to drought impacts. To date, there is little knowledge of which indicators are actually used in the many operational DEWS, why they were selected, and whether they have been compared with recorded local drought conditions and/or impacts. Also lacking is a comprehensive assessment of the state of monitoring of drought impacts. The aim of our article is therefore to answer the following research questions:

1. What types of drought indicators exist and which are used in operational DEWS?
2. What is the motivation behind the design of different indicators and their selection for operational use?
3. What are current practices for drought impact monitoring?

4. Have drought indicators been 'ground-truthed', i.e., compared with information representing local drought conditions or impacts?

Our research questions are addressed using two complementary approaches. Firstly, through a synthesis of the published literature on drought indicators and drought impacts, and secondly, through a survey targeted at providers of drought monitoring and early warning information. We close with a summary of current developments and trends regarding drought indicators and their usage in DEWS, and identify knowledge gaps and ways forward.]

[Methods]

[Given the many drought indicators that have been proposed for drought monitoring and the large number of review papers evaluating them, our focus was on synthesizing existing reviews rather than re-reviewing individual indicators. Tables 1 and 2 list the review articles we analyzed. To assess whether drought indicators have been ground-truthed (research question 4), we also collected original research papers dealing with testing of drought indicators, i.e., papers investigating how drought indicators are linked to other indicators or to drought impacts. The papers identified were classified according to the type of linkage (indicator-indicator versus indicator-impact), whether a novel indicator is proposed, and, in case of an indicator-impact linkage, the type of impact variable (Table 3). All papers were selected on the basis of expert knowledge and a snowball search using cross-references and forward citations of highly cited papers.

The online survey focused on (public) providers of drought monitoring and early warning information for public consumption, rather than water utilities or other entities that do monitoring for their own internal business purposes. The survey consisted of open and closed format questions regarding the type of organization and early warning system, reasons for the initiation of the system, use of drought indicators (individual and composite), reasons for the selection of indicators, status of impact monitoring, attempts to ground-truth indicators with impacts, and current indicator developments (the wording of many of the questions can be seen in the figures presented in this study). The invitation to participate in the survey was distributed via email to approximately 70 known DEWS providers and experts; the survey ran from the end of November 2014 until April 2015. We received 33 usable replies (excluding double entries from the same organization). The majority of participating organizations represented either universities/research institutions (48%) or governmental agencies (45%), followed by international (non-governmental) organizations (12%); multiple ticks were possible regarding the type of organization. The geographical coverage of the systems ranged from global to regional scale (global: 36%, continental: 6%, national: 27%, regional: 24%, other (e.g., basin scale): 6%). Several systems integrate different scales, e.g., global and continental, or national and regional scale; the preceding numbers only account for the largest named scale. Continental scale systems cover North and South America, Europe, Africa, and Australia. Most of the national systems cover European countries. Regional systems are distributed globally and either span several countries, or represent systems at sub-national scale. Many of the surveyed DEWS cover developed countries rather than developing countries.]

[What types of drought indicators exist and which are used in operational drought early warning systems?]

[Review of literature]

[Following the terminology used by the US National Drought Mitigation Center (NDMC), a 'drought indicator' can be a single parameter (e.g., rainfall or streamflow at a particular gauging station) or an index combining many kinds of data.¹⁷ An index typically involves computation, e.g., some sort of normalization or combining multiple parameters to produce a numerical index value. In this study we use 'drought indicator' as an overarching term to cover any parameters or indices that are used to characterize and quantify drought. A comprehensive synopsis of existing drought indicators is impractical given the vast (and growing) number of available indicators. Several review papers describe the method for calculation of different indicators and/or evaluate their strengths and limitations. While some of the selected reviews (Table 1) examine a wide range of indicators (e.g., 74 indicators reviewed by Zargar et al.¹⁸), others focus on certain types of indicators, or indicators for a specific region or purpose. Different classification schemes have been used to group drought indicators. A common classification is according to drought type, e.g., precipitation- and temperature-based indicators for meteorological drought, soil moisture or vegetation stress indicators for agricultural drought, and indicators based on streamflow, reservoir or groundwater levels for hydrological drought (see Dai¹⁹ or Zargar et al.¹⁸ for commonly-used indicators falling into these classes). Less common are indicators for socio-economic drought, which have been associated with imbalances in supply and demand of economic goods due to drought, leading to economic and social impacts.²⁰ We will return to socio-economic drought indicators in the section on impact monitoring.

Some indicators do not fit into these classification schemes. One such group is remote sensing indicators, which provide information on several variables, such as land surface temperature, cloud cover, soil moisture, and vegetation status. Remotely sensed vegetation stress indicators such as the Normalized Difference Vegetation Index (NDVI)²¹ and Vegetation Condition Index (VCI)²² are biophysical indicators of a lack of precipitation, but can also be seen as representing drought impacts. Poor vegetation health as a consequence of drought can cause losses in agriculture and forestry systems or a decline in ecosystem condition and thus loss of ecosystem services. Another group not fitting into the drought classification scheme is composite indicators, also termed joint, multivariate, comprehensive, combined, multi-scalar, aggregate, or hybrid indicators.¹⁴ Such indicators provide an integrated picture by including data on multiple types of drought. While commonly agreed composite indicators are by definition multi-dimensional, the boundaries determining which indicators fall into this category are quite fuzzy. We define a composite indicator as a blend of different stand-alone indicators. Examples are the US Drought Monitor (USDM)²³ or the Combined Drought Indicator (CDI).²⁴ For our review, indicators based on more than one input variable, such as the Standardized Precipitation Evaporation Index (SPEI)²⁵ or the Palmer Drought Severity Index (PDSI),²⁶ are considered to be a stand-alone or individual indicator rather than composite. One of the earliest approaches to designing a composite indicator was the USDM, which is based on six key stand-alone indicators and many supplementary indicators. The USDM is unique as it also incorporates local expert knowledge and impact information.²³ Hao and Singh¹⁴ review different composite indicators and discuss the strengths and weaknesses of each method. While composite indicators have the advantage of providing a comprehensive picture of drought, the selection of drought-related variables going into the composite indicator and the methods for combining this information require careful attention and evaluation.¹⁴

Indicators can also be categorized based on the methodology for calculation, the spatial and/or temporal resolution, or the data source. Methods for calculation include percent of normal,

cumulative anomaly, or percentiles of some drought variable. A trend over the last twenty years has been towards standardized indicators such as the Standardized Precipitation Index (SPI).²⁷ Standardized indicators can be thought of as a class of indicators, as the concept of standardization has been extended to nearly all drought-relevant variables: precipitation and evapotranspiration,²⁵ modeled grid cell runoff and routed streamflow,^{28,29} observed streamflow,³⁰ groundwater,³¹ snow melt and rainfall,³² and the PDSI.³³ Standardization has the advantage of consistent interpretability among the standardized indicator family, with the indicator value representing the number of standard deviations from the average cumulative deficit.³⁴ But limitations may arise from the methodology, such as record length, reference period and selection of a probability distribution for model fitting (see Núñez et al.³⁵ for a short review on limitations).

In contrast to the wealth of studies proposing novel indicators or reviewing their strengths and weaknesses, very few studies have investigated which indicators are used in operational DEWS. For the US, the NDMC created a searchable database of indicators used by states, according to their drought plans (<http://drought.unl.edu/Planning/DroughtPlans/StateDroughtPlans/PlansbyTrigger.aspx>). In a review of 33 drought plans for US states, Quiring³⁶ found that reservoir levels and PDSI were most often used, followed by precipitation, streamflow and other indicators. A recent review focusing on drought plans for the western United States revealed that several indicators are used, yet some states rely on certain “primary indicators”.^{37(p96)} For other regions, information on the use of drought indicators for operational purposes may be published in drought plans of water utilities or other entities. As far as we are aware there is no overarching review summarizing this information. Nevertheless, there is review literature that recommends indicators that could be used for operational purposes. For example, in 2009, the WMO held an Inter-Regional Workshop on Indices and Early Warning Systems for Drought in Lincoln, Nebraska, USA, to review drought indicators. The 54 workshop participants from 22 countries recommended that the SPI be used by national meteorological and hydrological services globally to characterize meteorological drought, and that separate recommendations be made for hydrological and agricultural drought.^{38]}

[Survey]

[Survey participants were asked which individual drought indicators they currently use for their DEWS, and whether they provide a composite indicator. Respondents could select between several common indicators per type of drought, or ‘Other’ (followed by a free text field to allow them to specify which) or ‘None’ (Figure 1). Results are discussed by drought type (socio-economic drought is assessed in the section on drought impact monitoring):

- Meteorological drought: The three most often used indicators are SPI, precipitation percentiles, and other indicators based only on precipitation, e.g. accumulated rainfall deficit, percent of normal. The SPEI is used by one third of the surveyed systems. For other indicators, participants named return period estimates for rainfall deficiencies, days without rain, heat related indicators and evaporation. The most common accumulation periods for meteorological indicators are 1, 3, 6, and 12 months (Figure 1). Several systems also provide precipitation-based indicators for accumulation periods up to 72 months. The mixture of short, intermediate, and long accumulation periods accounts for the need to monitor indicators relevant for a wide range of drought impacts with different response times.

- Agricultural drought, Soil moisture and vegetation conditions: soil moisture deficit/anomaly and NDVI are the most commonly used, followed by several other indicators such as the standardized soil moisture index, root stress, and remotely sensed vegetation stress indicators.
- Hydrological drought: Streamflow percentiles and reservoir levels are the most common. Other indicators include stock/farm pond conditions, water allocation levels, and mountain snowpack/snow water equivalent.
- Groundwater drought: 'None' was the most frequent reply, followed by groundwater level percentiles.

Overall, the percentage of 'None' and skipped answers increased moving from meteorological to hydrological, agricultural, and groundwater drought (see Figure 1). This highlights that those variables that are harder to measure and/or model at representative scales are underrepresented indicators in operational DEWS. This is a major gap given the fundamental need for indicators representing drought propagation in different domains of the hydrological cycle and at various spatial and temporal scales.

About 40% of the respondents produce a composite indicator. A variety of indicators go into each composite indicator, commonly covering several types of drought and/or data sources and/or scales. A few examples are the US Drought Monitor, the Multivariate Standardized Drought Index (MSDI), and indicators combining, for example, SPI, soil moisture anomalies, and FAPAR anomalies (Fraction of Absorbed Photosynthetically Active Radiation); PDSI, SPI, and NDVI; or precipitation, temperature, soil moisture, snow pack, and reservoir levels.]

[What is the motivation behind the design of different indicators and their selection for operational use?]

[Review of literature]

[There is relatively little literature on the motivations for indicator design or application. The rationale for proposing novel indicators cited in the literature is to design indicators that are superior to existing ones. For example, they seek to capture missing components of drought, make indicators more robust, increase spatial and/or temporal resolution, or enhance the applicability to certain regions, climate regimes, or sectors. Strategies for improvement include developing techniques for more data sparse areas or merging several indicators as an integrative assessment of drought (Hao and AghaKouchak³⁹; Ma et al.³³; Staudinger et al.³²; Narasimhan and Srinivasan⁴⁰; Zhang and Jia.⁴¹).

Drought monitoring based on a single drought-related variable is insufficient for detecting drought and its diverse consequences.^{13,14} The propagation of drought through the hydrological cycle is a major influence on how drought actually impacts society, and numerous studies have demonstrated that meteorological drought does not always equate to agricultural or hydrological drought (e.g., review by Van Loon⁴²). This lack of equivalence calls into question the purpose of indicators and what they are trying to measure, which in turn highlights a key aspect of monitoring, namely: How is drought understood and how is this understanding represented within current DEWS? With some exceptions (see for example Medd and Chappells⁴³; Vogel et al.⁴⁴; Hayman and Parks⁴⁵) the literature

reports little on the *framing* of drought – i.e. ways in which drought can be conceptualized and interpreted. Different understandings and diverse framings of drought will determine the aims, design, structure and content of DEWS, such as what is excluded and what indicators are required.⁴⁶ The framing of drought depends on the water governance context in a given country or region, including laws and policies related to water rights and on local understanding, as reflected in discourse in the media and among individuals and communities.^{47,48} Given this complexity, it may come as no surprise that indicators tend to represent a precipitation shortfall or hydrological shortage, and exclude social aspects and impacts.

As a result of this complexity, developing and selecting an appropriate indicator for decision making is problematic. An inadequate indicator may lead to a delayed or pre-emptive response,⁴⁹ perhaps with unintended consequences. The worst case is indicators that do not match impacts and experiences occurring on the ground, thus losing credibility with decision makers or the public. Thus, the main motivation for designing composite indicators is to reduce this risk by merging several indicators to obtain an integrated view of drought.]

[Survey]

[We asked the survey participants to rate the importance of reasons behind the selection of individual drought indicators (see Figure 2 for selectable response categories). Data availability and the timeliness of data were the most important reasons followed by simplicity of interpretation, demonstrating that pragmatism drives indicator choice. Good experiences from other organizations, common practices, stakeholder consultation, expert advice, and literature were all rated with intermediate importance. Costs were of relatively low importance, which could be explained by the use of freely available indicators. Other reasons noted were “long standing and well accepted local practices”, “[u]niversality of indicator when applied globally”, and prior investment in the initial development of the indicator, along with reasons linked to the predefined response categories (Figure 2).

An additional factor clearly influencing the choice of indicators is the purpose and targeted audience of the different operational DEWS. About half of the participants’ systems are specifically geared towards drought (45%) and the rest cover drought-related environmental conditions (e.g., water or vegetation status monitoring). More than two-thirds of the systems are not targeted to a specific user group or sector (e.g., agriculture, water supply, and/or the environment). Other participants made a distinction between the use for research, operational managers, decision makers, or the general public. One reply addressed changing demands over time: the DEWS “[u]sed to be geared toward the decision maker. Now [it] has to meet desires of everyone.”

Survey replies regarding the motivation for initiating the DEWS (Figure 3) further explain drought indicator selection. The most frequent reason was ‘Occurrence of (a) severe drought event(s)’, followed by a request by the government/governmental agencies/local authorities or by certain stakeholders (Figure 3). The key role of a severe drought event opening up windows of opportunity for policy and long-term risk management has been reported elsewhere.⁵⁰ It is possible that ad-hoc systems initiated following an event are strongly governed by practical issues and pragmatic choices. This would tie back to data availability and timeliness of data as the main motivations for the selection of indicators.]

[What are the current practices for drought impact monitoring?]

[Review of literature]

[To better understand the current state of drought impact monitoring, the challenges of tracking such information need to be highlighted. A key challenge is the many possible drought impacts (see Table 2 for an overview of review papers covering specific impact types), and differences in how people understand drought and perceive drought impacts. This arises because of different framings of different individuals or organizations within the drought system (see Hayman and Rickards⁴⁵) and their different roles giving rise to diverse experiences of the ‘same’ drought. As stated in the introduction, we define a drought impact as “an observable loss or change that occurred at a specific place and time because of drought”.¹⁵ The US NDMC distinguishes between physical manifestations of drought, such as a lack of precipitation, soil moisture, or water in hydrological systems, and impacts. Hence, low water levels in a river are an indicator of the physical manifestation of drought, not a drought impact. The resulting consequences (e.g., poor water quality, dead fish, and reduced tourism activity) are considered as impacts. The variability of concepts around impacts is discussed in Stahl et al.¹⁶ and Lackstrom et al.⁵¹

Another challenge is that impacts are a function of the vulnerability of the affected area, population, economic sector, or ecosystem, leading to differences in impact types, severity, and time of occurrence for similar drought characteristics.⁵ Impacts may not be visible, or may be disconnected spatially and temporally from the drought event, due to long response times (e.g., tree dieback), and occurrence outside of the main affected area because of the interconnectedness of industries and sectors.^{3,52} Indirect impacts complicate monitoring efforts. For example, reduced crop yield may trigger secondary effects such as food shortages, reduced income for farmers and agribusiness, increased prices for food and timber, unemployment, reduced government tax revenues, increased crime rates and mass migration.⁵ There is further complexity from the multi-causality of impacts. An example is crop damage caused by a combination of weather-related hazards rather than simply rainfall deficit.

These challenges mean drought impacts are not easy to measure, quantify, and/or monetize. Nevertheless, there are several sources of data related directly or indirectly to drought impacts, for example:

- Databases containing textual evidence of drought impacts from reports, newspaper articles, etc., such as the US Drought Impact Reporter;
- Risk management and loss data collected by governments and/or (re)insurance firms;
- Information on the number of affected people and economic damage of large drought events, such as the Emergency Events Database (EM-DAT, www.emdat.be);
- Crop yield statistics;
- Satellite measurements of vegetation stress;
- Databases on wildfire occurrence;
- Information on water use restrictions by water utilities, and;

- Monitoring of water quality and ecological impacts (e.g., the Environment Agency Drought Surveillance Network in England and Wales⁵³).

The above data varies across spatial and temporal resolution and scale, and whether it can be used as real-time information, or only for retrospective analysis. There is no system harmonizing impact data from different sources and/or organizations globally, as available for drought indicators, e.g., through the Global Drought Portal Data (<http://www.drought.gov/gdm/>). For most types of drought impacts (except for agriculture and other insurable risks) no data collection standards exist, resulting in low data availability and/or low consistency of information, e.g., Ding et al.⁵² Furthermore, reporting of impacts is ad-hoc and only when they are felt, rather than continuous monitoring of a system's state, e.g., Dollar et al.⁵⁴; Smith et al., 2014.⁵⁵ This prohibits a systematic assessment of drought impact evolution and the link of impacts to established drought indicators, both going in and coming out of drought. For these reasons, Lackstrom et al.⁵¹ identified impact monitoring as the 'missing piece' of drought early warning. The authors draw their conclusion based on a workshop on drought impact monitoring in the US, discussing opportunities, barriers, and best practices.

One advantage of impact monitoring using narrative accounts is that information on different impact types is collected and archived, e.g., impacts affecting a range of sectors, not only agriculture or water supply.^{5,56} The current benchmark of near real-time monitoring of drought impacts is the US Drought Impact Reporter (DIR; <http://droughtreporter.unl.edu>), which is an online tool for impact collection and reporting. The majority of impact reports in the DIR comes from accounts discovered via an automated media search.⁵ Other sources for populating the DIR are impact reporting by stakeholders, local authorities, the public, or volunteers; or tapping into other reporting systems. Recently, citizen science initiatives like CoCoRaHS (Community Collaborative Rain, Hail and Snow Network) have contributed to impact reporting.⁵⁵

Other regions also have text-based impact inventories, but the systems differ in terms of information channels used, search methods, and operational mode (real-time monitoring versus retrospective search of impacts, systems allowing for public retrieval of information and entries via a web interface versus systems in research mode). The Canadian Agroclimate Impact Reporter (<http://www.agr.gc.ca/atlas/air>) represents a near real-time system similar to the DIR, yet only focuses on agricultural impacts. It relies on a volunteer network of farmers who complete a monthly survey on any climate-related impacts they may have experienced. A database that is strongly modeled after the US DIR is the European Drought Impact report Inventory (EDII; <http://www.geo.uio.no/edc/droughtdb>).¹⁶ However, the EDII was designed as a research database focusing on impacts of past drought events.¹⁶ The majority of EDII entries are from published reports and papers that were assembled by researchers in retrospect, although the database has been recently opened up for public entries. A further European based system is the "Drought in the Media" portal by the European Drought Observatory (<http://edo.jrc.ec.europa.eu/edov2/php/index.php?id=1060>). As the name suggests, it makes use of a media monitor to retrieve information on droughts and related topics world-wide. For Australia, newspaper articles covering water-related issues have been collected within a research initiative.⁵⁷

Despite the potential to capture the diversity of drought impacts, text-based impact monitoring also has challenges.^{5,16,51,58,59} Examples are: (1) media selection or publishing bias of what impacts are reported; (2) identification of relevant search terms for retrieving articles, especially in different

languages, and; (3) the human factor of citizen science initiatives, e.g., fluctuating motivation of volunteers, varying user perception of impacts over time, and personal incentives or disincentives for reporting.^{51,60} Also, the question remains how to use such qualitative data as an indicator or source for testing, e.g., whether and how to quantify or otherwise incorporate it into DEWS. While the USDM already integrates impact information, further possibilities on how to use such data need to be explored.]

[Survey]

[We asked whether data on drought impacts are collected within the participants' systems, and if yes, what kind of data and how. Fifty-five percent of participants collect impact data, and 42% do not (Figure 1). Data on crop damage or yield are collected most frequently, followed by media reports. Other data, such as tree ring records, "satellite products" (presumably on vegetation stress), data on water quality, aquatic ecosystems, or demand, were also named. Respondents also stated impact data collection is fragmented and often ad-hoc during or after an event. Several respondents highlighted that other agencies or organizations collect such data, but the data are not transferred to or distributed/displayed by the DEWS. One respondent noted that data are gathered within individual research projects, yet there is no standardized collection and archiving. Other respondents were vague on whether impact data collection is a sporadic or permanent effort, and whether and how the data are used and/or disseminated. A few responses also demonstrated different understandings of the term 'impact', e.g., referring to low streamflow as an impact.]

[Have drought indicators been 'ground-truthed', i.e. compared with information representing local drought conditions, or impacts?]

[Review of literature]

[We analyzed 70 studies of drought indicator evaluation in the broadest sense, i.e. papers investigating how drought indicators are linked to other indicators, or to drought impacts (Table 3). About two thirds of these studies cross-compared the performance of different indicators. The most common aims of the studies we reviewed were to: (1) evaluate the spatial and temporal consistency of drought identification using multiple indicators; (2) address indicator uncertainties regarding underlying data sources and methodology of calculation (e.g., representativeness of remotely sensed soil moisture versus station-based data; (3) investigate lag times between different types of drought; (4) evaluate indicators for a certain region or application, or; (5) test the utility of a proposed indicator against existing ones. More than one third of the studies on indicator-indicator linkage proposed a novel indicator, either an individual or a composite one.

Less than half of the studies evaluated indicators with impact. Some studies investigated both indicator-indicator, and indicator-impact linkages, hence there is some double counting; remotely sensed vegetation indicators were treated as both indicators and impacts (Table 3). Different types of impact variables were used for indicator evaluation: about half of the studies used crop yield; ca. 20% used vegetation health based on remotely sensed indicators; ca. 20% used text-based impact data from the EDII or the US DIR; and roughly 10% used other, or several impact variables in parallel (e.g., tree rings, forest growth). Additionally, several studies used the US Drought Monitor for ground-truthing indicators. However, since the USDM is not a single impact variable but rather a blended indicator also incorporating some impact information, we considered these studies to fall

into the category of indicator-indicator linkage. Although our proportions are approximate, they demonstrate that the majority of studies rely on either historical crop yield or remotely sensed vegetation stress. Hence, mostly agricultural drought indicators are tested. Evaluation approaches for indicators relevant for societal, economic, and environmental impacts are sparse, highlighting a major gap in our understanding, given the widely discussed multifaceted nature of drought impacts.

Several common themes evolved from the analysis of the 70 studies. First, when a novel indicator is proposed, a common approach is to compare it to existing indicators, often well-known or benchmark indicators for certain regions (e.g., SPI, or the US Drought Monitor for US-wide applications). Additional evaluation with drought impacts is rare, and mostly conducted using crop yield (e.g., Narasimhan and Srinivasan⁴⁰; Potop⁶¹; Rhee et al.⁶²; Sepulcre-Canto et al.⁴⁹). Second, while there are numerous studies cross-comparing the performance of different indicators, there is no standard approach and methods for evaluation, and baseline indicators vary widely. Most studies either compared drought characteristics calculated by means of several indicators, or applied correlation or regression analysis. A small number of studies evaluated indicator performance using techniques for appraising skill, as typically used in forecast evaluation (e.g., Haslinger et al.⁶³; Kumar et al.⁶⁴). Third, most studies focus on some case study region, catchment, or country. Continental or global scale assessments are scarce but include Dai⁶⁵; Vicente-Serrano et al.⁶⁶; Vicente-Serrano et al.⁶⁷

As a result, generalizable information on the performance of indicators and their link to drought impacts is difficult to distill from the many studies reviewed. Some studies report similar findings regarding a specific indicator, such as better performance of the self-calibrated PDSI or Standardized PDI over PDSI,^{33,68} or SPEI outperforming SPI when linked to local-scale hydrological variables or drought impacts.^{58,67,69,70} Apart from this, one commonality seems to be that indicator performance and response times for different types of droughts or impacts are region specific. This was shown by several studies investigating the link between meteorological drought and streamflow,^{63,71–73} SPI and groundwater indicators,^{31,64} and between different indicators and crop yield, vegetation condition, or text-based impact reports.^{58,70,74,75} In addition to regional particularities, indicator performance was found to vary among crops,⁷⁶ and impact types (e.g., impacts on agriculture, water supply, or energy and industry).^{58,70,74} Hence, an overall lesson learned is that a regional and application-specific evaluation is necessary prior to selecting any drought indicator. This makes impact or local-scale water status observations a necessity, perhaps by including evaluation by stakeholders or expert elicitation, as has been practiced in the US (e.g., Steinemann⁷⁷; Steinemann and Cavalcanti⁷⁸; Svoboda et al.²³) and elsewhere.^{79,80}

[Survey]

[Seventy-three percent of respondents reported evaluating individual indicators with impacts (Figure 4). This is a high proportion given that a smaller percentage of respondents reported collecting impact data (55 percent). This suggests that impact data for evaluation are collected in retrospect, for a specific purpose or event, but these data are not continuously available/monitored. Composite indicators were less often evaluated with impact data (54%). Both qualitative (e.g., feedback from stakeholders, testing against local knowledge) and quantitative evaluations (e.g., statistical analysis) were reported. A quantitative evaluation was more common for individual indicators than for the composite ones. Evaluation approaches using stakeholder involvement may partly explain why

indicators have been assessed in terms of their meaning for impacts, even though formal impact monitoring is not as widespread. Examples of qualitative evaluation techniques are “occasional targeted questionnaires” to a wide user community, or “discussions with selected stakeholders on what indicators are used and what indicators are useful, and related to impacts”. The quantitative evaluation involved either: (1) small-scale comparison and/or statistical analysis regarding which indicators best correspond to crop yield, other impact variables, local-scale soil moisture/streamflow/reservoir levels, or indicators such as the USDM, or; (2) assessing the forecasting skill of indicators and/or their ability to predict historical drought events. One comment highlighted the need to balance between indicator skill and value for stakeholders when assessing the overall usefulness of indicators. Where there was no evaluation of indicators, the main reasons given were a lack of personnel and/or time constraints (Figure 4).]

[Current trends, knowledge gaps, and needs for future development]

[Results from survey and literature]

[To gain insight into current developments of indicator usage in operational DEWS, we asked whether survey participants intended to develop new or additional drought indicators in the future, and if so, what and why. Most participants (82%) replied ‘yes’ or ‘possibly’ and some commonalities emerged about trends and reasons for the design of further indicators. Several respondents planned to add existing commonly-used indicators (e.g. indicators representing different types of drought than those currently monitored, or vegetation status from satellite data). In addition, many participants stated their intention to work on the improvement of the currently-used indicators by, for example: (1) refining the methodology or models; (2) using more or better quality input datasets, and; (3) providing higher spatial or temporal resolution, and/or different format datasets (e.g., gridded data). Only a few novel indicators or tools are being developed that go beyond an improvement of methodology. An example is the design of a composite indicator, which appears to receive marked interest by the respondents. Six participants intend to develop such an indicator or explore possibilities in this respect, while 13 of the 33 surveyed systems already provide a composite indicator. Another factor driving indicator development is the requirement to simplify interpretation by users, and better address stakeholders’ needs and/or relevance for certain impacts, as mentioned several times. One participant stated that “[r]ather than developing new indicators (so many are available already), a better goal would be to provide tailored drought information for specific uses”.

Many of these trends are reflected in the current literature. Several participants referred to their own publications regarding indicator evaluation or design of novel indicators. However, research papers specifically addressing how to customize indicators for user needs are rare (but include, for example, Steinemann et al.⁸¹; Steinemann and Cavalcanti⁷⁸). Instead, recent research has: (1) extended the concept of indicator standardization and improved the standardization methodology (e.g., Stagge et al.³⁴; Vicente-Serrano et al.²⁵); (2) assessed multi-indicator drought climatologies at global or continental scales (e.g., Lloyd-Hughes and Saunders⁸²; Spinoni et al.⁸³; Touma et al.⁸⁴) because drought characteristics and derived drought climatologies based on several indicators may differ from those based on a single one^{8,23,78}; (3) designed composite indicators¹⁴, and; (4) utilized modelled and satellite-derived data for DEWS and integrated it into composite indicators to tackle challenges of data scarcity and human resources (e.g., Rhee et al.⁶²; Zhang and Jia⁴¹; Hao et al.⁸⁵), especially in developing countries (e.g., Anderson et al.⁸⁶; Dutra et al.⁸⁷).]

[Knowledge gaps and ways forward in DEWS research and applications]

[The above trends underline the perennial conflict between trying to provide tailored information for particular users or economic sectors, and at the same time providing information for everyone, often through a single composite indicator. On the one hand, blending several indicators may enhance interpretability for users of the systems, since the diverse information from potentially conflicting indicators is streamlined and simplified into one answer. On the other hand, any blending approach obviously involves the subjective choice of indicators, weights, and thresholds for delineation of intensity classes, which may make the interpretation less intuitive or relevant. Since a 'one-size-fits-all' DEWS does not exist, we advocate integrating knowledge from several sources and at different scales without losing detail at smaller spatial scales. The widely publicized USDM, for instance, is not meant to replace information from local water suppliers. Calls for a single source of information aim to simplify what might otherwise be a patchwork of local requirements, but circumstances certainly arise where local utilities' information provision is quite different from a typical single-source message of a larger scale DEWS. Any DEWS should thus seek to integrate rather than reduce complexity to provide meaningful information.

At the same time, we propose that decision-makers in any drought situation adopt an operational drought definition tailored to their own needs. One approach to achieve this, in addition to stakeholder engagement, may be ground-truthing drought indicators with local-scale information on drought conditions or impacts. One aspect of DEWS that is often not explicitly included but that was specifically studied in our survey and review is the connection of drought indicators to impacts. The analysis of research papers dealing with drought indicator evaluation showed that there are numerous studies assessing the performance of (novel) indicators against others, yet evaluation with local-scale water status or impact data other than crop yield is rare. One main reason hindering local-scale indicator testing is a lack of widespread monitoring of (1) variables representing hydrological and groundwater drought, as revealed by the survey, and (2) drought impacts. The scarcity of water status observations, especially for groundwater, reflects the common focus on drought seen through the lens of rainfall and soil moisture that can be easily (remotely) monitored and/or modelled. In general, the lack of hydrological indicators probably reflects a lack of widely accessible, shared hydrometric data at the regional, national or international scales (e.g., Hannah et al.⁸⁸; Viglione et al.⁸⁹) rather than a complete lack of such observations.

While the survey replies showed that efforts are made to collect impact data, the data are rather fragmented and often do not feed directly into the DEWS (see also Lackstrom⁵¹). Wilhite and Glantz (1985)³ found that "most scientific research related to drought has emphasized the physical over the societal aspects of drought".^{3(p119)} Three decades later, impact monitoring to better integrate societal aspects into early warning is still in its infancy, except for a few advanced systems like the US DIR and some initiatives in other countries. Incorporating observer networks may be a way forward to better integrate local knowledge into high-tech drought monitoring. One such example is the previously mentioned CoCoRaHS network in the US, but arguably the greatest potential for observer networks to advance drought monitoring is in developing countries. In data-poor environments with less available DEWS infrastructure, but where mobile communication uptake is high, citizen science can fill in gaps in in-situ networks and provide a link between on-the-ground impacts and large-scale data from earth observation. Examples are projects to utilize cell phones to report drought impacts (e.g. practiced in Ethiopia⁹⁰ and Somalia⁹¹) or the concept of 'Paysan Observateur' (observing

farmers) in Mali.⁹² Collectively, such initiatives offer a way of re-framing drought to include societal impacts, but a task remains on how indicators and impacts can be integrated. Recent initiatives in water management suggest developing social learning systems (see Blackmore and Ison⁹³) may be a way to enable diverse stakeholders to engage with monitoring and early warning systems and practices in order to improve their utility for a wide community of end users.

Although not specifically surveyed and reviewed, ecology and ecosystem services are additional under-represented aspects that emerged during the research and discussion. Although many drought indicators implicitly include terrestrial and freshwater ecosystems – for example, through monitoring soil moisture that is critical for plant growth – explicit indicators of ecological drought are rare. An exception is in the UK where there has been some progress toward developing indicators of ecosystem health for use in drought monitoring⁵⁴. However, one could question whether droughts are ‘harmful’ to ecosystems. Much depends on the human context: if culturally important ecosystems are lost, we may consider the drought to be ecologically and culturally negative. However, there is evidence that droughts eliminate weak members of species and prevent growth of invasive species, and so can have a positive impact on the ecosystem.⁹⁴ Particular challenges to incorporating ecological issues include recovery of ecosystems after drought.⁹⁵ Droughts may negatively affect services to people from ecosystems, such as recreation from boating or fishing in rivers, although these are not strictly ecological indicators. Extended severe droughts, or frequent droughts, may cause an ecosystem to go beyond its threshold tolerances and therefore transition into new ecosystem types⁹⁶. This usually causes a loss of ecosystem services that society relies upon. Although not a specific focus of our review and survey, we note that ecosystem services are absent from current DEWS indicators and appraisal of those indicators.]

Conclusion

[The aim of this work was to revisit drought indicators that are used in drought monitoring and early warning. Our synthesis of literature on drought indicators and impacts together with our survey of providers of drought early warning information tackled questions on the operational use of indicators, the motivation behind the design and selection of different indicators, current practices for drought impact monitoring, and any related ground-truthing of drought indicators with local drought conditions, and/or impacts. In summary, there is considerable variety in the indicators used by operational DEWS, though this variety is not nearly as wide as the range of indicators that have been published. Moreover, certain trends exist: common drought indicators, such as the SPI, are used very widely, and approximately 40% of the surveyed DEWS also provide a composite indicator. The survey confirmed that providers of monitoring and early warning services are constrained by pragmatic considerations such as variables that are easy to measure, and readily available in a timely manner. Perhaps less expected was that more than half of the survey participants collect impact data, although in the literature this has been referred to as the ‘missing piece’ of drought monitoring. A closer look at the replies revealed that very few systematic approaches exist and thus data collection efforts are mostly fragmented. As a result, impact data are not widely used in operational systems, e.g. for ground-truthing indicators – an exception being agricultural drought, where yield or vegetation stress data have been used to test indicators.

The two complementary approaches of literature synthesis and surveying DEWS providers allowed us to develop an integrated picture of the current state of drought indicator research and

practices in operational DEWS. This revealed key knowledge gaps, and particular challenges for future development of DEWS. There is a need for indicators representing drought propagation in different domains of the hydrological cycle and at various spatial and temporal scales, systematic impact data collection for ground-truthing indicators, and better understanding of drought's various economic consequences. Consideration of environmental impacts is still in early stages, and ecosystem services have yet to be integrated into monitoring and early warning frameworks. We also note that the underlying framing of drought – a major consideration in terms of how DEWS are designed and their ascribed purpose – remains largely unexplored in the literature. Citizen science initiatives and other social learning approaches that explore drought framing and DEWS design offer opportunities to explore multiple understandings of drought impacts and improve indicator design and use. While large-scale, big-picture, integrated indicators such as the US Drought Monitor are valuable, we see additional need for further research and development of DEWS systems tracing drought's cascading effects through specific ecological, economic and social contexts.]

Acknowledgements

[We particularly thank all participants of the survey and the GDIS community and the 2014 Pasadena workshop (<http://www.wcrp-climate.org/gdis-wkshp-2014-about>) participants for helping with contact identification. This study is an outcome of the international Belmont Forum/G8HORC's Freshwater Security programme project DrIVER (Drought Impacts: Vulnerability thresholds in monitoring and Early warning Research). We acknowledge funding towards the DrIVER project from the German Research Foundation DFG (project no. STA-632/2-1), NERC (grant number: NE/L010038/1), NSF (ICER-1342949), and CSIRO Land and Water.]

References

1. IPCC. Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation. A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change. Field, C.B., V. Barros, T.F. Stocker, D. Qin, D.J. Dokken, K.L. Ebi, M.D. Mastrandrea, K.J. Mach, G.-K. Plattner, S.K. Allen, M. Tignor PMM, ed. 2012.
2. Wilhite D. Quantification of agricultural drought for effective drought mitigation and preparedness: Key issues and challenges. In Sivakumar MVK, Motha RP, Wilhite DA, Wood DA, eds. *Proceedings of the WMO/UNISDR Expert Group Meeting on Agricultural Drought Indices, 2-4 June 2010, Murcia, Spain*. Geneva: World Meteorological Organization (WMO); 2011:219.
3. Wilhite D, Glantz M. Understanding: the drought phenomenon: the role of definitions. *Water Int* 1985,10(3):111-120. doi:10.1080/02508068508686328
4. Lloyd-Hughes B. The impracticality of a universal drought definition. *Theor Appl Climatol* October 2013. doi:10.1007/s00704-013-1025-7
5. Wilhite DA, Svoboda MD, Hayes MJ. Understanding the complex impacts of drought: A key to enhancing drought mitigation and preparedness. *Water Resour Manag* 2007,21:763-774. doi:10.1007/s11269-006-9076-5
6. Logar I, van den Bergh JCJM. Methods for Assessment of the Costs of Droughts. ConHaz WP5 Report. 2011. conhaz.org/CONHAZ_REPORT_WP05_1_FINAL.pdf
7. Wilhite DA, Buchanan-Smith M. Drought as a Hazard: Understanding the Natural and Social Context. In Wilhite DA, ed. *Drought and Water Crisis: Science, Technology, and Management*

- Issues*. Boca Raton: CRC Press (Taylor & Francis Group); 2005:3-29.
8. Sivakumar MVK, Motha RP, Wilhite DA, Wood DA. Agricultural Drought Indices. In Sivakumar MVK, Motha RP, Wilhite DA, Wood DA, eds. *Proceedings of the WMO/UNISDR Expert Group Meeting Agricultural Drought Indices, 2-4 June 2010, Murcia, Spain*. Geneva: World Meteorological Organization (WMO); 2011:219.
 9. WMO. Drought Monitoring and Early Warning: Concepts, Progress and Future Challenges. 2006. www.wamis.org/agm/pubs/brochures/WMO1006e.pdf
 10. UNISDR. Terminology on Disaster Risk Reduction. 2009.
 11. Belal A-A, El-Ramady HR, Mohamed ES, Saleh AM. Drought risk assessment using remote sensing and GIS techniques. *Arab J Geosci* 2012,7(1):35-53. doi:10.1007/s12517-012-0707-2
 12. Knutson CL, Hayes MJ, Philipps T, Western Drought Coordination Council P and MWG. How to Reduce Drought Risk. 1998.
 13. Mishra AK, Singh VP. A review of drought concepts. *J Hydrol* 2010,391(1):202-216.
 14. Hao Z, Singh VP. Drought characterization from a multivariate perspective: A review. *J Hydrol* 2015,527:668-678. doi:10.1016/j.jhydrol.2015.05.031
 15. NDMC. Drought Impact Reporter Help. <http://public.droughtreporter.unl.edu/help/>. Accessed September 30, 2015
 16. Stahl K, Kohn I, Blauhut V, et al. Impacts of European drought events: insights from an international database of text-based reports. *Nat Hazards Earth Syst Sci Discuss* 2015,3(9):5453-5492. doi:10.5194/nhessd-3-5453-2015
 17. NDMC. Glossary. <http://drought.unl.edu/DroughtBasics/Glossary.aspx>. Accessed September 30, 2015
 18. Zargar A, Sadiq R, Naser B, Khan FI. A review of drought indices. *Environ Rev* 2011,19:333-349. doi:10.1139/A11-013
 19. Dai A. Drought under global warming: a review. *Wiley Interdiscip Rev Clim Chang* 2011,2(1):45-65. doi:10.1002/wcc.81
 20. American Meteorological Society. Statement on Drought. <https://www2.ametsoc.org/ams/index.cfm/about-ams/ams-statements/statements-of-the-ams-in-force/drought/>. Published 2013. Accessed October 8, 2015
 21. Tucker CJ. Red and photographic infrared linear combinations for monitoring vegetation. *Remote Sens Environ* 1979,8(2):127-150. doi:10.1016/0034-4257(79)90013-0
 22. Kogan FN. Global Drought Watch from Space. *Bull Am Meteorol Soc* 1997,78(4):621-636.
 23. Svoboda M, LeComte D, Hayes M, et al. The drought monitor. *Bull Am Meteorol Soc* 2002,83(8):1181-1190.
 24. European Drought Observatory. PRODUCT FACT SHEET: Combined Drought Indicator – EUROPE. 2013. <http://edo.jrc.ec.europa.eu/edov2/php/index.php?id=1101>
 25. Vicente-Serrano SM, Beguería S, López-Moreno JI. A multiscalar drought index sensitive to global warming: the standardized precipitation evapotranspiration index. *J Clim* 2010,23(7):1696-1718.
 26. Palmer WC. Keeping Track of Crop Moisture Conditions, Nationwide: The New Crop Moisture Index. *Weatherwise* 1968,21(4):156-161. doi:10.1080/00431672.1968.9932814

27. McKee TB, Doesken NJ, Kleist J. The relationship of drought frequency and duration to time scales. In *Preprints, 8th Conference on Applied Climatology*. Anaheim, California; 1993:179-184.
28. Shukla S, Wood AW. Use of a standardized runoff index for characterizing hydrologic drought. *Geophys Res Lett* 2008,35:L02405.
29. Vidal J-P, Martin E, Franchistéguy L, Habets F, Soubeyroux J-M, Blanchard M, Baillon M. Multilevel and multiscale drought reanalysis over France with the Safran-Isba-Modcou hydrometeorological suite. *Hydrol Earth Syst Sci* 2010,14(3):459-478. doi:10.5194/hess-14-459-2010
30. Vicente-Serrano SM, López-Moreno JI, Beguería S, Lorenzo-Lacruz J, Azorin-Molina C, Morán-Tejeda E. Accurate computation of a streamflow drought index. *J Hydrol Eng* 2011,17(2):318-332.
31. Bloomfield JP, Marchant BP. Analysis of groundwater drought using a variant of the Standardised Precipitation Index. *Hydrol Earth Syst Sci Discuss* 2013,10(6):7537-7574.
32. Staudinger M, Stahl K, Seibert J. A drought index accounting for snow. *Water Resour Res* September 2014:n/a - n/a. doi:10.1002/2013WR015143
33. Ma M, Ren L, Yuan F, Jiang S, Liu Y, Kong H, Gong L. A new standardized Palmer drought index for hydro-meteorological use. *Hydrol Process* 2014,28(23):5645-5661. doi:10.1002/hyp.10063
34. Stagge JH, Tallaksen LM, Gudmundsson L, Van Loon AF, Stahl K. Candidate Distributions for Climatological Drought Indices (SPI and SPEI). *Int J Climatol* February 2015. doi:10.1002/joc.4267
35. Núñez J, Rivera D, Oyarzún R, Arumí JL. On the use of Standardized Drought Indices under decadal climate variability: Critical assessment and drought policy implications. *J Hydrol* June 2014. doi:10.1016/j.jhydrol.2014.05.038
36. Quiring SM. Developing objective operational definitions for monitoring drought. *J Appl Meteorol Climatol* 2009,48(6):1217-1229.
37. Fontaine MM, Steinemann AC, Hayes MJ. State Drought Programs and Plans: Survey of the Western United States. *Nat Hazards Rev* 2014,15(1):95-99. doi:10.1061/(ASCE)NH.1527-6996.0000094
38. Hayes M, Svoboda M, Wall N, Widhalm M. The Lincoln declaration on drought indices: universal meteorological drought index recommended. *Bull Am Meteorol Soc* 2011,92(4):485-488.
39. Hao Z, AghaKouchak A. A Nonparametric Multivariate Multi-Index Drought Monitoring Framework. *J Hydrometeorol* 2014,15(1):89-101. doi:10.1175/JHM-D-12-0160.1
40. Narasimhan B, Srinivasan R. Development and evaluation of Soil Moisture Deficit Index (SMDI) and Evapotranspiration Deficit Index (ETDI) for agricultural drought monitoring. *Agric For Meteorol* 2005,133(1):69-88.
41. Zhang A, Jia G. Monitoring meteorological drought in semiarid regions using multi-sensor microwave remote sensing data. *Remote Sens Environ* 2013,134:12-23. doi:10.1016/j.rse.2013.02.023
42. Van Loon AF. Hydrological drought explained. *Wiley Interdiscip Rev Water* 2015,2(4):359-392. doi:10.1002/wat2.1085
43. Medd W, Chappells H. Drought, demand and the scale of resilience: challenges for

- interdisciplinarity in practice. *Interdiscip Sci Rev* 2007,32(3):233-248.
doi:10.1179/030801807X211748
44. Vogel C, Koch I, Van Zyl K. "A Persistent Truth"—Reflections on Drought Risk Management in Southern Africa. *Weather Clim Soc* 2010,2(1):9-22. doi:10.1175/2009WCAS1017.1
 45. Hayman P, Rickards L. Drought, Climate Change, Farming, and Science: The Interaction of Four Privileged Topics. In Botterill L, Cockfield G, eds. *Drought, Risk Management and Policy: Decision-Making Under Uncertainty*. Florida: CRC Press; 2013:45-70.
 46. Dessai S, Sims C. Public perception of drought and climate change in southeast England. *Environ Hazards* 2010,9(4):340-357. doi:10.3763/ehaz.2010.0037
 47. Grafton RQ, Pittock J, Davis R, et al. Global insights into water resources, climate change and governance. *Nat Clim Chang* 2013,3:315-321. doi:10.1038/nclimate1746
 48. Piure A. Markets, Water Shares and Drought: Lessons from Australia. What Can the Water Industry in England and Wales Learn from Australia's Water Reform Story? 2014. <http://www.anglianwater.co.uk/about-us/statutory-reports/markets-water-shares-and-drought-lessons-from-australia.aspx>. Accessed February 22, 2016
 49. Sepulcre-Canto G, Horion S, Singleton A, Carrao H, Vogt J. Development of a Combined Drought Indicator to detect agricultural drought in Europe. *Nat Hazards Earth Syst Sci* 2012,12(11):3519-3531. doi:10.5194/nhess-12-3519-2012
 50. Pulwarty RS, Sivakumar MVK. Information systems in a changing climate: Early warnings and drought risk management. *Weather Clim Extrem* 2014,3:14-21. doi:10.1016/j.wace.2014.03.005
 51. Lackstrom K, Brennan A, Ferguson D, et al. The Missing Piece: Drought Impacts Monitoring. Workshop Report Produced by the Carolinas Integrated Sciences & Assessments Program and the Climate Assessment for the Southwest. 2013.
 52. Ding Y, Hayes MJ, Widhalm M. Measuring economic impacts of drought: a review and discussion. *Disaster Prev Manag An Int J* 2011,20(4):343-446. doi:10.1108/09653561111161752
 53. Environment Agency. Drought Response: Our Framework for England. 2015. https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/440728/National_Drought_Framework.pdf
 54. Dollar E, Edwards FK, Stratford C, et al. Monitoring and Assessment of Environmental Impacts of Droughts: Literature Synthesis. August 2013. <https://www.gov.uk/government/publications/monitoring-and-assessment-of-environmental-impacts-of-droughts-literature-synthesis>. Accessed October 14, 2015
 55. Smith KH, Svoboda M, Hayes M, Reges H, Doesken N, Lackstrom K, Dow K, Brennan A. Local Observers Fill In the Details on Drought Impact Reporter Maps. *Bull Am Meteorol Soc* 2014,95(11):1659-1662. doi:10.1175/1520-0477-95.11.1659
 56. Stahl K, Blauhut V, Kohn I, et al. A European Drought Impact Report Inventory (EDII): Design and Test for Selected Recent Droughts in Europe, DROUGHT-R& SPI Technical Report No. 3. 2012. <http://www.eu-drought.org/technicalreports/3>
 57. Wei J, Wei Y, Western A, Skinner D, Lyle C. Evolution of newspaper coverage of water issues in Australia during 1843-2011. *Ambio* October 2014. doi:10.1007/s13280-014-0571-2
 58. Bachmair S, Kohn I, Stahl K. Exploring the link between drought indicators and impacts. *Nat Hazards Earth Syst Sci* 2015,15(6):1381-1397. doi:10.5194/nhess-15-1381-2015

59. Dow K. News coverage of drought impacts and vulnerability in the US Carolinas, 1998–2007. *Nat Hazards* 2010,54(2):497-518.
60. Meadow AM, Crimmins MA, Ferguson DB. Field of Dreams, or Dream Team?: Assessing Two Models For Drought Impact Reporting in the Semiarid Southwest. *Bull Am Meteorol Soc* 2013,94:1507-1517. doi:10.1175/BAMS-D-11-00168.1
61. Potop V. Evolution of drought severity and its impact on corn in the Republic of Moldova. *Theor Appl Climatol* 2011,105(3-4):469-483. doi:10.1007/s00704-011-0403-2
62. Rhee J, Im J, Carbone GJ. Monitoring agricultural drought for arid and humid regions using multi-sensor remote sensing data. *Remote Sens Environ* 2010,114(12):2875-2887.
63. Haslinger K, Koffler D, Schöner W, Laaha G. Exploring the link between meteorological drought and streamflow: Effects of climate-catchment interaction. *Water Resour Res* 2014,50(3):2468-2487. doi:10.1002/2013WR015051
64. Kumar R, Musuuza JL, Van Loon AF, Teuling AJ, Barthel R, Ten Broek J, Mai J, Samaniego L, Attinger S. Multiscale evaluation of the standardized precipitation index as a groundwater drought indicator. *Hydrol Earth Syst Sci Discuss* 2015,12(8):7405-7436. doi:10.5194/hessd-12-7405-2015
65. Dai A. Characteristics and trends in various forms of the Palmer Drought Severity Index during 1900–2008. *J Geophys Res* 2011,116(D12):D12115. doi:10.1029/2010JD015541
66. Vicente-Serrano SM, Camarero JJ, Azorin-Molina C. Diverse responses of forest growth to drought time-scales in the Northern Hemisphere. *Glob Ecol Biogeogr* 2014,23(9):1019-1030. doi:10.1111/geb.12183
67. Vicente-Serrano SM, Beguería S, Lorenzo-Lacruz J, Camarero JJ, López-Moreno JI, Azorin-Molina C, Revuelto J, Morán-Tejeda E, Sanchez-Lorenzo A. Performance of drought indices for ecological, agricultural, and hydrological applications. *Earth Interact* 2012,16(10):1-27.
68. Wells N, Goddard S, Hayes MJ. A Self-Calibrating Palmer Drought Severity Index. *J Clim* 2004,17(12):2335-2351. doi:http://dx.doi.org/10.1175/1520-0442(2004)017<2335:ASPDSI>2.0.CO;2
69. McEvoy DJ, Huntington JL, Abatzoglou JT, Edwards LM. An Evaluation of Multiscalar Drought Indices in Nevada and Eastern California. *Earth Interact* 2012,16(18):1-18. doi:10.1175/2012EI000447.1
70. Stagge JH, Kohn I, Tallaksen LM, Stahl K. Modeling drought impact occurrence based on meteorological drought indices in Europe. *J Hydrol* 2015,530:37-50. doi:10.1016/j.jhydrol.2015.09.039
71. Lorenzo-Lacruz J, Vicente-Serrano SM, González-Hidalgo JC, López-Moreno JI, Cortesi N. Hydrological drought response to meteorological drought in the Iberian Peninsula. *Clim Res* 2013,58:117-131. doi:10.3354/cr01177
72. Vicente-Serrano SM, López-Moreno JI. Hydrological response to different time scales of climatological drought: an evaluation of the Standardized Precipitation Index in a mountainous Mediterranean basin. *Hydrol Earth Syst Sci* 2005,9(5):523-533.
73. Rahiz M, New M. Does a rainfall-based drought index simulate hydrological droughts? *Int J Climatol* 2014,34:2853-2871. doi:10.1002/joc.3879
74. Blauhut V, Gudmundsson L, Stahl K. Towards pan-European drought risk maps: quantifying the link between drought indices and reported drought impacts. *Environ Res Lett* 2015,10(1):014008. doi:10.1088/1748-9326/10/1/014008

75. Quiring SM, Ganesh S. Evaluating the utility of the Vegetation Condition Index (VCI) for monitoring meteorological drought in Texas. *Agric For Meteorol* 2010,150(3):330-339.
76. Potopová V, Štěpánek P, Možný M, Türkott L, Soukup J. Performance of the standardised precipitation evapotranspiration index at various lags for agricultural drought risk assessment in the Czech Republic. *Agric For Meteorol* 2015,202:26-38. doi:10.1016/j.agrformet.2014.11.022
77. Steinemann A. Drought Information for Improving Preparedness in the Western States. *Bull Am Meteorol Soc* 2014,95(6):843-847. doi:10.1175/bams-d-13-00067.1
78. Steinemann AC, Cavalcanti LFN. Developing multiple indicators and triggers for drought plans. *J Water Resour Plan Manag* 2006,132(3):164-174. doi:10.1061/(ASCE)0733-9496(2006)132:3(164)
79. Kruse S, Seidl I, Staehli M. Informationsbedarf zur Früherkennung von Trockenheit in der Schweiz. *Wasser Energ Luft* 2010,102(4):4.
80. Stölzle M, Stahl K. Wassernutzung und Trockenheitsindikatoren in Baden-Württemberg. *Standort-Zeitschrift für Angew Geogr* 2011,35(3):94-101.
81. Steinemann A, Iacobellis SF, Cayan DR. Developing and Evaluating Drought Indicators for Decision-Making. *J Hydrometeorol* 2015,16(4):1793-1803. doi:10.1175/JHM-D-14-0234.1
82. Lloyd-Hughes B, Saunders MA. A drought climatology for Europe. *Int J Climatol* 2002,22(13):1571-1592. doi:10.1002/joc.846
83. Spinoni J, Naumann G, Vogt J, Barbosa P. European drought climatologies and trends based on a multi-indicator approach. *Glob Planet Change* 2015,127:50-57. doi:10.1016/j.gloplacha.2015.01.012
84. Touma D, Ashfaq M, Nayak M, Kao S-C, Diffenbaugh NS. A Multi-model and Multi-index Evaluation of Drought Characteristics in the 21st Century. *J Hydrol* 2015,526:196-207. doi:10.1016/j.jhydrol.2014.12.011
85. Hao C, Zhang J, Yao F. Combination of multi-sensor remote sensing data for drought monitoring over Southwest China. *Int J Appl Earth Obs Geoinf* 2015,35:270-283. doi:10.1016/j.jag.2014.09.011
86. Anderson WB, Zaitchik BF, Hain CR, Anderson MC, Yilmaz MT, Mecikalski J, Schultz L. Towards an integrated soil moisture drought monitor for East Africa. *Hydrol Earth Syst Sci* 2012,16(8):2893-2913. doi:10.5194/hess-16-2893-2012
87. Dutra E, Wetterhall F, Di Giuseppe F, Naumann G, Barbosa P, Vogt J, Pozzi W, Pappenberger F. Global meteorological drought – Part 1: Probabilistic monitoring. *Hydrol Earth Syst Sci* 2014,18(7):2657-2667. doi:10.5194/hess-18-2657-2014
88. Hannah DM, Demuth S, van Lanen HAJ, Looser U, Prudhomme C, Rees G, Stahl K, Tallaksen LM. Large-scale river flow archives: importance, current status and future needs. *Hydrol Process* 2011,25(7):1191-1200. doi:10.1002/hyp.7794
89. Viglione A, Borga M, Balabanis P, Blöschl G. Barriers to the exchange of hydrometeorological data in Europe: Results from a survey and implications for data policy. *J Hydrol* 2010,394(1-2):63-77. doi:10.1016/j.jhydrol.2010.03.023
90. McDonald A. Rural Ethiopians use mobile phones to ring the alarm on drought. Oxfam Horn, East and Central Africa Blog. <http://www.oxfamblogs.org/eastafrica/?p=6017>. Published 2013. Accessed October 11, 2015

91. Demie L, Bekele A, Majwa P, Raj AJ. An Innovative Community Based Early Warning System That Uses Picture-Based Indicators and the Web Is the Way Forward for Ethiopia-Somaliland. 2011. http://www.fao.org/fileadmin/user_upload/drought/docs/REGLAP_magazine.pdf
92. UNISDR. Drought Risk Reduction, Framework and Practices, Contributing to the Implementation of the Hyogo Framework for Action. 2009.
93. Blackmore C, Ison R. Designing and developing learning systems for managing systemic change in a climate change world. In Wals, Arjen E. J. and Corcoran PB, ed. *Learning for Sustainability in Times of Accelerating Change: Wageningen Academic Publishers Books*. Wageningen: Wageningen Academic Publishers; 2012:347-363. doi:<http://dx.doi.org/10.3920/978-90-8686-757-8>
94. Everard M. The importance of periodic droughts for maintaining diversity in the freshwater environment. In *Freshwater Forum*, 7, 1, 33-50; 1996.
95. Oliver TH, Marshall HH, Morecroft MD, Brereton T, Prudhomme C, Huntingford C. Interacting effects of climate change and habitat fragmentation on drought-sensitive butterflies. *Nat Clim Chang* 2015,5(10):941-945. doi:10.1038/nclimate2746
96. Banerjee O, Bark R, Connor J, Crossman ND. An ecosystem services approach to estimating economic losses associated with drought. *Ecol Econ* 2013,91:19-27. doi:10.1016/j.ecolecon.2013.03.022
97. Pedro-Monzonís M, Solera A, Ferrer J, Estrela T, Paredes-Arquiola J. A Review of Water Scarcity and Drought Indexes in Water Resources Planning and Management. *J Hydrol* 2015,527:482-493. doi:10.1016/j.jhydrol.2015.05.003
98. AghaKouchak A, Farahmand A, Melton FS, Teixeira J, Anderson MC, Wardlow BD, Hain CR. Remote sensing of drought: progress, challenges and opportunities. *Rev Geophys* 2015,53(2):452-480. doi:10.1002/2014RG000456
99. Zhang Y, Peng C, Li W, Fang X, Zhang T, Zhu Q, Chen H, Zhao P. Monitoring and estimating drought-induced impacts on forest structure, growth, function, and ecosystem services using remote-sensing data: recent progress and future challenges. *Environ Rev* 2013,21(2):103-115. doi:10.1139/er-2013-0006
100. Quiring. Monitoring Drought: An Evaluation of Meteorological Drought Indices. *Geogr Compass* 2009,3(1):64-88. doi:10.1111/j.1749-8198.2008.00207.x
101. White DH, Walcott JJ. The role of seasonal indices in monitoring and assessing agricultural and other droughts: a review. *Crop Pasture Sci* 2009,60(7):599. doi:10.1071/CP08378
102. Heim Jr RR. A review of twentieth-century drought indices used in the United States. *Bull Am Meteorol Soc* 2002,83(8):1149-1165.
103. Keyantash J, Dracup JA. The Quantification of Drought: An Evaluation of Drought Indices. *Bull Am Meteorol Soc* 2002,83:1167-1180.
104. Economic Research Service United States Department of Agriculture. U.S. Drought 2012: Farm and Food Impacts. <http://www.ers.usda.gov/topics/in-the-news/us-drought-2012-farm-and-food-impacts.aspx>. Accessed October 10, 2015
105. Mosley LM. Drought impacts on the water quality of freshwater systems; review and integration. *Earth-Science Rev* 2015,140:203-214. doi:10.1016/j.earscirev.2014.11.010
106. Matthews WJ, Marsh-Matthews E. Effects of drought on fish across axes of space, time and ecological complexity. *Freshw Biol* 2003,48(7):1232-1253. doi:10.1046/j.1365-2427.2003.01087.x

107. Stanke C, Kerac M, Prudhomme C, Medlock J, Murray V. Health Effects of Drought: a Systematic Review of the Evidence. *PLoS Curr Disasters* 2013. doi:10.1371/currents.dis.7a2cee9e980f91ad7697b570bcc4b004.
108. Centers for Disease Control and Prevention, U.S. Environmental Protection Agency NO and AA and AWWA. When Every Drop Counts: Protecting Public Health during Drought Conditions - a Guide for Public Health Professionals. 2010.
109. Anderson MC, Hain C, Otkin J, Zhan X, Mo K, Svoboda M, Wardlow B, Pimstein A. An Intercomparison of Drought Indicators Based on Thermal Remote Sensing and NLDAS-2 Simulations with U.S. Drought Monitor Classifications. *J Hydrometeorol* 2013,14(4):1035-1056. doi:10.1175/JHM-D-12-0140.1
110. Carrão H, Singleton A, Naumann G, Barbosa P, Vogt J V. An Optimized System for the Classification of Meteorological Drought Intensity with Applications in Drought Frequency Analysis. *J Appl Meteorol Climatol* 2014,53(8):1943-1960. doi:10.1175/JAMC-D-13-0167.1
111. Ceglar A, Kajfež-Bogataj L. Simulation of maize yield in current and changed climatic conditions: Addressing modelling uncertainties and the importance of bias correction in climate model simulations. *Eur J Agron* 2012,37(1):83-95. doi:10.1016/j.eja.2011.11.005
112. Chandrasekar K, Sessa Sai MVR. Monitoring of late-season agricultural drought in cotton-growing districts of Andhra Pradesh state, India, using vegetation, water and soil moisture indices. *Nat Hazards* 2014,75(2):1023-1046. doi:10.1007/s11069-014-1364-4
113. Choi M, Jacobs JM, Anderson MC, Bosch DD. Evaluation of drought indices via remotely sensed data with hydrological variables. *J Hydrol* 2013,476:265-273. doi:10.1016/j.jhydrol.2012.10.042
114. Dieker E, Lanen HAJ van, Svoboda M. Comparison of Three Drought Monitoring Tools in the USA. WATCH Technical Report No. 25. WATCH 2010.
115. Edossa DC, Babel MS, Das Gupta A. Drought Analysis in the Awash River Basin, Ethiopia. *Water Resour Manag* 2009,24(7):1441-1460. doi:10.1007/s11269-009-9508-0
116. Gao Y, Markkanen T, Thum T, Aurela M, Lohila A, Mammarella I, Hagemann S, Aalto T. Assessing various drought indicators in representing drought in boreal forests in Finland. *Hydrol Earth Syst Sci Discuss* 2015,12(8):8091-8129. doi:10.5194/hessd-12-8091-2015
117. Gouveia C, Trigo RM, DaCamara CC. Drought and vegetation stress monitoring in Portugal using satellite data. *Nat Hazards Earth Syst Sci* 2009,9(1):185-195. doi:10.5194/nhess-9-185-2009
118. Gu Y, Hunt E, Wardlow B, Basara JB, Brown JF, Verdin JP. Evaluation of MODIS NDVI and NDWI for vegetation drought monitoring using Oklahoma Mesonet soil moisture data. *Geophys Res Lett* 2008,35(22).
119. Guttman NB. COMPARING THE PALMER DROUGHT INDEX AND THE STANDARDIZED PRECIPITATION INDEX. *J Am Water Resour Assoc* 1998,34(1):113-121. doi:10.1111/j.1752-1688.1998.tb05964.x
120. Jain VK, Pandey RP, Jain MK, Byun H-R. Comparison of drought indices for appraisal of drought characteristics in the Ken River Basin. *Weather Clim Extrem* 2015,8:1-11. doi:10.1016/j.wace.2015.05.002
121. Ji L, Peters AJ. Assessing vegetation response to drought in the northern Great Plains using vegetation and drought indices. *Remote Sens Environ* 2003,87(1):85-98.
122. Kumar MN. On the use of Standardized Precipitation Index (SPI) for drought intensity

- assessment. *Meteorol Appl* 2009,16:381-389. doi:10.1002/met.136
123. Li B, Rodell M. Evaluation of a model-based groundwater drought indicator in the conterminous U.S. *J Hydrol* 2015,526:78-88. doi:10.1016/j.jhydrol.2014.09.027
 124. Morid S, Smakhtin V, Moghaddasi M. Comparison of seven meteorological indices for drought monitoring in Iran. *Int J Climatol* 2006,26(7):971-985. doi:10.1002/joc.1264
 125. Naumann G, Dutra E, Barbosa P, Pappenberger F, Wetterhall F, Vogt J V. Comparison of drought indicators derived from multiple data sets over Africa. *Hydrol Earth Syst Sci* 2014,18(5):1625-1640. doi:10.5194/hess-18-1625-2014
 126. Potop V, Možný M, Soukup J. Drought evolution at various time scales in the lowland regions and their impact on vegetable crops in the Czech Republic. *Agric For Meteorol* 2012,156:121-133. doi:10.1016/j.agrformet.2012.01.002
 127. Scaini A, Sánchez N, Vicente-Serrano SM, Martínez-Fernández J. SMOS-derived soil moisture anomalies and drought indices: a comparative analysis using in situ measurements. *Hydrol Process* 2015,29(3):373-383. doi:10.1002/hyp.10150
 128. Shukla S, Steinemann AC, Lettenmaier DP. Drought Monitoring for Washington State: Indicators and Applications. *J Hydrometeorol* 2011,12(1):66-83. doi:10.1175/2010JHM1307.1
 129. Tadesse T, Wardlow B. Assessing the Vegetation Condition Impacts of the 2011 Drought across the US Southern Great Plains Using the Vegetation Drought Response Index (VegDRI). *J Appl Meteorol Climatol* 2015,54:153-169. doi:10.1175/JAMC-D-14-0048.1
 130. Vasiliades L, Loukas A. Hydrological response to meteorological drought using the Palmer drought indices in Thessaly, Greece. *Desalination* 2009,237(1):3-21.
 131. Vicente-Serrano SM, Gouveia C, Camarero JJ, et al. Response of vegetation to drought time-scales across global land biomes. *Proc Natl Acad Sci* 2013,110(1):52-57.
 132. Xia Y, Ek MB, Mocko D, Peters-Lidard CD, Sheffield J, Dong J, Wood EF. Uncertainties, Correlations, and Optimal Blends of Drought Indices from the NLDAS Multiple Land Surface Model Ensemble. *J Hydrometeorol* 2014,15(4):1636-1650. doi:10.1175/JHM-D-13-058.1
 133. Anderson MC, Hain C, Wardlow B, Pimstein A, Mecikalski JR, Kustas WP. Evaluation of Drought Indices Based on Thermal Remote Sensing of Evapotranspiration over the Continental United States. *J Clim* 2011,24(8):2025-2044. doi:10.1175/2010JCLI3812.1
 134. Bloomfield JP, Marchant BP. Analysis of groundwater drought building on the standardised precipitation index approach. *Hydrol Earth Syst Sci* 2013,17(12):4769-4787. doi:10.5194/hess-17-4769-2013
 135. Gu Y, Brown JF, Verdin JP, Wardlow B. A five-year analysis of MODIS NDVI and NDWI for grassland drought assessment over the central Great Plains of the United States. *Geophys Res Lett* 2007,34(6):L06407. doi:10.1029/2006GL029127
 136. Ma M, Ren L, Singh VP, Tu X, Jiang S, Liu Y. Evaluation and application of the SPDI-JDI for droughts in Texas, USA. *J Hydrol* 2015,521:34-45. doi:10.1016/j.jhydrol.2014.11.074
 137. Martínez-Fernández J, González-Zamora A, Sánchez N, Gumuzzio A. A soil water based index as a suitable agricultural drought indicator. *J Hydrol* 2015,522:265-273. doi:10.1016/j.jhydrol.2014.12.051
 138. Mendicino G, Senatore A, Versace P. A Groundwater Resource Index (GRI) for drought monitoring and forecasting in a mediterranean climate. *J Hydrol* 2008,357(3-4):282-302. doi:10.1016/j.jhydrol.2008.05.005

139. Rajsekhar D, Singh VP, Mishra AK. Multivariate drought index: An information theory based approach for integrated drought assessment. *J Hydrol* 2015,526:164-182. doi:10.1016/j.jhydrol.2014.11.031
140. Trnka M, Dubrovský M, Svoboda M, Semerádová D, Hayes M, Žalud Z, Wilhite D. Developing a regional drought climatology for the Czech Republic. *Int J Climatol* 2009,29(6):863-883. doi:10.1002/joc.1745
141. Vicente-Serrano SM, Beguería S, López-Moreno JI, Angulo M, El Kenawy A. A New Global 0.5° Gridded Dataset (1901–2006) of a Multiscalar Drought Index: Comparison with Current Drought Index Datasets Based on the Palmer Drought Severity Index. *J Hydrometeorol* 2010,11(4):1033-1043. doi:10.1175/2010JHM1224.1
142. Ziese M, Schneider U, Meyer-Christoffer A, Schamm K, Vido J, Finger P, Bissolli P, Pietzsch S, Becker A. The GPCP Drought Index – a new, combined and gridded global drought index. *Earth Syst Sci Data* 2014,6(2):285-295. doi:10.5194/essd-6-285-2014
143. Diodato N, Bellocchi G. Drought stress patterns in Italy using agro-climatic indicators. *Clim Res* 2008,36(1):53. doi:10.3354/cr00726
144. Hlavinka P, Trnka M, Semerádová D, Dubrovský M, Žalud Z, Možný M. Effect of drought on yield variability of key crops in Czech Republic. *Agric For Meteorol* 2009,149(3-4):431-442. doi:10.1016/j.agrformet.2008.09.004
145. Kattelus M, Salmivaara A, Mellin I, Varis O, Kummu M. An evaluation of the Standardized Precipitation Index for assessing inter-annual rice yield variability in the Ganges-Brahmaputra-Meghna region. *Int J Climatol* 2015. doi:10.1002/joc.4489
146. Mavromatis T. Drought index evaluation for assessing future wheat production in Greece. *Int J Climatol* 2007,27(7):911-924. doi:10.1002/joc.1444
147. Quiring SM, Papakryiakou TN. An evaluation of agricultural drought indices for the Canadian prairies. *Agric For Meteorol* 2003,118(1):49-62.
148. Rossi S, Niemeier S. Monitoring droughts and impacts on the agricultural production: Examples from Spain. In López-Francos A. (comp.) L-FA (collab.), ed. *Economics of Drought and Drought Preparedness in a Climate Change Context*. Options Mé. Zaragoza: CIHEAM / FAO / ICARDA / GDAR / CEIGRAM / MARM; 2010:35-40. <http://om.ciheam.org/article.php?IDPDF=801322>
149. Sun L, Mitchell S, Davidson A. Multiple drought indices for agricultural drought risk assessment on the Canadian prairies. *Int J Climatol* 2012. doi:10.1002/joc.2385
150. Bachmair S, Svensson C, Hannaford J, Barker LJ, Stahl K. A quantitative analysis to objectively appraise drought indicators and model drought impacts. *Hydrol Earth Syst Sci Discuss* 2015,12(9):9437-9488. doi:10.5194/hessd-12-9437-2015
151. Vicente-Serrano SM, Camarero JJ, Azorin-Molina C. Diverse responses of forest growth to drought time-scales in the Northern Hemisphere. *Glob Ecol Biogeogr* 2014,23(9):1019-1030. doi:10.1111/geb.12183

]

Figure captions

[Figure 1: Survey replies on the use of individual indicators in the drought early warning systems for different types of drought and impact data collection.

Figure 2: Survey replies on the importance of reasons for selecting the currently used individual indicators. Participants were asked to rate all reasons that apply. The dots represent the mean importance score, the bars the range; n= number of replies per category.

Figure 3: Survey replies on the reasons for initiating the drought early warning system (several ticks possible). The category 'Request by the government' represents 'Request by the government/governmental agencies/local authorities' in the original survey.

Figure 4: Survey replies on evaluating the selected drought indicators with drought impacts. Participants could tick both qualitative and quantitative evaluation, and several categories regarding the reasons for no evaluation.]

Tables

[Table 1: Reviews on drought indicators. Drought indicator type: V=various, M=meteorological, A=agricultural, H=hydrological (including groundwater), C=composite, O=other (e.g., remote sensing indicators)]

| Authors | Year | Scope | | Drought indicator type | | | | | | Topic |
|-------------------------------------|------|---------------------|--------------------------------|------------------------|---|---|---|---|---|---|
| | | Focus on indicators | Indicators among other aspects | V | M | A | H | C | O | |
| Hao and Singh ¹⁴ | 2015 | x | | | | | | | x | Composite indicators: methodologies, strengths and limitations |
| Pedro-Monzonis et al. ⁹⁷ | 2015 | x | | x | | | | | x | Drought and water scarcity indicators in the context of water resources planning |
| AghaKouchak et al. ⁹⁸ | 2015 | x | | | | | | | x | Drought monitoring using remote sensing observations |
| Van Loon ⁴² | 2015 | | x | x | | | x | | | Several aspects of drought including different indicators |
| Zhang et al. ⁹⁹ | 2013 | x | | | | | | | x | Remote sensing indicators in the context of impacts on forest ecosystems |
| Belal et al. ¹¹ | 2012 | | x | x | | | | | x | Several indicators, focus on remote sensing |
| Sivakumar et al. ⁸ | 2011 | x | | | | x | | | | WMO/UNISDR Expert Group Meeting: article collection on agricultural drought indices (usage, strengths and limitations) |
| Zargar et al. ¹⁸ | 2011 | x | | x | | | | | | Comprehensive review of 74 drought indicators |
| Dai ¹⁹ | 2011 | | x | | | | | | | Several aspects of drought including different indicators |
| Mishra and Singh ¹³ | 2010 | | x | x | | | | | | Several indicators: description, strengths and limitations |
| Quiring ¹⁰⁰ | 2009 | x | | | x | | | | | Meteorological drought indicators: description, strengths and limitations |
| White and Walcott ¹⁰¹ | 2009 | x | | x | | | | | | Various indicators, focus on relation to impacts on rural Australia |
| Heim ¹⁰² | 2002 | x | | x | | | | | | Major drought indicators used in the US |
| Keyantash and Dracup ¹⁰³ | 2002 | x | | x | | | | | | Several indicators: description, strengths and limitations |

Table 2: Reviews on drought impacts

| Authors | Year | Drought impact type | | | Topic |
|---|------|---------------------|---------------------------|------------|---|
| | | Various | Economic/ agricultural | Ecological | |
| Lackstrom et al. ⁵¹ | 2013 | x | | | Impact monitoring |
| Smith et al. ⁵⁵ | 2015 | x | | | Impact monitoring |
| Stahl et al. ¹⁶ | 2015 | x | | | Impact monitoring |
| Logar and van den Bergh ⁶ | 2011 | x | | | Cost assessment of drought damage |
| Ding et al. ⁵² | 2013 | | x | | Full scope of economic impacts |
| Economic Research Service, US Department of Agriculture ¹⁰⁴ | | | x | | Case study: impacts on US agriculture 2012 |
| Mosley ¹⁰⁵ | 2015 | | x | x | Impacts on water quality |
| Dollar et al. ⁵⁴ | 2013 | | | x | Ecological impacts focusing on rivers, lakes, wetlands and ponds |
| Matthews and Marsh- Matthews ¹⁰⁶ | 2003 | | | x | Impacts on fish |
| Stanke et al. ¹⁰⁷ | 2015 | | | | x Impacts on public health |
| Centers for Disease Control and Prevention, U.S. Environmental Protection Agency ¹⁰⁸ | 2010 | | | | x Impacts on public health |

Table 3: Original research papers assessing the link between different indicators, or between indicators and impacts; n=number of studies

| Linkage type | n | Categorization | n | Studies |
|---------------------|----|---|----|--|
| Indicator-indicator | 55 | Between existing indicators | 35 | Anderson et al. ¹⁰⁹ ; Carrao et al. ¹¹⁰ ; Ceglar et al.¹¹¹ ; Chandrasekar and Sesha Sai ¹¹² ; Choi et al.¹¹³ ; Dai ⁶⁵ ; Dieker et al.¹¹⁴ ; Edossa et al. ¹¹⁵ ; Gao et al. ¹¹⁶ ; Gouveia et al.¹¹⁷ ; Gu et al.¹¹⁸ ; Guttman ¹¹⁹ ; Haslinger et al. ⁶³ ; Jain et al. ¹²⁰ ; Ji and Peters¹²¹ ; Keyantash and Dracup ¹⁰³ ; Kumar ¹²² ; Kumar et al. ⁶⁴ ; Li and Rodell ¹²³ ; Lorenzo-Lacruz et al. ⁷¹ ; McEvoy et al. ⁶⁹ ; Morid et al. ¹²⁴ ; Naumann et al. ¹²⁵ ; Potop et al.¹²⁶ ; Quiring ¹⁰⁰ ; Quiring and Ganesh⁷⁵ ; Rahiz and New ⁷³ ; Scaini et al. ¹²⁷ ; Shukla et al. ¹²⁸ ; Tadesse and Wardlow¹²⁹ ; Vasiliades and Loukas ¹³⁰ ; Vicente-Serrano and López-Moreno ⁷² ; Vicente-Serrano et al.⁶⁷ ; Vicente-Serrano et al.¹³¹ ; Xia et al. ¹³² |
| | | Between a proposed novel indicator and existing ones | 20 | Anderson et al. ¹³³ ; Bloomfield and Marchant ¹³⁴ ; Gu et al.¹³⁵ ; Hao et al. ⁸⁵ ; Hao and AghaKoucha ³⁹ ; Ma et al. ¹³⁶ ; Ma et al. ³³ ; Martínez-Fernández et al.¹³⁷ ; Mendicino et al. ¹³⁸ ; Narasimhan and Srinivasan⁴⁰ ; Potop⁶¹ ; Rajsekhar et al. ¹³⁹ ; Rhee et al.⁶² ; Staudinger et al. ³² ; Trnka et al. ¹⁴⁰ ; Vicente-Serrano et al. ²⁵ ; Vicente-Serrano et al. ¹⁴¹ ; Wells et al. ⁶⁸ ; Zhang and Jia ⁴¹ ; Ziese et al. ¹⁴² |
| Indicator-impact | 31 | Crop yield | 14 | Ceglar et al.¹¹¹ ; Diodato and Bellocchi ^{143*} ; Hlavinka et al. ¹⁴⁴ ; Kattelus et al. ¹⁴⁵ ; Martínez-Fernández et al. ^{137*} ; Mavromatis ¹⁴⁶ ; Narasimhan and Srinivasan ^{40*} ; Potop^{61*} ; Potop et al.¹²⁶ ; Potopova et al. ⁷⁶ ; Quiring and Papakryiakou ¹⁴⁷ ; Rhee et al. ⁶² ; Rossi and Niemeyer ¹⁴⁸ ; Sun et al. ^{149*} |
| | | Remotely sensed vegetation stress | 7 | Choi et al.¹¹³ ; Gouveia et al.¹¹⁷ ; Gu et al.^{135*} ; Gu et al.¹¹⁸ ; Ji and Peters¹²¹ ; Quiring and Ganesh⁷⁵ ; Vicente-Serrano et al. ¹³¹ |
| | | Text-based data | 6 | Bachmair et al. ¹⁵⁰ ; Bachmair et al. ⁵⁸ ; Blauhut et al. ⁷⁴ ; Dieker et al.¹¹⁴ ; Stagge et al. ⁷⁰ ; Tadesse and Wardlow¹²⁹ |
| | | Other impact variables or several in parallel | 4 | Núñez et al. ³⁵ ; Sepulcre-Canto et al. ^{49*} ; Vicente-Serrano et al. ⁶⁷ ; Vicente-Serrano et al. ¹⁵¹ |

Colored font: study additionally assesses indicator-impact linkage;

*Impact variable used for evaluating the proposed novel indicator

]

Further Reading/Resources

[Table 1 and 2]

Related Articles

| DOI | Article title |
|-------------------|--|
| 10.1002/wat2.1085 | Hydrological drought explained |
| 10.1002/wcc.81 | Drought under global warming: a review |
| | |