

Open Research Online

The Open University's repository of research publications and other research outputs

Microstructural shock features in Lunar Mg-suite accessory phases

Conference or Workshop Item

How to cite:

Cernok, Ana; White, Lee Francis; Darling, James; Dunlop, Joseph; Johnson, Diane and Anand, Mahesh (2017). Microstructural shock features in Lunar Mg-suite accessory phases. In: Shock metamorphism in terrestrial and extra-terrestrial rocks (Miljković, Katarina ed.), 26 Jun - 2 Jul 2017, Curtin University, Perth, Western Australia.

For guidance on citations see [FAQs](#).

© [not recorded]

Version: Accepted Manuscript

Link(s) to article on publisher's website:

<http://www.sserviaustralia.org/event/shock-metamorphism-in-terrestrial-and-extra-terrestrial-rocks/>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

Microstructural shock features in Lunar Mg-suite accessory phases


ANA ČERNOK^{1*}, LEE FRANCIS WHITE², JAMES DARLING², JOSEPH DUNLOP², DIANE JOHNSON¹, AND MAHESH ANAND¹

¹Open University, School of Planetary Sciences, Walton Hall, Milton Keynes, MK7 6AA, United Kingdom

*Corresponding author: ana.cernok@open.ac.uk

²University of Portsmouth, School of Earth & Environmental Sciences, Burnaby Road, Portsmouth, PO1 3QL, United Kingdom

The lunar Mg-suite comprise plutonic highland rocks that represent an episode of crustal evolution following the primordial differentiation of the Moon [e.g. 1] and cover crystallization ages from 4.43 - 4.1 Ga [2]. We have examined thin sections of a shocked norite sampled by Apollo 17 mission and targeted apatite, merrillilite, and baddeleyite in an effort to constrain microstructural variation induced by shock metamorphism. Studied samples were assigned a high index of petrological pristinity but have experienced different levels of shock metamorphism [3]. Therefore, these materials present a unique opportunity to critically understand how impact processes have effected the microstructural evolution of these key accessory phases, constraining their effect on distribution of volatiles (Cl, F, OH) and radiogenic ages (U-Th-Pb) [e.g. 4-6]. Applied techniques include scanning electron microscopy (SEM), cathodoluminescence (CL), electron backscatter diffraction (EBSD), and Raman Spectroscopy. The preliminary dataset includes high-resolution (< 300 nm step size) EBSD maps of nine spatially associated apatite, five merrillite, and four baddeleyite grains, surrounded by different minerals. Apatite is found to form planar and non-planar fractures, with domains of sub-micrometer granular features that appear amorphous at the length scales of EBSD (~80 x 40nm). Texture component maps show up to 25° of misorientation within a single grain, evidence of severe crystal-plastic deformation. While these grains are heavily deformed, we conclude that no recrystallization of apatite has taken place. Merrillite is predominately granular, yielding randomly orientated subgrains indicative of solid-state or melt-recrystallization, with coarser granular textures clearly being evidence of a relatively late stage formation. A small number of merrillite grains contain extensive (< 30 °) crystal lattice deformation. Baddeleyite has degraded crystallinity and is often found to be amorphous at EBSD scale. Based on EBSD and CL analysis, its microtexture can be related to that of group 2 to 3 reported for Martian baddeleyites [6], which are believed to have experienced substantial loss of radiogenic Pb.. CL features show lighter cloudy regions in the areas of lower crystallinity as indexed by EBSD, indicating a high density of defects [e.g. 7]. No high pressure polymorph of any phase was detected by Raman spectroscopy. Raman spectra of all samples complement the EBSD pattern by indicating that well-crystallized sub-micrometer sized crystallites are surrounded by amorphous material. Merrillite shows lower shock impedance than apatite, but both higher than that of baddeleyite. These structural complexities cannot be visualised using backscatter electron (BSE) imaging alone. The evidence of microstructural changes in response to shock metamorphism of widely used accessory minerals should be taken into account when interpreting the volatile content and crystallization ages of lunar highland samples.



References: [1] Papike J. J. et al. (1996) *GCA*, 60, 3967-3978. [2] Shearer C.K. et al. (2015) *AM*, 100, 294-325. [3] Warren P.H. (1993) *AM*, 78, 360-376. [4] McCubbin F.M. et al. (2015) *AM*, 100, 1668-1707. [5] Adcock et al. (2017) *Nature Comm.*, 14667. [6] Darling J. R. et al. (2016), *EPSL*, 444, 1-12. [7] Niihara et al. (2012), *EPSL* 341-344, 195-210.