

# Open Research Online

---

The Open University's repository of research publications and other research outputs

## Fate of pathogens in a simulated bioreduction system for livestock carcasses

### Journal Item

How to cite:

Gwyther, Ceri L.; Jones, David L.; Golyshin, Peter N.; Edwards-Jones, Gareth and Williams, A. Prysor (2012). Fate of pathogens in a simulated bioreduction system for livestock carcasses. *Waste Management*, 32(5) pp. 933–938.

For guidance on citations see [FAQs](#).

© 2011 Elsevier

Version: Accepted Manuscript

Link(s) to article on publisher's website:

<http://dx.doi.org/doi:10.1016/j.wasman.2011.10.031>

---

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

---

[oro.open.ac.uk](http://oro.open.ac.uk)

1 **Fate of pathogens in a simulated bioreduction system for livestock carcasses**

2

3 Ceri L. Gwyther<sup>a</sup>, David L. Jones<sup>a</sup>, Peter N. Golyshin<sup>b</sup>, Gareth Edwards-Jones<sup>a</sup>, A. Prysor

4 Williams<sup>a</sup>\*

5 <sup>a</sup> *School of Environment, Natural Resources & Geography, College of Natural Sciences,*

6 *Bangor University, Gwynedd, LL57 2UW, UK*

7 <sup>b</sup> *School of Biological Sciences, College of Natural Sciences, Bangor University,*

8 *Gwynedd, LL57 2UW, UK*

9

10 \*Corresponding author: Tel.: +44 1248 382637; Fax: +44 1248 354997; E-mail:

11 prysor.williams@bangor.ac.uk

12

13

1 **ABSTRACT**

2 The EU Animal By-Products Regulations generated the need for novel methods of  
3 storage and disposal of dead livestock. Bioreduction prior to rendering or incineration has  
4 been proposed as a practical and potentially cost-effective method; however, its  
5 biosecurity characteristics need to be elucidated. To address this, *Salmonella enterica*  
6 (serovars Senftenberg and Poona), *Enterococcus faecalis*, *Campylobacter jejuni*,  
7 *Campylobacter coli* and a *lux*-marked strain of *Escherichia coli* O157 were inoculated  
8 into laboratory-scale bioreduction vessels containing sheep carcass constituents. Numbers  
9 of all pathogens and the metabolic activity of *E. coli* O157 decreased significantly within  
10 the liquor waste over time, and only *E. faecalis* remained detectable after three months.  
11 Only very low numbers of *Salmonella* spp. and *E. faecalis* were detected in bioaerosols,  
12 and only at initial stages of the trial. These results further indicate that bioreduction  
13 represents a suitable method of storing and reducing the volume of livestock carcasses  
14 prior to ultimate disposal.

15

16 **Keywords:** agriculture; animal slaughterhouse waste; fallen livestock; meat waste;  
17 zoonoses

18

1 **1. Introduction**

2 In order to reduce the risk of further outbreaks of animal diseases such as bovine  
3 spongiform encephalopathy and foot and mouth disease, the European Union introduced  
4 the Animal By-Products Regulations (EC/1774/2002) in 2003 (Anon, 2009). These  
5 regulations sought to improve biosecurity across all aspects of the livestock sector, from  
6 production to waste disposal. Since their implementation, the options available to most  
7 farmers to dispose of fallen (dead) livestock have been effectively limited to either  
8 rendering or incineration, whereas previously most fallen stock was buried. The  
9 regulations have led to animosity within the agricultural industry due to the considerable  
10 costs and biosecurity concerns associated with centralised collection and rendering or  
11 incineration of fallen stock (Bansback, 2006; Gwyther et al., 2011). Indeed, there is call  
12 for both a change in legislation and the development of alternative methods of disposal  
13 (Bansback, 2006).

14 Bioreduction is a novel technology that has shown potential as a viable option for  
15 storing and pre-treating fallen stock prior to disposal (Williams et al., 2009). Bioreduction  
16 is the aerobic biodegradation of animal by-products in a partially sealed vessel, where the  
17 contents are mildly heated and aerated and ultimately disposed of via the permitted route  
18 for ‘Category 1’ material in accordance to the EU ABPR (i.e. via incineration or  
19 rendering). The process has been shown to reduce the volume of waste and hence the  
20 frequency of collection and associated disposal cost, as well as being a practical method  
21 for industry (Williams et al., 2009).

22 Dead livestock may harbour a range of zoonotic agents (Milnes et al., 2008), and  
23 current methodologies for their disposal in Europe (e.g. incineration and rendering)

1 depend on high temperatures to deactivate pathogens; however, bioreduction operates at a  
2 mesophilic temperature (approx. 40 °C) and does not utilise any chemical disinfection  
3 procedure. Rather, the active aeration coupled with the competitive and antagonistic  
4 effects of the prevalent microbes are hypothesised to reduce pathogen levels (Williams et  
5 al., 2009). For bioreduction to be approved under the revised EU ABPR (EC/1069/2009)  
6 (as described in Annex IV of EU implementing Regulation EC 142/2011) as an  
7 alternative method of storing fallen stock prior to disposal, the fate of pathogens within  
8 the system must be elucidated and the evidence presented to the European Food Safety  
9 Authority (EFSA), which then decide whether to ratify the system for industry use  
10 (Bohm, 2008). EFSA stipulate that novel disposal methods should lead to a 5-log  
11 reduction in the numbers of two indicator organisms representing bacterial pathogens,  
12 *Salmonella enterica* serovar Senftenberg (hereafter called *S. Senftenberg*) and  
13 *Enterococcus faecalis* (Bohm, 2008). A previous field-scale study on bioreduction of  
14 sheep recovered negligible numbers of pathogens (Williams et al., 2009), but the initial  
15 pathogen concentration was not high enough to validate whether or not a 5-log reduction  
16 in numbers had occurred. Whilst it is preferable to assess the fate of pathogens at field-  
17 scale, the logistics of growing and handling the large volumes of pathogens needed to  
18 gain a sufficient concentration in the bioreduction vessels would be problematic. Further,  
19 EFSA guidelines state that simulated systems can be used as a proxy of field-scale  
20 systems provided that they are representative of actual conditions (EFSA, 2008).

21 The aim of this work was to validate the effectiveness of bioreduction in reducing  
22 numbers of introduced pathogens in a laboratory-scale system. By applying the criteria  
23 stipulated by EFSA for ratifying novel *disposal* methods to a simulated *storage* process

1 that is bioreduction, this study will help verify whether bioreduction represents a  
2 biosecure method of containing fallen stock prior to disposal. In addition to *S.*  
3 Senftenberg and *Enterococcus faecalis*, additional microorganisms (*Campylobacter* spp.,  
4 *E. coli* O157, and other *Salmonella* strains) were also tested as they represent common  
5 zoonotic pathogens that may be introduced with carcasses into bioreduction vessels.

6

## 7 **2. Materials and methods**

### 8 *2.1 Vessel design*

9 Laboratory-scale versions of the bioreduction vessels described by Williams et al.  
10 (2009) were constructed using 5 l polypropylene containers; 19 cm high × 13 cm wide ×  
11 26 cm long. These mini bioreducer vessels (MBVs) were placed within a darkened  
12 incubator set to 40 °C (± 2 °C) and the contents continuously aerated at a maximum rate  
13 of 6 l min<sup>-1</sup>. To negate microbial contamination and odour, the outflow from the MBVs  
14 were passed through a commercial disinfectant (20% Trigene; Medichem, Kent, UK) and  
15 then an odour trap containing activated charcoal before being vented into a Category II  
16 biosafety cabinet (Fig. 1).

17

### 18 *2.2 Trial management*

19 The inoculated MBVs ( $n = 3$ ) were managed in a similar way to the field-scale  
20 bioreduction vessels (Williams et al., 2009). Specifically, an initial volume of water (2.2  
21 l; equivalent to 2,800 l under field-scale conditions) was added so that each MBV was  
22 just under half filled. A total of  $231 \pm 1.5$  g of sheep carcass components (equivalent to

1 300 kg under field-scale conditions) were added to each MBV. Components (sourced  
2 locally) comprised of muscle, bone, fat (in an intact ‘chop’), pelt, blood, stomach  
3 contents, wool and liver, in proportions representative of a sheep carcass (MLC, 2006), to  
4 give a final volume of approximately 2.5 l, determined by volume marks on each MBV  
5 and by weight. Particle size was uniform across all MBVs, up to a maximum of  
6 approximately 100 mm (bone), with components being distributed uniformly within each  
7 MBV. A commercial catalyst with sodium alginate as the active ingredient (Gel-60<sup>®</sup>,  
8 Biopolym, Spain) was added at the recommended dose of 1 g catalyst to 1 kg of carcass  
9 (Williams et al., 2009). The final ratio of liquid to solids was 10:1. Each MBV was  
10 aerated as described previously. Water levels were visually inspected every other day and  
11 fresh water added so that two-thirds of the carcass contents (by height) were covered  
12 throughout the trial (Gutiérrez et al., 2003; Lobera et al., 2007). When individual carcass  
13 components were no longer recognisable, water was added to prevent drying out of the  
14 remaining organic matter. Control MBVs ( $n = 2$ ) were managed under the same  
15 conditions but without the addition of pathogens.

16

## 17 2.3 Inoculation

### 18 2.3.1 Bacterial strains

19 *S. Senftenberg* (NCTC13385) was obtained from HPA culture collections (Health  
20 Protection Agency, Salisbury, UK) and *Enterococcus faecalis* (ATCC 29212) was  
21 obtained from Oxoid (Oxoid Ltd., Hampshire, UK). *Salmonella enterica* (ser) Poona  
22 (hereafter called *S. Poona*) (NCTC4840) was obtained from Oxoid; a *lux*-marked strain of  
23 *E. coli* O157 (3704 Tn5 *luxCDABE*) and an environmental strain of *E. coli* O157 (#3704)

1 (both non-pathogenic but accurately reflecting survival of pathogenic strains) were used  
2 to represent verocytotoxic *E. coli* (Kudva et al., 1998; Ritchie et al., 2003); and  
3 *Campylobacter jejuni* (6035) and *Campylobacter coli* (6168) were kindly donated by the  
4 University of Aberdeen (Ogden et al., 2009). All media were bought from Oxoid unless  
5 otherwise stated.

6

### 7 2.3.2 Microbiological preparation and inoculation

8 All bacterial strains were grown from frozen stock, with each (except  
9 *Campylobacter*) grown overnight in tryptone soya broth (TSB; CM0129) then  
10 subsequently combined with other strains of the same organism and incubated overnight  
11 in fresh TSB in an orbital shaker (150 rev min<sup>-1</sup>) at 37 °C. *C. coli* and *C. jejuni* were  
12 grown overnight in Bolton Broth (CM0983) containing lysed horse blood (SR0048) and a  
13 supplement containing cefoperazone, vancomycin, trimethoprim and cycloheximide  
14 (SR0183) in a microaerobic environment at 41.5 °C and subsequently combined and  
15 again incubated overnight. The microaerobic environments were obtained using  
16 anaerobic jars and CampyGen sachets (CN0025).

17 The final concentration of each inoculum was obtained by serially diluting and  
18 plating onto selective agar: *Salmonella* on Xylose-Lysine-Desoxycholate Agar (XLD,  
19 CM0469); *E. coli* O157 on Sorbitol MacConkey Agar (CM0813) containing cefixime and  
20 potassium tellurite (CT-SMAC; SR0172); *E. faecalis* on Slanetz and Bartley Medium  
21 (SBM; CM0377) and *Campylobacter* on modified Charcoal Cefoperazone Deoxycholate  
22 agar (mCCDA; CM0739) containing the supplement cefoperazone and amphotericin B  
23 (SR0155). SBM plates were incubated for 48 hours at 37 °C whereas mCCDA plates



1 were incubated microaerobically for 48 hours at 41.5 °C. All other plates were incubated  
2 at 37 °C for 24 hours. Each treatment MBV was subsequently inoculated so that the  
3 concentration of micro-organisms per ml of liquid was as follows: 7.91 log<sub>10</sub> CFU of  
4 *Salmonella*, 7.89 log<sub>10</sub> CFU of *E. faecalis*, 7.5 log<sub>10</sub> CFU of *E. coli* O157 and 6.81 log<sub>10</sub>  
5 CFU of *Campylobacter*.

6

## 7 2.4 *Liquor waste*

### 8 2.4.1 *Microbiological analysis*

#### 9 2.4.1.1 *Enumeration*

10 Liquor samples (25 ml) were recovered directly from each MBV using a 5 ml  
11 pipette on days 0, 3, 23, 56 and 84 and analysed on the same day. Samples were  
12 homogenised in a Seward 400 stomacher (Seward Ltd., Worthing, UK) for 1 min at 230  
13 rev min<sup>-1</sup> with 225 ml maximum recovery diluent (MRD, CM0733) then serially diluted  
14 in MRD. Samples were enumerated as described previously for each pathogen. Total  
15 viable counts (TVC) were enumerated using the pour plate techniques based on and BS  
16 EN ISO 4833:2003 (FSA, 2009a) using standard plate count agar (PCA, CM0463). All  
17 presumptive colonies were sub-cultured onto nutrient agar and incubated at 37 °C for 24  
18 hours, whilst presumptive *Campylobacter* colonies were incubated microaerobically at  
19 41.5 °C for 48 hours. *E. coli* O157, *Campylobacter* and *Salmonella* spp. were confirmed  
20 using latex agglutination (DR0620, DR0150 and FT0203, respectively) with further  
21 biochemical tests using Microbact™ GNB 12A (MB1132) and sub-culturing onto  
22 *Salmonella* chromogenic agar (CM1007 containing supplement SR0194) for *Salmonella*

1 spp.. Confirmation of *E. faecalis* was performed using glucose agar (Sigma, 16447) and  
2 subsequently sub-culturing onto bile aesculin agar (CM0888).

3

#### 4 2.4.1.2 Enrichment

5 Where *Salmonella*, *E. coli* O157 and *Campylobacter* were not detected by  
6 enumeration, enrichment was used to confirm the absence of these bacteria. Enrichment  
7 of samples for *Salmonella* spp. was based on ISO standards 6579:2002 (FSA, 2009b) and  
8 *Campylobacter* samples were enriched based on the ISO 10272-1:2006(E) method (HPA,  
9 2007). For the enrichment of *E. coli* O157, 20 ml of mTSB (CM0989 containing  
10 supplement SR0190) was added to 5 ml of liquor and shaken in an orbital shaker for 6  
11 hours (37 °C, 150 rev min<sup>-1</sup>) after which time 0.1 ml of the enriched culture was streaked  
12 onto duplicate plates of CT-SMAC. Plates were incubated and presumptive colonies  
13 confirmed as described previously.

14

#### 15 2.4.2 Activity of *E. coli* O157

16 At each sampling time-point, a 1 ml aliquot from each MB was placed into a plastic  
17 luminometer cuvette and its luminescence [relative light units (RLU)] determined using a  
18 SystemSURE 18172 luminometer (Hygiena Int., Watford, UK).

19

#### 20 2.5 Bioaerosol analysis

21 Bioaerosol samples were taken on days 0, 24, 57, and 85. Samples were obtained  
22 from gases before they passed through the commercial disinfectant using selective agar

1 plates in an Andersen Air Sampler 2000 (Andersen Instruments Inc; Atlanta, Georgia,  
2 USA). The pump was connected to the mini bioreducer by silicone tubing and activated  
3 for 30 min at a flow rate of 10 l min<sup>-1</sup>. Plates were arranged randomly with each sampling  
4 date but were consistent between replicates on the same sampling date. *E. faecalis* were  
5 captured using Slanetz and Bartley Medium, *Salmonella* using Brilliant Green Agar  
6 (BGA, CM0263), *Campylobacter* using mCCDA, *E. coli* O157 using CT-SMAC, and  
7 TVC using PCA. All plates were incubated as described previously whilst the BGA was  
8 incubated at 37 °C for 24 h.

9

## 10 *2.6 Data Analysis*

11 Samples where micro-organisms were detected only via enrichment were allocated  
12 an arbitrary value of half the detection limit. To avoid analysing data with zero values  
13 (i.e. those not detected by enrichment), all data were log<sub>10</sub> (y + 1) transformed. Data were  
14 analysed using SPSS v15.0 (SPSS Inc., Chicago, USA). Normality of the microbiological  
15 data was tested using the Kolmogorov-Smirnov test and means analysed using either  
16 related samples *t*-test if normal or Wilcoxon signed rank test if non-normal.

17

## 18 **3. Results**

### 19 *3.1 Waste degradation*

20 At the end of the trial, the reduction in volume of carcass components in each  
21 vessel was similar (88.2 ± 3.7% of that initially added). The discernable animal remains

1 were predominantly identified as stomach content although there were also some fatty  
2 deposits and small fragments of bone.

3

### 4 3.2 Microbiological characteristics

5 The controls were found to have natural populations of *Salmonella* spp., *E. faecalis*,  
6 and *Campylobacter* spp. but no *E. coli* O157 were detected. Survival of the introduced  
7 *Salmonella* spp. and *E. faecalis* in the treatment MBVs followed similar survival patterns  
8 to natural populations in the controls (Fig. 2). Although numbers of both *Salmonella* spp.  
9 and *E. faecalis* reduced markedly over the three month trial, the dynamics of survival  
10 differed between both micro-organisms. Specifically, *Salmonella* spp. numbers remained  
11 relatively stable until day 54, after which they significantly declined ( $P < 0.05$ ) so that  
12 they could only be detected by enrichment at the end of the trial period (Fig. 2A).  
13 Numbers of *E. faecalis* generally decreased more steadily throughout the trial, although  
14 had recovered somewhat in the control MBs towards the latter stages (Fig. 2B) ( $P > 0.05$ ).

15 A significant ( $P < 0.05$ ) decline in both the numbers and activity of *E. coli* O157  
16 were seen in the inoculated MBVs (Fig. 3A), culminating in a 5-log reduction by day 84  
17 and luminescence values falling to below background levels. *Campylobacter* spp.  
18 numbers declined significantly ( $P < 0.05$ ) within the first three days followed by an  
19 increase in numbers in both the control and inoculated MBVs on day 23 and a subsequent  
20 decrease (Fig. 3B). On day 56, *Campylobacter* spp. numbers in the inoculated vessels had  
21 reduced by greater than 5-log values and remained so until the end of the trial. TVC  
22 values dropped initially in the inoculated MBVs, but then recovered and stabilised  
23 towards latter stages of the trial (Fig. 4). In the control MBVs, there was a non-significant

1 increase in numbers up to day 23 ( $P > 0.05$ ), followed by a gradual decrease to values  
2 very similar to those in the treatment bioreducers.

3

### 4 3.3 Bioaerosols

5 No pathogens were recovered as bioaerosols from the control bioreducers. Low  
6 numbers of *Salmonella* spp. and *E. faecalis* were detected as bioaerosols in initial stages  
7 of the trial from the inoculated MBVs; although no *Salmonella* were detected after the  
8 first sampling date and numbers of *E. faecalis* decreased considerably with each sampling  
9 date until they were undetectable (Table 1). Although this mimicked the decline in mean  
10 concentration of *E. faecalis* within the liquor, the relationship between bioaerosol and  
11 liquor counts was not statistically significant ( $P > 0.05$ ; data not shown). Neither *E. coli*  
12 O157 nor *Campylobacter* were detected within any bioaerosol samples. TVC values  
13 increased towards the middle and latter stages of the trial in both the inoculated and  
14 control systems and were statistically similar throughout ( $P > 0.05$ ); although numbers  
15 recovered were more variable with time.

16

## 17 4. Discussion

18 This trial was conducted over three months as it has been shown that this is the time  
19 required for most of the carcass components to degrade within a bioreduction system.  
20 Bioreduction has already proved to be effective at reducing the volume of carcass  
21 material to be disposed (Williams et al., 2009) and the findings of this trial supported this  
22 as even the bone material largely degraded. Although the system was designed to

1 accurately mimic field-scale bioreduction, it should be remembered that the surface area  
2 of the contents added to the MBVs was proportionately greater than that of an entire  
3 carcass and that the aeration rate was greater; caution is therefore needed when  
4 extrapolating the rates of degradation to field-scale.

5 In keeping with previous work on bioreduction (Gutiérrez et al., 2003, Williams et  
6 al., 2009), a catalyst product, Gel-60<sup>®</sup> (Biopolym, Spain), was used in this trial. Gel-60<sup>®</sup>  
7 is based on sodium alginate and it is claimed that the hydrolysis of fibre, proteins, poly-  
8 and oligo-saccharides within it provides ready substrates for the indigenous microbial  
9 community, encouraging microbial growth and degradation of the animal carcass  
10 (Gutiérrez et al., 2003). However, the efficacy of Gel-60<sup>®</sup> has recently been trialled and  
11 was found to induce no affect on degradation rates (data not shown).

12 At the end of the bioreduction process, there remains a nutrient-rich liquid that  
13 should be disposed of via the permitted route for 'Category 1' material according to the  
14 EU ABPR (Williams et al., 2009). Lobera et al. (2007) recommend retaining a proportion  
15 of the liquor to facilitate the bioreduction of future carcasses, as is advised to inoculate  
16 new compost or anaerobic digestion plants (Gerardi, 2003; Sundberg and Johnson, 2005).  
17 In addition to reducing the volume for final disposal, the advantage of liquid waste  
18 compared to solid is that it can be removed easily from the vessel by use of a vacuum  
19 pump. Relevant biosecurity and environmental factors to consider with bioreduction and  
20 the liquor waste generated have been explored and compared to other methods of carcass  
21 storage and disposal by Gwyther et al. (2011).

22 EFSA stipulate that novel disposal methods for animal by-products should lead to a  
23 5-log reduction in the numbers of *S. Senftenberg* and *E. faecalis* as a measure of

1 biosecurity (Bohm, 2008). Bioreduction isn't however a disposal method for fallen stock,  
2 but rather a storage system designed to reduce the volume of waste in a safe, contained  
3 environment prior to ultimate disposal via incineration or rendering (Williams et al.,  
4 2009). Over the three month trial period, all microorganisms, with the exception of *E.*  
5 *faecalis*, had reduced by 5-log values, although *E. faecalis* had also notably decreased by  
6 over 4-log values. However, due to the low numbers of *Salmonella* spp. identified on the  
7 final sampling date, it is recommended that future trials go beyond the three month period  
8 in order to gain a better insight into the fate of the organism. A carcass storage system  
9 similar to bioreduction, though anaerobic, was trialled by Gutiérrez et al. (2003) and a  
10 number of commensal pathogens, including *Salmonella* spp., could not be detected after  
11 55 days. Although their results concur with the ones obtained in this current study, a  
12 direct comparison isn't possible due to the difference in aeration status of the systems  
13 trialled and because Gutiérrez et al. (2003) looked at presence or absence of pathogens,  
14 rather than numbers. EFSA also stipulate that novel disposal methods for animal by-  
15 products should lead to a 3-log reduction in the numbers of suitable indicator viruses  
16 (Bohm, 2008). Viruses were not analysed in this trial due to logistical issues, but the fate  
17 of porcine parvovirus will be tested in future field-scale trials of bioreduction.

18 *Campylobacter* spp. was predicted to be absent from all samples due to its penchant  
19 for microaerophilic conditions. Indeed, temporary technical issues resulted in a brief  
20 reduction in aeration rate on day 23 which corresponds with an observed increase in  
21 *Campylobacter* spp. numbers; thereafter numbers reduced considerably and none were  
22 recovered at the latter stages of the trial (Fig. 3B). TVC values were initially lower in

1 inoculated MBVs than the sum of both *Salmonella* and *E. faecalis*; though this is likely to  
2 reflect the differences in the nature of the agars and plating methods used.

3 Williams et al. (2009) hypothesised that microbial competition and predation  
4 reduced the population of pathogens within bioreduction vessels. Numerous other studies  
5 have showed the reduction of pathogens (including *S. Senftenberg*) in a range of wastes is  
6 promoted by competition from naturally present antagonistic microbes (Ceustermans et  
7 al., 2007; Hussong et al., 1985; Pietronave et al., 2004; Sidhu et al., 2001). However,  
8 some species of *Enterococcus* are able to produce bacteriocins that give them a  
9 competitive edge over other organisms (Fisher and Philips, 2009). Together with its  
10 ability to survive a wide range of environmental conditions (Fisher and Philips, 2009),  
11 this may explain why *E. faecalis* could still be recovered at the end of the trial; although  
12 it too may have decreased further had the trial period been extended.

13 Many pathogens can enter a viable but non-culturable state (VBNC) when under  
14 environmental stress and this may lead to underestimation of numbers when using  
15 culturing methods. However, bacteria containing the *lux* gene that have entered a VBNC  
16 state can still be detected in real-time by measuring bioluminescence (Duncan et al.,  
17 1994; Ritchie et al., 2003). Luminescence directly reports on bacterial metabolic activity  
18 which represents a prerequisite for host infection (Unge et al., 1999). Bioluminescence  
19 measurements showed that there was a concomitant decrease in both numbers and  
20 metabolic activity of *E. coli* O157; hence conditions within the vessels were not  
21 conducive to the organism's proliferation. Although currently not widely available,  
22 further trials with additional constructed *lux*-marked pathogens would be worthwhile to  
23 determine the effects of bioreduction on a greater range of target organisms.



1           Only low levels of bioaerosols were detected from the simulated bioreduction  
2 systems. Andersen samplers recover only the culturable fraction of micro-organisms and  
3 various molecular techniques have occasionally been shown to be useful in analysing  
4 bioaerosols in different environments (Fallschissel et al., 2009; Junhui et al., 1997; Maron  
5 et al., 2005). However, Andersen samplers are the preferred industry choice for sampling  
6 bioaerosols from other types of organic wastes (Stagg et al., 2010), and the large volume  
7 of air processed at each sampling point was deemed more than sufficient to detect the  
8 presence of bioaerosols. Our findings support others studies (Adkin et al., 2010) that  
9 propose bioreduction provides negligible risk of hazardous bioaerosol generation.

10           Where possible, more than one strain or serotype of each pathogen was added to the  
11 simulated bioreduction systems so as to negate potential inter-strain variation in survival.  
12 Whilst natural strains of pathogens may show greater resistance to environmental stresses  
13 than experimental cultures, it is unlikely that natural populations of the microaerophilic  
14 organisms or facultative anaerobes such as those used in this study would be any better  
15 adapted to cope with the process of forced aeration and other stresses encountered within  
16 a bioreduction system (Williams et al., 2009). This was also evident in our results as  
17 survival patterns of introduced strains were similar to natural strains within controls for  
18 all micro-organisms. However, molecular methods (e.g. automated ribosomal intergenic  
19 spacer analysis fingerprinting) may differentiate between strains (Cardinale et al., 2004)  
20 and therefore could be used to elucidate such points in future trials.

21           The rate of carcass degradation is likely to be largely governed by biological  
22 activity and the composition of the microbial community. Utilising molecular techniques  
23 may elucidate the changes in microbial population and whether such changes affect the

1 rate of degradation. Such information may be valuable in developing novel biological  
2 catalyst products to further improve the process. Further work at field-scale is also  
3 necessary to validate the energy demand of bioreduction and to identify where gains in  
4 efficiencies could be made.

5

## 6 **5. Conclusions**

7 This work indicates that bioreduction is efficient at containing pathogens from  
8 carcass material and hence that the system could potentially be suitably secure to store  
9 fallen stock prior to ultimate disposal. Further investigation at field-scale level that also  
10 includes other relevant organisms (e.g. indicator viruses) is required so that the system  
11 can be soundly considered for industry use and incorporation into the revised EU Animal  
12 By-Products Regulations (1069/2009).

13

## 14 **Acknowledgments**

15 We are grateful to BPEX, the Welsh Assembly Government and to Hybu Cig Cymru –  
16 Meat Promotion Wales for funding the work. We thank Sarah Chesworth for her technical  
17 assistance throughout the study.

18

19

1 **References**

- 2 Adkin A., Matthews D., Hope J., Maddison B.C., Somerville R.A., Pedersen J., 2010.  
3 Risk of escape of prions in gaseous emissions from on-farm digestion vessels.  
4 *Vet. Rec.* 167, 28-29.
- 5 Anon., 2009. The Animal By-Products Regulations, Vol. 1069/2009, European  
6 Commission, Brussels.
- 7 Bansback, B., 2006. Independent Review of the National Fallen Stock Scheme and  
8 Company. [http://archive.defra.gov.uk/foodfarm/byproducts/documents/nfsco-](http://archive.defra.gov.uk/foodfarm/byproducts/documents/nfsco-review.pdf)  
9 [review.pdf](http://archive.defra.gov.uk/foodfarm/byproducts/documents/nfsco-review.pdf) (accessed 12/08/11).
- 10 Bohm, R., 2008. The experimental validation and the organisms to be considered in the  
11 context of the ABP Regulation. European Food Safety Authority, Parma, Italy.
- 12 Cardinale, M., Brusetti, L., Quatrini, P., Borin, S., Puglia, A.M., Rizzi, A., Zanardini, E.,  
13 Sorlini, C., Corselli, C., Daffonchio, D., 2004. Comparison of different primer  
14 sets for use in automated ribosomal intergenic spacer analysis of complex  
15 bacterial communities. *Appl. Environ. Microbiol.* 70, 6147-6156.
- 16 Ceustermans, A., De Clerq D., Aertsen, A., Michiels, C., Coosemans, J., Ryckeboer, J.,  
17 2007. Inactivation of *Salmonella* Senftenberg strain W 775 during composting of  
18 biowastes and garden wastes. *J. Appl. Microbiol.* 103, 53-64.
- 19 Duncan, S., Glover, L.A., Killham, K., Prosser, J.I., 1994. Luminescence-based detection  
20 of activity of starved and viable but nonculturable bacteria. *Appl. Environ.*  
21 *Microbiol.* 60, 1308-1316.
- 22 EFSA, 2008. Guidelines for applications for new alternative methods of disposal or use  
23 of animal by-products under regulation (EC) no 1774/2002. The Health and

1 Consumer Protection Directorate-General and the European Food Safety  
2 Authority SANCO/2806/2008, Parma, Italy.

3 Fallschissel, K., Kampfer, P., Jackel, U., 2009. Direct detection of *Salmonella* cells in the  
4 air of livestock stables by real-time PCR. *Ann Occup Hyg* 53, 859-868.

5 Fisher, K., Phillips, C., 2009. The ecology, epidemiology and virulence of *Enterococcus*.  
6 *Microbiology* 155, 1749-1757.

7 FSA, 2009a. Total Aerobic Plate Counts. Food Standards Agency publications, London,  
8 UK.

9 FSA, 2009b. Detection of *Salmonella* for red meat samples (Based on the ISO  
10 6579:2002 Standard Protocol). Food Standards Agency publications, London,  
11 UK.

12 Gerardi, M.H., 2003. *The Microbiology of Anaerobic Digesters*. John Wiley & Sons Inc.

13 Gutiérrez, C., Ferrández, F., Andujar, M., Martín, J., Clemente, P., Lobera, J., 2003.  
14 Results of the preliminary study into: Physicochemical and bacteriological  
15 parameters of the hydrolisation of non-ruminant animal carcasses with bio-  
16 activators. Murcia University, Spain.

17 Gwyther, C.L., Williams, A.P., Golyshin, P.N., Edwards-Jones, G., Jones, D.L., 2011  
18 The environmental and biosecurity characteristics of livestock carcass disposal  
19 methods: A review. *Waste Manag.* 31, 767-778.

20 HPA, 2007. Identification of *Campylobacter* species. Health Protection Agency  
21 publications, London, UK.

22 Hussong, D., Burge, W.D., Enkiri, N.K., 1985. Occurrence, growth, and suppression of  
23 salmonellae in composted sewage-sludge. *Appl. Environ. Microbiol.* 50, 887-893.

1 Junhui, Z., Ruifu, Y., Fengxiang, C., Meiling, C., Hong, C., 1997. Polymerase chain  
2 reaction analysis of laboratory generated bioaerosols. *Aerobiologia*. 13, 7-10.

3 Kudva, I.T., Blanch, K., Hovde, C.J., 1998. Analysis of *Escherichia coli* O157:H7  
4 survival in ovine or bovine manure and manure slurry. *Appl. Environ. Microbiol.*  
5 64, 3166-3174.

6 Lobera, J.B., González, M., Sáez, J., Montes, A., Clemente, P., Quiles, A., Crespo, F.,  
7 Alonso, F., Carrizosa, J.A., Andújar, M., Martínez, D., Gutiérrez, C., 2007. Final  
8 report about the results on monogastric animal corpse hydrolyzation: Experience  
9 based on pigs production. Report submitted to the European Commission.

10 Maron, P.A., Lejon, D.P.H., Carvalho, E., Bizet, K., Lemanceau, P., Ranjard, L., Mougél  
11 C., 2005. Assessing genetic structure and diversity of airborne bacterial  
12 communities by DNA fingerprinting and 16S rDNA clone library. *Atmos Environ*  
13 39, 3687-3695.

14 Milnes, A.S., Stewart, I., Clifton-Hadley, F.A., Davies, R.H., Newell, D.G., Sayers, A.R.,  
15 Cheasty, T., Cassar, C., Ridley, A., Cook, A.J.C., Evans, S.J., Teale, C.J., Smith,  
16 R.P., McNally, A., Toszeghy, M., Fütter, R., Kay, A., Paiba, G.A., 2008.  
17 Intestinal carriage of verocytotoxigenic *Escherichia coli* O157, *Salmonella*,  
18 thermophilic *Campylobacter* and *Yersinia enterocolitica*, in cattle, sheep and pigs  
19 at slaughter in Great Britain during 2003. *Epidemiol. Infect.* 136, 739-751.

20 MLC, 2006. A review of the red meat offal & by-products industry. Meat and Livestock  
21 Commission publications, Warwickshire, UK.

22 Ogden, I.D., Dallas, J.F., MacRae, M., Rotariu, O., Reay, K.W., Leitch, M., Thomson,  
23 A.P., Sheppard, S.K., Maiden, M., Forbes, K.J., Strachan, N.J.C., 2009.

1           *Campylobacter* excreted into the environment by animal sources: prevalence,  
2           concentration shed, and host association. *Foodborne Pathog. Dis.* 6, 1161-1170.

3   Pietronave, S., Fracchia, L., Rinaldi, M., Martinotti, M.G., 2004. Influence of biotic and  
4           abiotic factors on human pathogens in a finished compost. *Water Res* 38, 1963-  
5           1970.

6   Ritchie, J.M., Campbell, G.R., Shepherd, J., Beaton, Y., Jones, D., Killham, K., Artz,  
7           R.R.E., 2003. A stable bioluminescent construct of *Escherichia coli* O157:H7 for  
8           hazard assessments of long-term survival in the environment. *Appl. Environ.*  
9           *Microbiol.* 69, 3359-3367.

10   Sidhu, J., Gibbs, R.A., Ho, G.E., Unkovich. I., 2001. The role of indigenous  
11           microorganisms in suppression of *Salmonella* regrowth in composted biosolids.  
12           *Water Res.*, 35, 913-920.

13   Stagg, S., Bowry, A., Kelsey, A., Crook, B., 2010. Bioaerosol emissions from waste  
14           composting and the potential for workers' exposure. Health and Safety Executive  
15           publications, London, UK.

16   Sundberg, C., Jonsson, H. 2005. Process inhibition due to organic acids in fed-batch  
17           composting of food waste - influence of starting culture. *Biodegradation*, 16, 205-  
18           213.


19   Unge, A., Tombolini, R., Mølbak, L., Jansson, J. K., 1999. Simultaneous monitoring of  
20           cell number and metabolic activity of specific bacterial populations with a dual *gfp*-  
21           *luxAB* marker system. *Appl. Environ. Microbiol.*, 65, 813-821.

- 1 Williams AP, Edwards-Jones G, Jones D.L, 2009. In-vessel bioreduction provides an
- 2 effective storage and pre-treatment method for livestock carcasses prior to final
- 3 disposal. *Bioresour. Technol.* 100, 4032-4040.


**Table 1:** Detection of bioaerosols from the inoculated mini bioreduction vessels (CFU m<sup>-3</sup>). Values represent means ± standard error of the mean. ND denotes not detected.

Day	<i>Salmonella</i> spp.	<i>E. faecalis</i>	<i>E. coli</i> O157	<i>Campylobacter</i> spp.	Total Viable Counts
0	31.9 ± 21.2	15.6 ± 10.9	ND	ND	15.7 ± 6.3
24	ND	4.4 ± 4.4	ND	ND	147.8 ± 126.7
57	ND	1.1 ± 1.1	ND	ND	2925.6 ± 2917.2
85	ND	ND	ND	ND	41.3 ± 31.3







**Figure 1:** Experimental layout of the mini bioreduction vessels (MBV). Each MBV was attached to a large disinfection Trigene bottle via silicone tubing with the exception of the second control MBV which was attached to its own Trigene bottle and vacuum pump.



**Figure 2:** Changes in numbers of *Salmonella* spp. (A) and *Enterococcus faecalis* (B) over time in the inoculated and control mini bioreduction vessels. The dashed line represents a 5-log reduction in numbers from the original starting concentration. Values represent means  $\pm$  standard error of the mean.



**Figure 3:** Changes in numbers and metabolic activity (as measured by relative light units of luminescence) of *Escherichia coli* O157 in the inoculated mini bioreduction vessels (A) and numbers of *Campylobacter* spp. (B) over time in the inoculated and control mini bioreduction vessels. The dashed line represents a 5-log reduction in numbers from the original starting concentration. No *E. coli* O157 was present in the controls. Values represent means  $\pm$  standard error of the mean.



**Figure 4:** Changes in numbers of Total Viable Counts on standard plate count agar over time in the inoculated and control mini bioreduction vessels. Values represent means  $\pm$  standard error of the mean.