

# Open Research Online

---

The Open University's repository of research publications and other research outputs

## Surface-based 3D measurements of small aeolian bedforms on Mars

Conference or Workshop Item

How to cite:

Balme, Matthew; Robson, Ellen; Barnes, Robert; Butcher, Frances; Fawdon, Peter; Huber, Ben; Ortner, Thomas; Paar, Gerhard; Traxler, Christoph; Bridges, John; Gupta, Sanjeev and Vago, Jorge (2018). Surface-based 3D measurements of small aeolian bedforms on Mars. In: EGU General Assembly 2018, 8-13 Apr 2018, Vienna, Austria.

For guidance on citations see [FAQs](#).

© 2018 The Authors

Version: Version of Record

Link(s) to article on publisher's website:

<https://meetingorganizer.copernicus.org/EGU2018/EGU2018-15847.pdf>

---

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

---

[oro.open.ac.uk](https://oro.open.ac.uk)



## Surface-based 3D measurements of small aeolian bedforms on Mars

Matthew Balme (1), Ellen Robson (2), Robert Barnes (3), Frances Butcher (1), Peter Fawdon (1), Ben Huber (4), Thomas Ortner (5), Gerhard Paar (4), Christoph Traxler (5), John Bridges (6), Sanjeev Gupta (3), and Jorge Vago (7)

(1) Open University, Dept. of Physical Sciences, Milton Keynes, United Kingdom (matt.balme@open.ac.uk), (2) School of Geography, Earth and Environmental Sciences, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK, (3) Department of Earth Sciences and Engineering, Imperial College London, London SW7 2AZ, UK, (4) Joanneum Research, Institute for Information and Communication Technologies, Steyrergasse 17, 8010 Graz, Austria, (5) vrVis Forschungs GmbH., Donau-City-Strasse 11, 1220 Vienna, Austria, (6) Space Research Centre, University of Leicester, Leicester LE1 7RH, UK, (7) European Space Agency, Keplerlaan 1, 2200 AG Noordwijk, The Netherlands

Recent aeolian bedforms such as dunes and ripples are common on Mars and occur across a scale range of centimetres to kilometres. Aeolian bedforms provide a mobility hazard to Mars rovers, such as the ESA ExoMars rover, which will launch in 2020 to either Mawrth Vallis or Oxia Planum. However, both sites contain numerous aeolian bedforms, mainly with simple ripple-like morphologies, and wavelengths of meters to tens of metres. The larger examples are ‘Transverse Aeolian Ridges’ (TARs), a few metres high and up to a few tens of metres across. TARs present a serious, but recognized and (hopefully) avoidable, rover mobility hazard that can be assessed using orbital data. Both sites also host smaller (1-4 m across) bedforms of similar ripple-like morphology, but it is unknown whether these bedforms will be traversable by the ExoMars rover: they are too small to be resolved in even the best Digital Terrain Models (DTMs) available.

Here, we estimate the heights of these small bedforms by comparison with similar features observed by previous Mars rovers. We use rover-based stereo imaging from the NASA Mars Exploration Rover (MER) Opportunity and PRo3D software, a 3D visualisation and analysis tool, to measure the size and height of ripple-like bedforms in the Meridiani Planum region of Mars. These are good analogues for the smaller bedforms at the ExoMars rover candidate landing sites.

We find that bedform height scales linearly with length (as measured across the bedform, perpendicular to the crest ridge) with a ratio of about 1:15. Using 25cm/pixel HiRISE images, we measured small aeolian bedforms at the Oxia Planum candidate landing site, finding them to be similar in length and morphology to those at Meridiani Planum. This allows us to infer the heights of bedforms at the ExoMars sites simply by measuring their lengths in HiRISE images. These data can then be used to explore the traversability of this site.

Our method suggests that most of the bedforms studied in Oxia Planum have ridge crests higher than 15 cm, but lower than 25 cm. Hence, if the tallest bedforms the ExoMars rover will be able to safely cross are only 15 cm high, then the Oxia Planum site studied here contains mostly impassable bedforms. However, if the rover can safely traverse 25 cm high bedforms, then most bedforms here will be smaller than this threshold. Finally, our results show that the mini-TARs have length/height ratios similar to TARs in general. Hence, these bedforms should probably be classified simply as ‘small TARs’, and are not a discrete population or sub-type of aeolian bedforms.