

# Open Research Online

---

The Open University's repository of research publications and other research outputs

## Potential Identification of sublimation-driven downslope mass movement on Mercury

Conference or Workshop Item

How to cite:

Malliband, C. C.; Conway, S. J.; Rothery, D. A. and Balme, M. R. (2018). Potential Identification of sublimation-driven downslope mass movement on Mercury. In: Mercury: Current and Future Science of the Innermost Planet, 1-3 May 2018, Columbia, Maryland, USA.

For guidance on citations see [FAQs](#).

© [\[not recorded\]](#)

Version: Version of Record

---

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's [data policy](#) on reuse of materials please consult the policies page.

---

[oro.open.ac.uk](http://oro.open.ac.uk)

**POTENTIAL IDENTIFICATION OF SUBLIMATION-DRIVEN DOWNSLOPE MASS MOVEMENT ON MERCURY.** C.C. Malliband<sup>1</sup>, S.J. Conway<sup>\*2</sup>, D.A. Rothery<sup>1</sup>, and M.R. Balme<sup>1</sup>, <sup>1</sup>School of Physical Sciences, The Open University, Milton Keynes, MK7 6AA, UK (chris.malliband@open.ac.uk), <sup>2</sup>CNRS, LPG, Université de Nantes, France (susan.conway@univ-nantes.fr) <sup>\*</sup>presenting author

**Introduction:** Mass movement has been recognised on many solar system bodies. Evidence of mass movement on Mercury has previously been limited to a single documented example, found in the pyroclastic vent NE of Rachmaninoff crater. Here we present the identification of a second example in a small, recent impact crater,  $\approx 285$  km north of Nabokov.

**Mass movement at NE Rachmaninoff Vent:** Attention was drawn to the slope features in the NE Rachmaninoff vent (Fig. 1) on the MESSENGER website, but so far as we are aware there has been no formal study. The features are downslope erosion-deposition systems with an alcove at the head, chute and a fan at the base. Feature heads appear to develop in a stratigraphic layer of brighter material (Fig. 1B). This brighter material appears to be related to hollows [1].

**Slope features in unnamed crater ( $\approx 285$  km N of Nabokov):** The newly identified slope features are in a  $\approx 12$  km diameter simple impact crater. The crater is

surrounded by low reflectance material and has an area of hollows on the NE crater rim. The slope features start just below the crater rim, in a bright stratigraphic layer. This may be similar to the hollow forming layer in NE Rachmaninoff.

**Current work:** We are examining other areas with steep slopes and catalogued hollows, and are performing a global survey to identify any further examples. We will compare the topography of these features to erosion-deposition systems on Mars [2,3] and Vesta [4]. Our working hypothesis is that these downslope movements are caused by sublimation.

**References:** [1] Blewett D.T. et al. (2011) *Science*, 333, 1856–1859. [2] Conway S.J. et al. (2015) *Icarus*, 253, 189–204. [3] Brusnikin E.S. et al. (2016) *Icarus*, 278, 52–61. [4] Scully J.E.C. et al. (2015) *EPSL*, 411, 151–163.

**Acknowledgements:** This work is partially supported by the CNES, in preparation for BepiColombo.




Fig. 1: A,B,C Gully-like slope features in vent NE of Rachmaninoff. A: Context image centered on  $36^{\circ}\text{N}$ ,  $63.8^{\circ}\text{E}$  showing widespread slope features. Box shows location of B. (NAC: EN1003843866M) B: High resolution (6.4m/pixel) image showing source at bright layer (NAC: EN1028933034M) C: Enhanced color. D, E, F. Newly discovered slope features at  $8^{\circ}\text{S}$ ,  $55^{\circ}\text{E}$ . D, enhanced color; E, high incidence angle (NAC: EN0252295266M) F, low incidence angle. Note bright, possibly hollow forming, material, high in crater wall (NAC: EN1028933034M).