

# Open Research Online

---

The Open University's repository of research publications and other research outputs

## Asymptotically approaching the Moore bound for diameter three by Cayley graphs

### Journal Item

How to cite:

Bachratý, Martin; Šiagiová, Jana and Širáň, Jozef (2019). Asymptotically approaching the Moore bound for diameter three by Cayley graphs. *Journal of Combinatorial Theory, Series B*, 134 pp. 203–217.

For guidance on citations see [FAQs](#).

© 2018 Elsevier Inc.



<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Version: Accepted Manuscript

Link(s) to article on publisher's website:  
<http://dx.doi.org/doi:10.1016/j.jctb.2018.06.003>

---

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

---

[oro.open.ac.uk](http://oro.open.ac.uk)

# Asymptotically approaching the Moore bound for diameter three by Cayley graphs

Martin Bachratý

*Comenius University, Bratislava, Slovakia*

Jana Šiagiová

*Slovak University of Technology, Bratislava, Slovakia*

Jozef Širáň

*Open University, U.K., and Slovak University of Technology, Slovakia*

---

## Abstract

The largest order  $n(d, k)$  of a graph of maximum degree  $d$  and diameter  $k$  cannot exceed the Moore bound, of the form  $M(d, k) = d^k - O(d^{k-1})$  for  $d \rightarrow \infty$  and any fixed  $k$ . Known results in finite geometries on generalised  $(k+1)$ -gons imply, for  $k = 2, 3, 5$ , existence of an infinite sequence of values of  $d$  such that  $n(d, k) = d^k - o(d^k)$ . Thus, for  $k = 2, 3, 5$  the Moore bound can be asymptotically approached in the sense that  $\limsup_{d \rightarrow \infty} n(d, k)/M(d, k) = 1$ ; moreover, no such result is known for any other value of  $k \geq 2$ . The corresponding graphs are, however, far from vertex-transitive, and there appears to be no obvious way to extend them to vertex-transitive graphs giving the same type of asymptotic result.

The second and the third author (2012) proved by a direct construction that the Moore bound for diameter  $k = 2$  can, in a similar sense, be asymptotically approached by *Cayley* graphs. Subsequently, the first and the third author (2015) showed that the same construction can be derived from generalised triangles with polarity.

By a detailed analysis of regular orbits of suitable groups of automorphisms of graphs arising from polarity quotients of incidence graphs of generalised quadrangles with polarity, we prove that for an infinite set of values of  $d$  there exist Cayley graphs of degree  $d$ , diameter 3, and order  $d^3 - O(d^{2.5})$ . The Moore bound for diameter 3 can thus as well be asymptotically approached by Cayley graphs. We also show that this method does not extend

to constructing Cayley graphs of diameter 5 from generalised hexagons with polarity.

*Keywords:* Degree, Diameter, Cayley graph, Generalized quadrangle, Suzuki-Tits oval.

*2000 MSC:* 05C12 (primary), 05B25, 05C25 (secondary).

---

## 1. Introduction

For positive integers  $d$  and  $k$  let  $n(d, k)$  be the largest order of a graph of maximum degree  $d$  and diameter  $k$ . It is well known that the value of  $n(d, k)$  cannot exceed the Moore bound  $M(d, k) = 1 + d + d(d-1) + \dots + d(d-1)^{k-1}$ . Setting trivial cases aside, for  $d \geq 3$  and  $k \geq 2$  we have  $n(d, k) = M(d, k)$  only for  $k = 2$  and  $d = 3, 7$ , and possibly 57, the unique graphs for the first two degrees being the Petersen and the Hoffman-Singleton graph [11, 2, 5]. For a survey of results about (im)possibility of getting ‘close’ to the Moore bound for the remaining values of  $d$  and  $k$  we refer to [18]. The main driving forces in this field appear to be the question of Bermond and Bollobás [3] if for arbitrarily large  $c$  there exist  $d, k$  such that  $n(d, k) < M(d, k) - c$ , and the problem of Delorme [6] of determining the value of  $\limsup_{d \rightarrow \infty} n(d, k)/d^k = \limsup_{d \rightarrow \infty} n(d, k)/M(d, k)$  for every fixed  $k \geq 2$ .

Regarding the question of Bermond and Bollobás, a substantial progress has recently been made in [9] by proving that for any fixed  $d$  and any  $c > 0$  the order of the largest *vertex-transitive*  $d$ -regular graph of diameter  $k$  is smaller than  $M(d, k) - c$  for almost all  $k$ . The best available result on Delorme’s problem is his own observation [6] that  $\limsup_{d \rightarrow \infty} n(d, k)/M(d, k) = 1$  for  $k \in \{2, 3, 5\}$ . This follows by taking polarity quotients of the incidence graphs of generalised  $n$ -gons admitting a polarity (cf. [17]) for  $n \in \{3, 4, 6\}$ , respectively. Despite having a fairly large automorphism group compared to their order, these graphs are not even regular and by [1] there appears to be no obvious way to extend them to vertex-transitive graphs by just adding edges in the case of diameter  $k = 2$ .

For the remaining diameters the best currently known results on Delorme’s problem are much weaker but still far from easy to prove. We know that  $\limsup_{d \rightarrow \infty} n(d, 4)/M(d, 4) \geq 1/4$  by [7], and from [4] we have  $\limsup_{d \rightarrow \infty} n(d, k)/M(d, k) \geq (1.6)^{-k}$  for  $k \geq 6$ , where 1.6 can be replaced by 1.57 for  $k \equiv -1, 0, 1 \pmod{6}$ .

In the light of the above-mentioned result of [9] addressing the question of Bermond and Bollobás for vertex-transitive graphs, it is natural to ask if the Moore bound can be *asymptotically approached*, in the sense of Delorme's limit superior being equal to 1, by *vertex-transitive*, or even *Cayley* graphs. The importance of this direction of research is underscored by the fact that, from the practical point of view, computer generation of record large graphs of given degree and diameter is almost exclusively limited to searching over Cayley graphs in cases when the degree or diameter exceed values manageable by other methods; cf. [14] and the online tables [25].

To this end, for  $d \geq 3$  and  $k \geq 2$  we let  $vt(d, k)$  and  $Cay(d, k)$  denote the largest order of a vertex-transitive and a Cayley graph, respectively, of degree  $d$  and diameter  $k$ ; clearly,  $vt(d, k) \geq Cay(d, k)$ . The task now is to estimate the values of  $\limsup_{d \rightarrow \infty} vt(d, k)/M(d, k)$  and  $\limsup_{d \rightarrow \infty} Cay(d, k)/M(d, k)$  for  $k \geq 2$  and, specifically, to determine if the Moore bound can be asymptotically approached by vertex-transitive or Cayley graphs for diameters 2, 3 and 5. Here, however, the available results are scarcer and, expectedly, not as good as those for  $n(d, k)$ . Let us begin with  $k \geq 3$ . In the vertex-transitive case, the digraphs of [10] yield, after ignoring edge directions,  $\lim_{d \rightarrow \infty} vt(d, k)/M(d, k) \geq 2^{-k}$  for every  $k \geq 3$ . For Cayley graphs, constructions of [15, 16] give  $\lim_{d \rightarrow \infty} Cay(d, k)/M(d, k) \geq k \cdot 3^{-k}$  for every  $k \geq 3$ , with improvements of the lower bounds by [24] to  $3 \cdot 2^{-4}$ ,  $32 \cdot 5^{-4}$  and  $25 \cdot 4^{-5}$  for  $k = 3, 4$  and  $5$ , respectively.

For  $k = 2$  the strongest finding so far in this category is the fact that  $\limsup_{d \rightarrow \infty} Cay(d, 2)/M(d, 2) = 1$ , showing that the Moore bound for diameter 2 can be asymptotically approached by Cayley graphs. This result was obtained in [22] by a direct construction of certain Cayley graphs of one-dimensional affine groups over finite fields of characteristic 2. Later in [1] it was shown that the construction of [22] is equivalent to extending a regular orbit of a polarity quotient of the incidence graph of a generalised triangle under the action of a suitable group.

Our aim is to show that the Moore bound for diameter 3 can also be asymptotically approached by Cayley graphs, that is, to prove the equality  $\limsup_{d \rightarrow \infty} Cay(d, 3)/M(d, 3) = 1$ . In fact, we prove that for an infinite set of values of  $d$  there exist Cayley graphs of degree  $d$ , diameter 3, and order  $d^3 - O(d^{2.5})$ . The method is a variant of the one used in [1], namely, extension of a regular orbit of a suitable subgroup of the automorphism group of a polarity quotient of the incidence graph of a generalised quadrangle. Details, however, are much more subtle and complex in comparison with those of [1].

We also show that an extension of this method is not feasible for proving an analogous result for diameter 5.

The paper is organised as follows. In Section 2 we review basic concepts on the finite generalised quadrangles with polarity and their automorphisms. These are used in Section 3 to study incidence in auxiliary graphs obtained from the incidence graphs of finite generalised quadrangles by polarity. In Section 4 we investigate induced subgraphs of the auxiliary graphs obtained as orbits of suitable groups of automorphisms. The induced subgraphs are finally extended in Section 5 to give Cayley graphs which give  $\limsup_{d \rightarrow \infty} \text{Cay}(d, 3)/M(d, 3) = 1$ . We conclude by showing that applying this scenario to generalised hexagons with polarity does not produce Cayley graphs asymptotically approaching the Moore bound for diameter 5.

## 2. Generalised quadrangles, polarity and symmetries

Let  $q$  be a prime power and let  $F = GF(q)$  be the Galois field of order  $q$ . As usual, let  $F^+$  and  $F^*$  be the additive and the multiplicative group of  $F$ . We begin by recalling the projective geometry  $PG(3, q)$  whose points are the 1-dimensional subspaces of  $F^4$  minus the origin (sometimes called projective vectors), that is, equivalence classes  $[\mathbf{x}]$  of non-zero quadruples  $\mathbf{x} = (x_0, x_1, x_2, x_3) \in F^4$ , with two quadruples  $\mathbf{x}$  and  $\mathbf{y}$  equivalent if  $y_i = tx_i$  for some  $t \in F^*$  and every  $i \in \{0, 1, 2, 3\}$ .

For further considerations we also need to involve a skew-symmetric bilinear form  $Q : F^4 \times F^4 \rightarrow F$ ; in classical texts such as [19] this form is usually taken to be  $(\mathbf{x}, \mathbf{y}) \mapsto x_0y_1 - x_1y_0 + x_2y_3 - x_3y_2$  but here we prefer to work with an equivalent form given by  $Q(\mathbf{x}, \mathbf{y}) = x_0y_3 - x_3y_0 + x_1y_2 - x_2y_1$ . Note that this choice of  $Q$  is obtained from the ‘classical’ one application of the permutation  $(0)(132)$ , that is, by cyclically permuting the last three coordinates of the input vectors as shown. A subset  $S$  of  $PG(3, q)$  is *totally isotropic* (with respect to  $Q$ ) if for any two points  $[\mathbf{x}], [\mathbf{y}] \in S$  we have  $Q(\mathbf{x}, \mathbf{y}) = 0$ . Note that all the  $q^3 + q^2 + q + 1$  points are totally isotropic themselves, and an easy counting argument shows that there are exactly  $q^3 + q^2 + q + 1$  totally isotropic lines (that is, totally isotropic 2-dimensional subspaces of  $F^4$  with the origin removed) of  $PG(3, q)$ .

Total isotropy helps us introduce an important incidence geometry within  $PG(3, q)$ , standardly denoted  $W(q)$ ; see e.g. [17, 20]. Points of  $W(q)$  are the points of  $PG(3, q)$ , lines of  $W(q)$  are the totally isotropic lines of  $PG(3, q)$ , and incidence is defined by containment as in  $PG(3, q)$ . Every line of  $W(q)$

contains  $q + 1$  points (as in  $PG(3, q)$ ) and, by counting, every point of  $W(q)$  lies on  $q + 1$  lines of  $W(q)$ . The incidence structures  $W(q)$  are prominent examples of *generalised quadrangles* [17, 20].

A polarity  $\pi$  of  $W(q)$  is an involutory mapping that sends the point set of  $W(q)$  onto its line set and vice versa, with the property that for any two points  $u, v$  lying on a line  $\ell$  of  $W(q)$ , the lines  $\pi(u)$  and  $\pi(v)$  intersect at the point  $\pi(\ell)$ . By a classical result of Tits [23], the incidence structure  $W(q)$  admits a polarity if and only if  $q$  is an odd power of 2. We describe such a polarity next, following [19] but using the skew-symmetric bilinear form  $Q$  introduced earlier.

From now on and throughout the paper, let  $q = 2^{2n+1}$  for some positive integer  $n$ . Let  $\omega = 2^{n+1}$  and let  $\sigma$  be the automorphism of  $F = GF(q)$  given by  $\sigma(x) = x^\omega$ , so that  $\sigma^2(x) = x^2$ . For every point  $u = [\mathbf{x}] \in W(q)$  let  $c = x_0x_3 + x_1x_2$  and for every pair of points  $[\mathbf{x}], [\mathbf{y}] \in W(q)$  we let  $\delta_{ij} = x_iy_j + x_jy_i$  for any distinct  $i, j \in \{0, 1, 2, 3\}$ ; dependence of  $c$  on  $[\mathbf{x}]$  and of  $\delta_{ij}$  on  $[\mathbf{x}], [\mathbf{y}]$  will always be assumed but not shown in the notation. With this in mind, a polarity on  $W(q)$  can be introduced as follows.

**Proposition 2.1.** *For a point  $u = [\mathbf{x}] \in W(q)$  let  $\pi(u)$  be the set of all non-zero vectors of  $F^4$  spanned by the totally isotropic set of four vectors*

$$(0, x_0^\omega, x_1^\omega, c^{\omega/2}), (x_0^\omega, 0, c^{\omega/2}, x_2^\omega), (x_1^\omega, c^{\omega/2}, 0, x_3^\omega) \text{ and } (c^{\omega/2}, x_2^\omega, x_3^\omega, 0). \quad (1)$$

*In the reverse direction, for a line  $\ell$  of  $W(q)$  through a pair of distinct points  $[\mathbf{x}]$  and  $[\mathbf{y}]$  let  $\pi(\ell)$  be the point  $[\mathbf{z}] \in W(q)$  with coordinates given by*

$$z_0 = \delta_{01}^{\omega/2}, z_1 = \delta_{02}^{\omega/2}, z_2 = \delta_{13}^{\omega/2} \text{ and } z_3 = \delta_{23}^{\omega/2}. \quad (2)$$

*Then  $\pi(u)$  is a line of  $W(q)$ , and the mapping  $\pi$  is a polarity on  $W(q)$ .*

**Proof.** The reasoning is the same as in the proof of Theorem 12.7.1 of [19, p. 566], up to subscripts. Namely, the choice of the bilinear form  $Q$  introduced in the first paragraph (made upon a suggestion of an anonymous referee) the formulae (1) and (2) presented here differ from the ones in [19, p. 565-566] as a result of a systematic application of the permutation (0)(132) to the subscripts of coordinates of points while keeping the definition of  $\delta_{ij}$  the same in [19] and here. We give here an independent sketch of the crucial part of the argument in our coordinate setting.

Let  $u = [\mathbf{x}]$  and  $v = [\mathbf{y}]$  be a pair of distinct points on a line  $\ell$  in  $W(q)$ . To prove the result it is sufficient to show that  $\pi^2 = id$  (which we leave out as it is routine) and that the lines  $\pi(u)$  and  $\pi(v)$  intersect at the point  $\pi(\ell)$ .

By (2),  $\pi(\ell)$  is the point  $[\mathbf{z}]$  with homogeneous coordinates  $z_0 = \delta_{01}^{\omega/2}$ ,  $z_1 = \delta_{02}^{\omega/2}$ ,  $z_2 = \delta_{13}^{\omega/2}$  and  $z_3 = \delta_{23}^{\omega/2}$ , where  $\delta_{ij} = x_i y_j + x_j y_i$ . The lines  $\pi(u)$  and  $\pi(v)$  are determined by a pair of linearly independent vectors from a set of four vectors given by (1) and we will illustrate the process just on  $\pi(u)$ . If, say,  $x_0 \neq 0$ , then the line  $\pi(u)$  is determined by the first two vectors in (1), which form the two rows of the matrix

$$\begin{pmatrix} 0 & x_0^\omega & x_1^\omega & c^{\omega/2} \\ x_0^\omega & 0 & c^{\omega/2} & x_2^\omega \end{pmatrix} \quad (3)$$

The homogeneous Plücker coordinates  $\text{Pl}(\pi(u)) = [p_{01}, p_{23}, p_{02}, p_{13}, p_{03}, p_{12}]$  of the line  $\pi(u)$  are then determined by evaluating, for  $0 \leq i < j \leq 3$ , the two-by-two sub-determinant  $p_{ij}$  of the matrix in (3) corresponding to the column indices  $i, j$  (and then, if convenient, homogenizing by any non-zero multiple in  $F$ ). It may be checked that in our case we obtain

$$\text{Pl}(\pi(u)) = [x_0^\omega, x_3^\omega, x_1^\omega, x_2^\omega, c^{\omega/2}, c^{\omega/2}] \quad (4)$$

and since by the theory explained in Chapter 12 of [19] the Plücker coordinates do not depend on a particular choice of two vectors representing a line, the formula (4) is valid also if  $x_0 = 0$ .

Recall that we need to check that the point  $[\mathbf{z}] = \pi(\ell)$  with  $z_i$  as above lies on both the lines  $\pi(u)$  and  $\pi(v)$ . We will indicate the procedure only for  $\pi(u)$  with  $\text{Pl}(\pi(u)) = [p_{01}, p_{23}, p_{02}, p_{13}, p_{03}, p_{12}]$  given by (4), as the case of  $\pi(v)$  is completely analogous. By [19, p. 538, (12.4)], the point  $[\mathbf{z}]$  lies on the line with Plücker coordinates  $\text{Pl}(\pi(u))$  if and only if

$$z_i p_{jk} + z_j p_{ik} + z_k p_{ij} = 0 \quad \text{for every } i, j, k, \quad 0 \leq i < j < k \leq 3. \quad (5)$$

It may be checked that the four equalities in (5) are indeed satisfied and the only extra tool one needs for this is total isotropy determined by the skew-symmetric form  $Q$ , that is, the identity  $x_0 y_3 + x_3 y_0 + x_1 y_2 + x_2 y_1 = 0$ . As already mentioned, checking that the point  $\pi(\ell)$  lies also on the line  $\pi(v)$  is analogous; one just needs to swap the roles of  $x_i$  and  $y_i$  ( $0 \leq i \leq 3$ ).  $\square$

For every  $x, y \in F$  we let  $f(x, y) = x^{\omega+2} + xy + y^\omega$ . The set of matrices  $M(r; a, b)$  given, for all  $r \in F^*$  and  $a, b \in F$ , by

$$M(r; a, b) = \begin{pmatrix} 1 & a & b & f(a, b) \\ 0 & r & a^\omega r & (a^{\omega+1} + b)r \\ 0 & 0 & r^{\omega+1} & ar^{\omega+1} \\ 0 & 0 & 0 & r^{\omega+2} \end{pmatrix} \quad (6)$$

is closed under multiplication and forms a group  $G$  of order  $q^2(q-1)$ . We note that the upper triangular form of  $M(r; a, b)$  in (6), suggested by our anonymous referee as a consequence of the choice of the skew-symmetric form  $Q$ , is handy to work with and differs from the one given in [19, p. 572] by applying conjugation by a permutation matrix representing the permutation (0)(132) used before. One may verify that

$$M(r; a, b)M(s; c, d) = M(rs; as + c, ac^\omega s + bs^{\omega+1} + d); \quad (7)$$

and that  $M(r; a, b)$  also admits a factorization of the form

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & r & 0 & 0 \\ 0 & 0 & r^{\omega+1} & 0 \\ 0 & 0 & 0 & r^{\omega+2} \end{pmatrix} \begin{pmatrix} 1 & a & 0 & a^{\omega+2} \\ 0 & 1 & a^\omega & a^{\omega+1} \\ 0 & 0 & 1 & a \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & b & b^\omega \\ 0 & 1 & 0 & b \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad (8)$$

so that, in our notation, the product (8) may be written in the form

$$M(r; a, b) = M(r; 0, 0)M(1; a, 0)M(1; 0, b). \quad (9)$$

Observe that the subgroup generated by  $M(1; 0, b)$  for  $b \in F$  is isomorphic to  $F^+$  and every such matrix commutes with  $M(1; a, 0)$ . The set of the latter matrices is not closed under multiplication but  $M(1; a, 0)^2 = M(1, 0, a^{\omega+1})$  and so for  $a \neq 0$  the elements  $M(1; a, 0)$  have order four. The subgroup  $H$  formed by the matrices  $M(1; a, b)$  for  $a, b \in F$ , with  $|H| = q^2$  and with multiplication  $M(1; a, b)M(1; c, d) = M(1; a + c, b + d + ac^\omega)$ , cf. (7), is also known as the Suzuki 2-group. Moreover,  $H$  is normal in  $G$  and  $G \cong H \rtimes F^*$ , the semidirect product being induced by the homomorphism  $F^* \rightarrow \text{Aut}(H)$  in which  $r \in F^*$  induces the automorphism  $(a, b) \mapsto (ar, br^{\omega+1})$  of  $H$ .

The group  $G$  acts on  $W(q)$  as a group of collineation by right multiplication. In [23] J. Tits proved that the group of all collineations of  $PG(3, q)$  leaving the set  $\Omega = \{[0, 0, 0, 1]\} \cup \{[1, x, y, f(x, y)]; x, y \in F\}$  invariant is (isomorphic to) the Suzuki group  $Sz(q) = {}^2B_2(q)$ , a simple group of order  $q^2(q^2 + 1)(q - 1)$ . Moreover, in the above representation,  $G$  is the subgroup



of  $Sz(q)$  stabilizing the point  $[0, 0, 0, 1]$ . We note that  $\Omega$  is the set of absolute points with respect to  $\pi$ , that is, the set of points  $u$  for which  $u \in \pi(u)$ ; it is also known as the Suzuki-Tits ovoid. We reiterate that to compare these facts with those in [19, p. 573-574] one needs to apply the permutation (0)(132) to all coordinates.

The action of  $G$  on the points (and hence also on the lines) of  $W(q)$  is intransitive and a straightforward calculation shows that  $G$  has the following five orbits  $O_1 - O_5$  on points of  $W(q)$ :

- $O_1 = \{[1, x, y, f(x, y)]; x, y \in F\} = \Omega \setminus \{[0, 0, 0, 1]\}$ , of size  $q^2$ ;
- $O_2 = \{[0, 1, x, y]; x, y \in F\}$ , of size  $q^2$ ;
- $O_3 = \{[0, 0, 1, x]; x \in F\}$ , of size  $q$ ;
- $O_4 = \{[0, 0, 0, 1]\}$ , which is the unique fixed point of  $G$ ; and
- $O_5 = V \setminus (\cup_{i=1}^4 O_i)$ , of size  $q^2(q - 1)$ .

We point out the (for us) important fact that  $G$  acts regularly on the orbit  $O_5$ . To see this, note that  $|G| = |O_5|$ , so that it is sufficient to show that, say, the  $G$ -orbit of the point  $[1, 0, 0, 1] \in O_5$  is the whole of  $O_5$ . Indeed, by (7) we obtain  $[1, 0, 0, 1] \cdot M(r; a, b) = [1, a, b, f(a, b) + r^{\omega+2}] \in O_5$  and it is easy to see that for distinct triples  $(r; a, b), (r'; a', b') \in F^* \times F \times F$  the points  $[1, 0, 0, 1] \cdot M(r; a, b)$  and  $[1, 0, 0, 1] \cdot M(r'; a', b')$  are distinct.

### 3. The graph arising from factorisation by polarity

Let  $A(q)$  be the graph whose vertex set  $V$  is the set of all points  $u$  of  $W(q)$ , with two distinct vertices  $u, v$  adjacent in  $A(q)$  if  $u \in \pi(v)$ , which, by properties of  $\pi$  is equivalent to  $v \in \pi(u)$ . Observe that every vertex of  $A(q)$  is adjacent to  $q + 1$  or  $q$  vertices, and the degree of a vertex  $u$  in  $A(q)$  is  $q$  if and only if  $u \in \pi(u)$ , that is, if and only if  $u$  is an element of the Suzuki-Tits ovoid, of cardinality  $q^2 + 1$ . Note that we could have defined  $A(q)$  as the quotient graph of the bipartite point-line incidence graph of  $W(q)$  obtained by factorisation by the polarity  $\pi$ , that is, by identifying  $u$  with  $\pi(u)$  throughout and suppressing eventual edges between  $u$  and  $\pi(u)$ .

A basic property of the generalised quadrangle  $W(q)$  (cf. [20]) is that for any line  $\ell$  and any point  $u$  not on  $\ell$  there is a unique point  $u' \in \ell$  such that  $u$  and  $u'$  are collinear in  $W(q)$ . This immediately translates to the observation

that the diameter of the graph  $A(q)$  is equal to 3. Indeed, suppose that  $u, v \in V$  are vertices that are not adjacent in the graph  $A(q)$ . Then, since the line  $\ell = \pi(v)$  does not contain  $u$ , by the above property there is a (unique, which is not important for this argument) point  $u' \in \ell$  such that  $u, u' \in \ell'$  for some line  $\ell' \in W(q)$ . But letting  $\pi(\ell') = v'$  we have a path  $uv'u'v$  of length 3 in  $A(q)$ . (Note that, in the above argument, the vertices  $u$  and  $v$  might correspond to collinear points of  $W(q)$ , in which case we would have  $u$  and  $v$  joined by a path of length 2 and also by a path of length 3, giving rise to a cycle of length 5 in  $A(q)$ .)

For further analysis we will need a description of the neighbourhood  $N(u)$  of a few vertices  $u$  of  $A(q)$ , that is, the set of all  $v \in V$  adjacent to  $u$  in  $A(q)$ . By the adjacency rule in  $A(q)$ , a vertex  $u = [\mathbf{x}]$  is adjacent to precisely the vertices  $v = [\mathbf{y}] \neq [\mathbf{x}]$  that are points on the line  $\pi[\mathbf{x}]$  spanned by the vectors in (1). Informally, a projective vector  $[\mathbf{x}]$  is adjacent in  $A(q)$  precisely to the projective vectors in  $\pi[\mathbf{x}]$  distinct from  $\mathbf{x}$ . A concrete identification of  $\pi[\mathbf{x}]$  can be cumbersome in general but is not for the few vertices we need. We illustrate the process on the vertex  $[\mathbf{x}] = [1, 1, 1, 1]$ . Taking the second and the fourth vector in (1) one sees that  $\pi[\mathbf{x}]$  is generated by the two (totally isotropic) projective vectors  $[1, 0, 0, 1]$  and  $[0, 1, 1, 0]$ . Excluding the self-adjacency it follows that the vertex  $[1, 1, 1, 1]$  is adjacent in  $A(q)$  to the  $q$  vertices  $[0, 1, 1, 0]$  and  $[1, z, z, 1]$  for  $z \in F \setminus \{1\}$ , or, equivalently,  $N[1, 1, 1, 1] = \{[0, 1, 1, 0]\} \cup \{[1, z+1, z+1, 1]; z \in F^*\}$ . In a completely analogous way we obtain the following table.

$u \in V$	$N(u)$
$[0, 1, 0, 0]$	$\{[0, 0, 1, 0]\} \cup \{[1, 0, z, 0]; z \in F\}$
$[0, 0, 1, 0]$	$\{[0, 0, 0, 1]\} \cup \{[0, 1, 0, z]; z \in F\}$
$[0, 0, 0, 1]$	$\{[0, 0, 1, z]; z \in F\}$
$[0, 1, 1, 0]$	$\{[0, 0, 1, 1]\} \cup \{[1, 1, z, z]; z \in F\}$
$[0, 1, 0, 1]$	$\{[0, 0, 1, 0]\} \cup \{[1, 0, z, 1]; z \in F\}$
$[1, 0, 0, 0]$	$\{[0, 1, 0, 0]\} \cup \{[1, z, 0, 0]; z \in F^*\}$
$[1, 0, 0, 1]$	$\{[0, 1, 0, 1]\} \cup \{[1, z, 1, z]; z \in F\}$
$[1, 0, 1, 1]$	$\{[0, 1, 0, 1]\} \cup \{[1, z, 1, z+1]; z \in F^*\}$
$[1, 1, 1, 1]$	$\{[0, 1, 1, 0]\} \cup \{[1, z+1, z+1, 1]; z \in F^*\}$

Table 1: Neighbourhood of selected vertices of  $A(q)$ .

As the next step we show that the group  $G$  generated by the matrices  $M = M(r; a, b)$  introduced in (6) acts on the vertices of  $A(q)$  by right mul-

tiplication. This is clearly equivalent to the statement that  $\pi[\mathbf{x}M] = \pi[\mathbf{x}]M$  for every  $[\mathbf{x}] \in V$ . To verify this it is sufficient to restrict ourselves to a particular  $[\mathbf{z}]$  from each orbit  $O \in \{O_1, \dots, O_5\}$  listed in section 2. Indeed, if we choose a  $[\mathbf{z}] \in O$ , then, for every  $[\mathbf{x}] \in O$  we have  $[\mathbf{x}] = [\mathbf{z}M_z]$  for some  $M_z \in G$ . Provided we show that  $\pi[\mathbf{z}M'] = \pi[\mathbf{z}]M'$  for every  $M' \in G$ , we then have  $\pi[\mathbf{x}M] = \pi[\mathbf{z}M_zM] = \pi[\mathbf{z}]M_zM = \pi[\mathbf{z}M_z]M = \pi[\mathbf{x}]M$  for every  $M \in G$ . The verification can now be done by letting  $[\mathbf{z}]$  be the representatives  $[1, 0, 0, 0]$ ,  $[0, 1, 0, 0]$ ,  $[0, 0, 1, 0]$ ,  $[0, 0, 0, 1]$  and  $[1, 0, 0, 1]$  of the orbits  $O_1$  to  $O_5$ , respectively.

We illustrate the procedure on the computationally most demanding case when  $[\mathbf{z}] = [1, 0, 0, 1]$ . Instead of proving  $\pi[\mathbf{z}M] = \pi[\mathbf{z}]M$  we prove the equivalent equality  $[\mathbf{z}]M = \pi[\pi[\mathbf{z}]M]$ . Looking at Table 1 we see that both vertices  $[0, 1, 0, 1]$  and  $[1, 0, 1, 0]$  are neighbours of  $[\mathbf{z}] = [1, 0, 0, 1]$  in  $A(q)$ . This is equivalent to stating that  $\pi[\mathbf{z}]$  is the line of  $W(q)$  through the points  $[\mathbf{x}] = [0, 1, 0, 1]$  and  $[\mathbf{y}] = [1, 0, 1, 0]$ . Since  $G$  is a collineation group of  $W(q)$ , it follows that for every  $M = M(r; a, b)$  from (6) the point  $[\mathbf{z}]M = [1, a, b, f(a, b) + r^{\omega+2}]$  lies on the line  $\ell = \pi[\mathbf{z}]M$  through the points  $[\mathbf{x}]M = [0, r, a^\omega r, r^{\omega+2} + r(a^{\omega+1} + b)]$  and  $[\mathbf{y}]M = [1, a, b + r^{\omega+1}, f(a, b) + ar^{\omega+1}]$ . We now calculate the point  $\pi(\ell)$  by the procedure described in (2). A somewhat lengthy but straightforward verification shows that  $\delta_{01}^{\omega/2} = r^{\omega/2}$ ,  $\delta_{02}^{\omega/2} = r^{\omega/2}a$ ,  $\delta_{13}^{\omega/2} = r^{\omega/2}b$  and  $\delta_{23}^{\omega/2} = r^{\omega/2}(f(a, b) + r^{\omega+2})$ , so that  $[\delta_{02}^{\omega/2}, \delta_{31}^{\omega/2}, \delta_{03}^{\omega/2}, \delta_{21}^{\omega/2}] = [1, a, b, f(a, b) + r^{\omega+2}]$ . Thus,  $\pi(\ell) = [\mathbf{z}]M$  and hence  $\pi[\pi[\mathbf{z}]M] = [\mathbf{z}]M$ , which is what we wanted to establish. We leave out the details for the remaining choices of orbit representatives as they are similar (and easier).

#### 4. Strategy: Focusing on the subgraph induced by the regular orbit

From this point on we will focus on the subgraph  $B(q)$  of  $A(q)$  induced by the subset  $O_5$  of  $V$ . Since the group  $G$  acts regularly on the vertex set of  $B(q)$ , it follows [21] that  $B(q)$  is a Cayley graph for the group  $G$  and some generating set  $S$  for  $G$ . The procedure of turning a graph with a regular subgroup of its automorphism group into a Cayley graph is well known; the set  $S$  is uniquely determined and can be recovered by looking at the ‘local’ action of  $G$  on an arbitrarily chosen vertex. We will give details below but precede everything by outlining the general strategy of our proof.

**Strategy.** For an infinite set of degrees  $d$  we will construct a Cayley graph of degree  $d$ , order  $d^3 - O(d^{2.5})$  as follows:

(a) We will explicitly determine an isomorphism of  $B(q)$  onto a Cayley graph  $C(G, S)$  for our group  $G$  acting regularly on the vertex set  $O_5$  of  $B(q)$  (Lemma 4.1); this graph will have degree  $|S| = q - 1$  and order  $q^2(q - 1)$  for  $q = 2^{2n+1}$  but its diameter will still be larger than 3.

(b) We show that the generating set  $S$  can be extended to a set  $S^*$  by adding no more than  $\lceil \sqrt{q-1} \rceil + 4$  generators in such a way that the resulting Cayley graph  $Cay(G, S^*)$  will have the required properties; namely, it will have degree  $d = q + O(\sqrt{q})$ , order  $q^2(q - 1) = d^3 - O(d^{2.5})$ , and diameter 3 (by a combination of Lemma 4.2 and Theorem 5.1).

We begin by developing the item (a). Let  $u = [\mathbf{x}]$  be a fixed vertex of  $B(q)$ ; for later convenience we will fix the vertex  $u = [1, 0, 0, 1]$  of  $B(q)$  throughout. By the regular action of  $G$  on vertices of  $B(q)$ , for every neighbour  $w$  of  $u$  in  $B(q)$  there is exactly one  $g_w \in G$  represented by a matrix  $M_w$  as in (6) such that  $w = ug_w = [\mathbf{x}]M_w$ . To see what happens with adjacency in an arbitrary vertex, just apply an arbitrary element  $g \in G$  to this situation to conclude that a vertex  $ug$  is adjacent to  $wg = u(g_w g)$  for every neighbour  $w$  of  $u$ . Let now  $S$  be the set of all the  $g_w \in G$  where  $w$  ranges over all neighbours of  $u$  in  $B(q)$ . The regular action of  $G$  enables us to identify  $G$  with the vertex set of  $B(q)$  by means of the bijection  $g \mapsto ug$  and the same bijection gives an isomorphism of the Cayley graph  $C(G, S)$  onto  $B(q)$ ; note that this isomorphism maps the identity of  $G$  onto the fixed vertex  $u$ . The adjacency rule in  $C(G, S)$  follows from the above, namely,  $g \in G$  is adjacent to  $g_w g$  for every  $g_w \in S$ .

By Table 1 and the description of the orbit  $O_5$ , the neighbourhood of  $u$  in  $B(q)$  is the set  $\{[1, x, 1, x]; x \in F, x \neq 1\}$  of size  $q - 1$ . The elements  $g_w \in S$  are thus matrices  $M(r; a, b)$  from (6) such that  $[1, 0, 0, 1]M(r; a, b) = [1, x, 1, x]$ ,  $x \neq 1$ . A quick calculation reveals that we must have  $b = 1$ , and  $a, r$  are tied by the equation  $a^{\omega+2} + r^{\omega+2} = 1$ . This equation has a unique solution  $a = a(r) \neq 1$  for every  $r \in F^*$ , given by  $a(r) = (1 + r^{\omega+2})^{1-\omega/2}$  (recall that every element in  $F^*$  has a unique square root). One can verify that  $[1, 0, 0, 1]M(r; a(r), 1) = [1, a, 1, a]$ ,  $a \neq 1$ . We have therefore identified the generating set as  $S = \{M(r; a(r), 1); r \in F^*\}$ ; this set is closed under taking inverses by the above construction. Summing up, we have:

**Lemma 4.1.** *The graph  $B(q)$  is isomorphic to the Cayley graph  $C(G, S)$  of degree  $q - 1$  with the generating set  $S = \{M(r; a(r), 1); r \in F^*\}$ .  $\square$*

In contrast to  $A(q)$  the diameter of  $B(q)$  is larger than 3. This is a consequence of the following fact. Assume that  $v$  and  $w$  are vertices of  $A(q)$  such that  $v$  represents a point of  $W(q)$  and  $w$  a line of  $W(q)$  not containing the point and such that  $v$  and  $\pi(w)$  are not on a line of  $W(q)$ . Then, by properties of generalised quadrangles, there is a unique path from  $v$  to  $w$  of length 3 in  $A(q)$ , and no shorter path, and removing vertices from  $A(q)$  will destroy some of these paths. Our aim is to extend the generating set  $S$  of the Cayley graph  $C(G, S) \simeq B(q)$  by just a ‘few’ new generators to ensure that the new Cayley graph (which will contain  $B(q)$  as a spanning subgraph) has diameter 3. We thus need to identify pairs of vertices of  $B(q)$  that could end up at distance larger than 3 after the removal of the sets  $O_1 - O_4$  from  $A(q)$ . Since  $B(q)$  is a Cayley graph, it is sufficient to do this for pairs of vertices  $u, v \in O_5$  in which  $u = [1, 0, 0, 1]$  is our fixed vertex.

In the considerations below we will frequently refer to Table 1 without explicit alerts. In the graph  $A(q)$ , our fixed vertex  $u = [1, 0, 0, 1] \in O_5$  is adjacent to the two vertices  $u_1 = [1, 1, 1, 1] \in O_1$ ,  $u_2 = [0, 1, 0, 1] \in O_2$ , and has  $q - 1$  neighbours inside  $B(q)$ , all of the form  $[1, x, 1, x]$  for  $x \in F$ ,  $x \neq 1$ . The neighbour  $u_1 = [1, 1, 1, 1] \in O_1$  of  $u$  in  $A(q)$  is adjacent to  $u'_2 = [0, 1, 1, 0] \in O_2$  and to another  $q - 1$  vertices  $[1, z+1, z+1, 1]$ ,  $z \in F^*$ , in  $B(q)$ . It follows that when restricting to  $B(q)$  we lose the paths of length 2 from  $u$  via  $u_1$  to the  $q - 2$  vertices in the set  $L(u_1) := N(u_1) \setminus \{u, u'_2\} = \{[1, z+1, z+1, 1]; z \in F^* \setminus \{1\}\}$ . Similarly, the neighbour  $u_2 \in O_2$  of  $u$  in  $A(q)$  is adjacent to  $u_3 = [0, 0, 1, 0] \in O_3$ ,  $u'_1 = [1, 0, 1, 1] \in O_1$ , and has further  $q - 1$  neighbours  $[1, 0, z+1, 1]$ ,  $z \in F^*$ , in  $B(q)$ . Again, when restricting to  $B(q)$  we lose the paths of length 2 from  $u$  through  $u_2$  to the  $q - 2$  vertices in the set  $L(u_2) := N(u_2) \setminus \{u, u_3, u'_1\} = \{[1, 0, z+1, 1]; z \in F^* \setminus \{1\}\}$ .

The vertex  $u'_1$  is adjacent in  $A(q)$  to  $u_2 = [0, 1, 0, 1] \in O_2$  and to the  $q - 1$  vertices forming the set  $L(u'_1) := N(u'_1) \setminus \{u_2\} = \{[1, z, 1, z+1]; z \in F^*\}$ . The neighbourhood of  $u'_2$  in  $A(q)$  consists of  $u_1 = [1, 1, 1, 1] \in O_1$ ,  $u'_3 = [0, 0, 1, 1] \in O_3$ , and the  $q - 1$  vertices in the set  $L(u'_2) := N(u'_2) \setminus \{u_1, u'_3\} = \{[1, 1, z+1, z+1]; z \in F^*\}$ . This implies that when restricting our attention to the graph  $B(q)$  we lose the paths of length 3 of the form  $uu_1u'_2w$  for  $w \in L(u'_2)$  and  $uu_2u'_1w$  for  $w \in L(u'_1)$ .

We now pass onto the item (b) of the outlined strategy. So far we have identified four sets of vertices,  $L(u_1)$ ,  $L(u_2)$ ,  $L(u'_1)$  and  $L(u'_2)$ , to which we lose paths from  $u$  through  $u_1$  and  $u_2$  when considering  $B(q)$ ; we note that no vertex in  $B(q)$  is adjacent to  $u_3$ . For convenience of the reader and for later use in this and the forthcoming section we list the vertices  $u_i$ ,  $u'_i$  and the sets

$L(u_i)$  and  $L(u'_i)$  for  $i = 1, 2$  in a tabular form.

$u_1 = [1, 1, 1, 1] \in O_1$	$L(u_1) = \{[1, z+1, z+1, 1]; z \in F^* \setminus \{1\}\}$
$u'_1 = [1, 0, 1, 1] \in O_1$	$L(u'_1) = \{[1, z, 1, z+1]; z \in F^*\}$
$u_2 = [0, 1, 0, 1] \in O_2$	$L(u_2) = \{[1, 0, z+1, 1]; z \in F^* \setminus \{1\}\}$
$u'_2 = [0, 1, 1, 0] \in O_2$	$L(u'_2) = \{[1, 1, z+1, z+1]; z \in F^*\}$

Table 2: Vertices  $u_i$ ,  $u'_i$  and the sets  $L(u_i)$  and  $L(u'_i)$  for  $i = 1, 2$ .

In order to make up for the loss of the paths mentioned above we will suitably extend the generating set  $S$  in the Cayley graph  $C(G, S) \simeq B(q)$ . To do so we first determine the action of  $G$  on the four sets, which is equivalent to determining the vertex stabilisers in  $G$  of  $u_1$ ,  $u_2$ ,  $u'_1$  and  $u'_2$ .

**Lemma 4.2.** *The stabilisers in  $G$  of  $u_1$ ,  $u_2$ ,  $u'_1$  and  $u'_2$  are all cyclic and isomorphic to  $F^*$ . In more detail:*

- $Stab_G(u_1) = Stab_G(u'_2) = \{M(r; r+1, r+1); r \in F^*\} \simeq F^*$ , acting regularly on both  $L(u_1) \cup \{u\}$  and  $L(u'_2)$  by  $[1, z+1, z+1, 1] \cdot M(r; r+1, r+1) = [1, zr+1, zr+1, 1]$  and  $[1, 1, z+1, z+1] \cdot M(r; r+1, r+1) = [1, 1, zr^{\omega+1} + 1, zr^{\omega+1} + 1]$ .
- $Stab_G(u_2) = Stab_G(u'_1) = \{M(r; 0, 1+r^{\omega+1}); r \in F^*\} \simeq F^*$ , acting regularly on both sets  $L(u_2) \cup \{u\}$  and  $L(u'_1)$  by  $[1, 0, z+1, 1] \cdot M(r; 0, 1+r^{\omega+1}) = [1, 0, zr^{\omega+1}, 1]$  and  $[1, z, 1, z+1] \cdot M(r; 0, 1+r^{\omega+1}) = [1, zr, 1, zr + 1]$  for  $z \in F^*$ .

**Proof.** Calculation of the stabilisers from the equation  $[\mathbf{x}]M(r; a, b) = [\mathbf{x}]$  for  $[\mathbf{x}] \in \{u_1, u'_2, u_2, u'_1\}$  and solving for  $a, b$  in terms of  $r$  is straightforward. Regarding isomorphism of the stabilisers with  $F^*$ , for  $Stab_G(u_1) = Stab_G(u'_2)$  it is given by  $\theta_1 : r \mapsto M(r; r+1, r+1)$  since one can check that  $M(r; r+1, r+1)M(s; s+1, s+1) = M(rs; rs+1, rs+1)$  for all  $r, s \in F^*$ . The same type of isomorphism  $\theta_2 : r \mapsto M(r; 0, 1+r^{\omega+1})$  works for  $Stab_G(u_2) = Stab_G(u'_1)$ , as one can easily verify that  $M(r; 0, 1+r^{\omega+1})M(s; 0, 1+s^{\omega+1}) = M(rs; 0, 1+(rs)^{\omega+1})$  for every  $r, s \in F^*$ . The actions of the stabilisers on the four sets are obvious.  $\square$

## 5. Large Cayley graphs of diameter 3 and small degree

For any integer  $m \geq 3$  let  $c_m$  denote the smallest number of elements in a generating set  $X$  of a Cayley graph of a cyclic group  $C(Z_m, X)$  of diameter two. The best available general upper bound on  $c_m$ , which is, of course, the degree of  $C(Z_m, X)$ , is  $c_m \leq 2\lceil\sqrt{m}\rceil$ , see [8] for a short proof. Applying this to the cyclic group  $F^*$  of order  $m = q - 1$  we have the existence of a Cayley graph  $C(F^*, X)$  of diameter 2 with  $|X| = c_{q-1} \leq 2\lceil\sqrt{q-1}\rceil$ . Taking images of  $C(F^*, X)$  under the isomorphisms  $\theta_1$  and  $\theta_2$  from the proof of Lemma 4.2, that is, letting  $S_1 = \theta_1(X)$  and  $S_2 = \theta_2(X)$  and denoting  $H_1 = \text{Stab}_G(u_1) = \text{Stab}_G(u'_2)$  and  $H_2 = \text{Stab}_G(u_2) = \text{Stab}_G(u'_1)$ , we have constructed Cayley graphs  $C(H_1, S_1)$  and  $C(H_2, S_2)$  of degree  $c_{q-1}$  and diameter 2.

We are now ready to describe a suitable ‘small’ extension of our generating set  $S$  from Lemma 4.1 in the Cayley graph  $C(G, S) \simeq B(q)$  to obtain a graph of diameter three. Let  $M_1 = M(1; 1, 1)$ ,  $M_2 = M(1; 1, 0)$ ,  $S_3 = \{M_1, M_1^{-1}, M_2, M_2^{-1}\}$  and let  $S^* = S \cup S_1 \cup S_2 \cup S_3$ ; note that  $S^*$  is closed under taking inverses.

**Theorem 5.1.** *For every  $n \geq 1$  and  $q = 2^{2n+1}$  the Cayley graph  $C(G, S^*)$  of order  $q^2(q-1)$  and degree  $q + 2c_{q-1} + 3$  has diameter 3.*

**Proof.** The Cayley graph  $C(G, S^*)$  has order  $|G| = q^2(q-1)$ , degree  $|S|$  that evaluates to  $q + 2c_{q-1} + 3$ , and it contains the Cayley graph  $C(G, S) \simeq B(q)$  as a spanning subgraph. We may thus identify the vertex set of  $C(G, S^*)$  with the orbit  $O_5$  of the group  $G$  on the set  $V$  as we did in the isomorphism from  $C(G, S)$  onto  $B(q)$ .

We proceed by extending some of our earlier observations made for the fixed vertex  $u$  to arbitrary vertices of  $B(q)$ . By the action of  $G$  on vertices of the entire graph  $A(q)$  and by the regular action of the same group on the vertex set  $O_5$  of the graph  $B(q)$ , every vertex  $v \in O_5$  has one neighbour  $v_1 \in O_1$ , one neighbour  $v_2 \in O_2$ , and  $q-1$  neighbours within  $B(q)$ . Further, in  $A(q)$ , the vertex  $v_1$  is adjacent to one vertex  $v'_2 \in O_2$  and to a set  $L(v_1)$  of  $q-2$  vertices in  $B(q)$  distinct from  $v$ , and the vertex  $v_2$  has one neighbour  $v_3 \in O_3$ , another neighbour  $v'_1 \in O_1$  and  $q-2$  neighbours in a set  $L(v_2)$  of vertices of  $B(q)$  distinct from  $v$ . Moreover, by our isomorphism from the Cayley graph  $C(G, S)$  onto  $B(q)$  constructed earlier, the subsets  $L(v_1)$  and  $L(v_2)$  of  $O_5$  are  $g$ -images of the sets  $L(u_1)$  and  $L(u_2)$  for the element  $g \in G$  that carries  $u$  onto  $v = ug$ . Also, note that the unique vertex  $[0, 0, 0, 1]$  of  $O_4$

of degree  $q$  is incident only to  $q$  vertices in  $O_3$  and the neighbours of every vertex of  $O_3$  (except  $[0, 0, 0, 1]$ ) lie in  $O_2$ .

Equipped with this we now prove that the diameter of  $C(G, S^*)$  is 3. We know that it is sufficient to check distances from our fixed vertex  $u = [1, 0, 0, 1]$ . Since the diameter of  $A(q)$  is 3, it is sufficient to show that for every path  $P$  of length at most 3 in  $A(q)$  from  $u$  to an arbitrary vertex  $w \in O_5$  and passing through a vertex outside  $O_5$  the distance between  $u$  and  $w$  is still at most 3 in  $C(G, S^*)$ . By our earlier examination of possibilities, and referring also to Table 2 and the notation introduced in the previous paragraph, such a path  $P$  can only have one of the following eight forms:

- 1)  $P = uu_1w$ , with  $w \in L(u_1)$ ,
- 2)  $P = uu_2w$ , with  $w \in L(u_2)$ ,
- 3)  $P = uu_1w'w$  for some  $w' \in L(u_1)$ , with  $w \notin L(u_1) \cup \{u\}$ ,
- 4)  $P = uu_2w'w$  for some  $w' \in L(u_2)$ , with  $w \notin L(u_2) \cup \{u\}$ ,
- 5)  $P = uu_1u'_2w$ , with  $w \in L(u'_2)$ ,
- 6)  $P = uu_2u'_1w$ , with  $w \in L(u'_1)$ ,
- 7)  $P = uvv_1w$  for some  $v \in O_5$ , with  $w \in L(v_1)$ , or
- 8)  $P = uvv_2w$  for some  $v \in O_5$ , with  $w \in L(v_2)$ .

Since  $S_1, S_2 \subset S^*$ , we may apply Lemma 4.2 to conclude that the subgraphs of  $C(G, S^*)$  induced by the subsets  $L(u_1) \cup \{u\}$  and  $L(u_2) \cup \{u\}$  are isomorphic to the Cayley graphs  $C(H_1, S_1)$  and  $C(H_2, S_2)$ , both of diameter 2. This implies that the distance between  $u$  and  $w$  is at most 3 in the cases 1) – 4). Similarly, since  $H_2$  is also equal to  $Stab_G(u'_1)$  and  $H_1$  is equal to  $Stab_G(u'_2)$ , by Lemma 4.2 (which also describes the action of  $H_2$  and  $H_1$ ) the distances in  $C(G, S^*)$  between vertices in the sets  $L(u'_1)$  and  $L(u'_2)$  are at most 2. Having included the elements  $M_1 = M(1; 1, 1)$  and  $M_2 = M(1; 1, 0)$  in  $S^*$ , we have an edge from  $u = [1, 0, 0, 1]$  to  $uM_1 = [1, 1, 1, 0] \in L(u'_1)$  and an edge from  $u$  to  $uM_2 = [1, 1, 0, 0] \in L(u'_2)$ . It follows that we have paths of length 3 from  $u$  to  $w$  inside  $C(G, S^*)$  also in the cases 5) and 6).

Let  $v$  be a vertex as in the cases 7) and 8), incident with  $u$  in  $B(q)$ . By the regular action of  $G$  on the vertex set of  $B(q)$  there exists a (unique)  $g \in G$  such that  $v = ug$ . From the previous paragraph we know that, for  $i = 1, 2$ , the subgraph of  $C(G, S^*)$  induced by the vertex set  $L(u_i) \cup \{u\}$  is isomorphic to the Cayley graph  $C(H_i, S_i)$ . But since  $L(v_i) \cup \{v\} = (L(u_i) \cup \{u\})g$  for  $i = 1, 2$ , the subgraph of  $C(G, S^*)$  induced by the set  $L(v_i) \cup \{v\}$  is isomorphic to the one induced by the set  $L(u_i) \cup \{u\}$ , which is the Cayley graph  $C(H_i, S_i)$ . It follows that in the cases 7) and 8), for  $i = 1, 2$ , the  $vv_iw$ -part of the path  $P$  can be replaced by a path of length at most two in the subgraph of  $C(G, S^*)$



induced by the set  $L(v_i) \cup \{v\}$ , since the diameter of  $C(H_i, S_i)$  is 2. This completes the proof.  $\square$

Letting  $d = q + 2c_{q-1} + 3 \leq q + 4\lceil\sqrt{q-1}\rceil + 3$  for  $q = 2^{2n+1}$  and asymptotically expressing the order  $q^2(q-1)$  of  $C(G, S^*)$  in terms of  $d$ , we obtain:

**Corollary 5.2.** *There exists an infinite increasing sequence of values of  $d$  for which there are Cayley graphs of degree  $d$ , diameter 3 and order  $d^3 - O(d^{2.5})$  as  $d \rightarrow \infty$ .*  $\square$

We thus have  $\limsup_{d \rightarrow \infty} \text{Cay}(d, 3)/M(d, 3) = 1$ , that is, the Moore bound for diameter 3 can be asymptotically approached by Cayley graphs.

## 6. Remarks

Our construction of an infinite sequence of Cayley graphs of diameter 3, degree  $q + o(q)$  and order  $q^3 - o(q^3)$  was based on the existence of a group  $G$  of order  $q^2(q-1)$  for  $q = 2^{2n+1}$ , regular on the vertex set of a subgraph obtained from the point-line incidence graph of a generalised quadrangle  $W(q)$  by factorisation by a polarity. Moreover, the group  $G$  is (isomorphic to) a subgroup of index  $q^2 + 1$  of the Suzuki group  $\text{Sz}(q) = {}^2B_2(q)$ , a simple group of order  $q^2(q^2+1)(q-1)$ , acting doubly transitively on the Suzuki-Tits ovoid (of order  $q^2 + 1$ ) in the generalised quadrangle  $W(q)$ ; see [17] for more details.

It is tempting to consider an analogous approach for constructing an infinite sequence of Cayley graphs of diameter 5, degree  $q + o(q)$  and order  $q^5 - o(q^5)$  from a suitable group acting regularly on some subgraph obtained from the incidence graph  $I(q)$  of a *generalised hexagon*  $H(q)$  factored by a polarity. By known results summarised in [17], a generalised hexagon  $H(q)$  admits a polarity if and only if  $q = 3^{2n+1}$ ,  $n \geq 1$ . For such  $q$ , the graph  $I(q)$ , of diameter 5 and order  $q^5 + \dots + q + 1$ , has  $q^5 + q^4 + q^2 + q$  vertices of degree  $q + 1$  together with the  $q^3 + 1$  vertices of degree  $q$  that correspond to the Ree-Tits ovoid formed by the absolute (that is, self-polar) points of  $H(q)$ . It turns out that, up to field automorphisms, the automorphism group of  $I(q)$  can be identified with the collineation group that preserves the Ree-Tits ovoid, which is the Ree group  $\text{Re}(q) = {}^2G_2(q)$ .

Unfortunately, by the classification of maximal subgroups of the Ree groups [13, 12], the group  $\text{Re}(q) = {}^2G_2(q)$ , which is a simple group of order  $q^3(q^3+1)(q-1)$ , does not contain a subgroup of order  $O(q^5)$  for  $q \rightarrow \infty$ .

The approach that works for construction of large Cayley graphs of diameter 2 from generalised triangles in [1], and of diameter 3 from generalised quadrangles in this article, thus does not carry over to an analogous construction for diameter 5 from generalised hexagons.

**Acknowledgements.** The authors thank an anonymous referee for carefully reading the manuscript and insightfully suggesting a number of improvements regarding presentation of results on generalized quadrangles.

All three authors acknowledge support from the research grants APVV 0136-12, APVV-15-0220, VEGA 1/0026/16 and VEGA 1/0142/17.

- [1] M. Bachratý and J. Širáň, Polarity graphs revisited, *Ars Math. Contemp.* 8 (2015) 55–67.
- [2] E. Bannai and T. Ito, On finite Moore graphs, *J. Fac. Sci. Tokyo Univ.* 20 (1973) 191–208.
- [3] J.-C. Bermond and B. Bollobás, The diameter of graphs: A survey, *Congressus Numerantium* 32 (1981) 3–27.
- [4] E. Canale and J. Gómez, Superfluous edges and exponential expansions of deBruijn and Kautz digraphs, *Discr. Appl. Math.* 138 (2004) 303–331.
- [5] R.M. Damerell, On Moore graphs, *Proc. Cambridge Phil. Soc.* 74 (1973) 227–236.
- [6] C. Delorme, Large bipartite graphs with given degree and diameter, *J. Graph Theory* 8 (1985) 325–334.
- [7] C. Delorme, Examples of products giving large graphs with given degree and diameter, *Discrete Applied Mathematics* 37/38 (1992) 157–167.
- [8] G. Erskine, Diameter 2 Cayley graphs of dihedral groups, *Discrete Math.* 338 (2015) 1022–1024.
- [9] G. Exoo, R. Jajcay, M. Mačaj and J. Širáň, The defect of vertex-transitive graphs of given degree and diameter, submitted (2014).
- [10] V. Faber, J. W. Moore and W. Y. C. Chen, Cycle prefix digraphs for symmetric interconnection networks, *Networks* 23 (1993) 641–649.

- [11] A. J. Hoffman and R. R. Singleton, On Moore graphs with diameter 2 and 3, *IBM J. Res. Develop.* 4 (1960) 497–504.
- [12] P. B. Kleidman, The maximal subgroups of the Chevalley groups  $G_2(q)$  with  $q$  odd, the Ree groups  ${}^2G_2(q)$ , and their automorphism groups, *J. Algebra* 117 (1988) 30–71.
- [13] V. M. Levchuk and Ya. N. Nuzhin, The structure of Ree groups, *Algebra i Logika* 24 (1985) 26–41 (Russian); English translation, *Algebra and Logic* 24 (1985) 16–26.
- [14] E. Loz and J. Širáň, New record graphs in the degree-diameter problem, *Australasian J. Combin.* 41 (2008) 63–80.
- [15] H. Macbeth, J. Šiagiová and J. Širáň, Cayley graphs of given degree and diameter for cyclic, Abelian, and metacyclic groups, *Discrete Math.* 312 (2012) 94–99.
- [16] H. Macbeth, J. Šiagiová, J. Širáň and T. Vetrík, Large Cayley graphs and vertex-transitive non-Cayley graphs of given degree and diameter, *J. Graph Theory* 64 (2010) 87–98.
- [17] H. van Maldeghem, *Generalized polygons* (Birkhäuser, 1998).
- [18] M. Miller and J. Širáň, Moore graphs and beyond: A survey of the degree/diameter problem, *Electr. J. Combin.* 20 (2) (2013), Dynamic Survey DS14v2, 92 pp.
- [19] S. E. Payne, *Topics in Finite Geometry: Ovals, Ovoids and Generalized Quadrangles* (University of Colorado, Denver, 2007).
- [20] S. E. Payne and J. A. Thas, *Finite Generalized Quadrangles* (European Mathematical Society, 2009).
- [21] G. Sabidussi, On a class of fixed-point free graphs, *Proc. Amer. Math. Soc.* 9 (1958) 800–804.
- [22] J. Šiagiová and J. Širáň, Approaching the Moore bound for diameter two by Cayley graphs, *J. Combin. Theory Ser. B* 102 (2012) 470–473.
- [23] J. Tits, Ovoids et Groupes de Suzuki, *Arch. Math.* 13 (1962) 187–198.

- [24] T. Vetrík, Cayley graphs of given degree and diameters 3, 4 and 5, *Discrete Math.* 313 (2013) 213-216.
- [25] *Online tables of current largest graphs for the degree-diameter problem:*  
[www.eyal.com.au/wiki/](http://www.eyal.com.au/wiki/)