

# Open Research Online

---

The Open University's repository of research publications and other research outputs

## Completing the design spectra for graphs with six vertices and eight edges

### Journal Item

How to cite:

Forbes, Anthony D.; Griggs, Terry S. and Forbes, Kim A. (2018). Completing the design spectra for graphs with six vertices and eight edges. *Australasian Journal of Combinatorics*, 70(3) pp. 386–389.

For guidance on citations see [FAQs](#).

© [not recorded]

Version: Version of Record

Link(s) to article on publisher's website:

[https://ajc.maths.uq.edu.au/pdf/70/ajc.v70\\_p386.pdf](https://ajc.maths.uq.edu.au/pdf/70/ajc.v70_p386.pdf)

---

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

---

[oro.open.ac.uk](https://oro.open.ac.uk)

# Completing the design spectra for graphs with six vertices and eight edges

ANTHONY D. FORBES    TERRY S. GRIGGS

*School of Mathematics and Statistics  
The Open University  
Walton Hall, Milton Keynes MK7 6AA  
U.K.*

anthony.d.forbes@gmail.com    terry.griggs@open.ac.uk

KIM A. FORBES

*Research, Business and Innovation  
Kingston University  
Penrhyn Road, Kingston upon Thames, KT1 2EE  
U.K.*

kimaforbes@gmail.com

## Abstract

Apart from two possible exceptions, the design spectrum has been determined for every graph with six vertices and at most eight edges. The purpose of this note is to establish the existence of the two missing designs, both of order 32.

## 1 Introduction

Let  $G$  be a simple graph. If the edge set of a simple graph  $K$  can be partitioned into edge sets of graphs each isomorphic to  $G$ , we say that there exists a *decomposition* of  $K$  into  $G$ . In the case where  $K$  is the complete graph  $K_n$  we refer to the decomposition as a  $G$ -*design* of order  $n$ . The *design spectrum* of  $G$  is the set of non-negative integers  $n$  for which there exists a  $G$ -design of order  $n$ .

The design spectrum problem for small graphs has attracted attention. In particular, it has been solved for (i) all graphs with at most five vertices, (ii) all graphs with six vertices and at most seven edges, (iii) all graphs with six vertices and eight edges, with two possible exceptions, and (iv) eleven of the 21 graphs with six vertices and nine edges. See [1] and [2] for details and references. More recently, the spectrum problem was resolved for all of the remaining ten graphs with six vertices and nine edges, [4].

Figure 1: Graphs  $M_1$  and  $M_2$





Figure 2: Graph  $Q$



## 2 Design construction

The two possible exceptions for graphs with six vertices and eight edges are illustrated in Figure 1, and in each case it is order 32 that remains unresolved; see [5], where the graphs are denoted by  $M_1$  and  $M_2$ . To complete the task of determining the design spectra for 6-vertex, 8-edge graphs it suffices merely to exhibit designs of order 32 for  $M_1$  and  $M_2$ . However, it might be of some interest to explain how they were obtained; so we give brief details.

The number of graphs in each design is 62 and without a vertex of degree 1 there is no convenient automorphism that we can exploit. So we are left in each case with the task of assembling sixty-two 6-tuples of numbers from the set  $N = \{0, 1, \dots, 31\}$ .

Let  $Q$  denote the graph obtained by removing an edge from a  $K_4$  (Figure 2). Our primary objective is to obtain a decomposition  $\mathcal{D}$  of  $K_{32}$  labelled with  $N$  into 62 triangles and 62 copies of  $Q$ , also labelled from  $N$ .

A *partial Steiner triple system*  $\text{PSTS}(x)$  consists of a *point set*,  $X$  say, of cardinality  $x$  and a collection of *triples*, 3-element subsets of  $X$ , such that each unordered pair of elements from  $X$  occurs in at most one triple. We create by hill climbing, [6] (or see [3, §2.7.2]), a random  $\text{PSTS}(32)$ ,  $\mathcal{S}$ , with point set  $N$  and 62 triples. The triples of  $\mathcal{S}$  are the triangles of  $\mathcal{D}$ .

Next, we create the  $T$ - $K$  matrix  $\mathbf{M}$  for assembling the graphs  $Q$ . Denote by  $T$  the set of edges of the labelled  $K_{32}$  that are not present as pairs in the triples of  $\mathcal{S}$ .

To each ordered 4-tuple  $k = (a, b, c, d)$ ,  $a, b, c, d \in N$ , we associate a graph  $Q_k \cong Q$  with its vertices labelled by the elements of  $k$  as shown in Figure 2. Let  $K$  be the set of ordered 4-tuples  $k$  corresponding to distinct labelled  $Q_k$  such that all the edges of  $Q_k$  belong to  $T$ . The rows of  $\mathbf{M}$  are indexed by elements of  $T$  and the columns by elements of  $K$ . The matrix is defined by  $\mathbf{M}_{t,k} = 1$  if  $Q_k$  contains edge  $t$ ; otherwise  $\mathbf{M}_{t,k} = 0$ .

We attempt to solve  $\mathbf{M}\mathbf{v} = \mathbf{1}$  for  $\mathbf{v}$  a  $\{0, 1\}$  vector. If we are unsuccessful, we try another PSTS(32). Otherwise we recover the 4-tuples corresponding to 1s in  $\mathbf{v}$  and hence the graphs  $Q$  of our decomposition  $\mathcal{D}$ .

Finally, for  $\delta = 2, 3$  we pair off triangles and  $Q$ s in  $\mathcal{D}$  such that in each pair there is a common label attached to the triangle and a vertex of degree  $\delta$  in the  $Q$ . Thus we have the required decompositions of  $K_{32}$  into  $M_1$  when  $\delta = 3$  and  $M_2$  when  $\delta = 2$ . The results are presented in Section 3. Incidentally, by pairing triangles and  $Q$ s with no common vertices we obtain a  $Q \cup K_3$  design of order 32.

### 3 The designs

We represent the graphs by ordered 6-tuples  $(a, b, c, d, e, f)_{M_1}$  and  $(a, b, c, d, e, f)_{M_2}$ , where the letters correspond to vertices as illustrated in Figure 1. With vertex set  $Z_{32}$  the decompositions consist of 62 graphs each:

- $(8, 9, 3, 16, 23, 0)_{M_1}$ ,  $(5, 8, 6, 27, 25, 1)_{M_1}$ ,  $(20, 7, 0, 27, 1, 16)_{M_1}$ ,
- $(3, 19, 5, 20, 6, 0)_{M_1}$ ,  $(14, 20, 5, 22, 1, 4)_{M_1}$ ,  $(20, 21, 18, 31, 24, 2)_{M_1}$ ,
- $(8, 11, 14, 25, 19, 1)_{M_1}$ ,  $(26, 17, 8, 19, 27, 13)_{M_1}$ ,  $(30, 18, 8, 19, 0, 27)_{M_1}$ ,
- $(15, 11, 2, 27, 29, 0)_{M_1}$ ,  $(14, 13, 0, 15, 27, 3)_{M_1}$ ,  $(10, 14, 2, 19, 18, 4)_{M_1}$ ,
- $(14, 7, 16, 30, 26, 28)_{M_1}$ ,  $(9, 18, 14, 26, 22, 0)_{M_1}$ ,  $(23, 14, 17, 31, 3, 18)_{M_1}$ ,
- $(25, 21, 0, 14, 4, 6)_{M_1}$ ,  $(0, 11, 19, 31, 1, 24)_{M_1}$ ,  $(12, 10, 0, 16, 20, 26)_{M_1}$ ,
- $(26, 2, 0, 21, 1, 15)_{M_1}$ ,  $(5, 18, 0, 12, 16, 2)_{M_1}$ ,  $(19, 15, 6, 21, 27, 2)_{M_1}$ ,
- $(16, 13, 4, 19, 26, 3)_{M_1}$ ,  $(22, 16, 15, 31, 27, 1)_{M_1}$ ,  $(23, 16, 11, 28, 4, 20)_{M_1}$ ,
- $(24, 21, 3, 16, 4, 8)_{M_1}$ ,  $(5, 23, 15, 26, 10, 30)_{M_1}$ ,  $(28, 15, 3, 18, 0, 4)_{M_1}$ ,
- $(12, 24, 15, 27, 14, 6)_{M_1}$ ,  $(4, 21, 5, 27, 15, 17)_{M_1}$ ,  $(31, 27, 10, 25, 15, 30)_{M_1}$ ,
- $(17, 18, 2, 27, 20, 10)_{M_1}$ ,  $(12, 4, 9, 31, 21, 7)_{M_1}$ ,  $(31, 7, 3, 26, 2, 9)_{M_1}$ ,
- $(6, 18, 7, 31, 9, 1)_{M_1}$ ,  $(22, 18, 13, 24, 5, 11)_{M_1}$ ,  $(13, 31, 5, 28, 20, 8)_{M_1}$ ,
- $(6, 2, 13, 28, 16, 30)_{M_1}$ ,  $(18, 1, 11, 29, 25, 16)_{M_1}$ ,  $(6, 24, 10, 26, 20, 29)_{M_1}$ ,
- $(22, 25, 7, 26, 8, 10)_{M_1}$ ,  $(26, 4, 11, 30, 29, 10)_{M_1}$ ,  $(19, 25, 12, 28, 23, 9)_{M_1}$ ,
- $(29, 22, 4, 19, 31, 8)_{M_1}$ ,  $(2, 3, 4, 29, 8, 12)_{M_1}$ ,  $(23, 25, 2, 29, 27, 6)_{M_1}$ ,
- $(21, 22, 6, 23, 28, 8)_{M_1}$ ,  $(30, 22, 2, 28, 12, 23)_{M_1}$ ,  $(12, 22, 3, 17, 13, 11)_{M_1}$ ,
- $(28, 12, 1, 29, 9, 27)_{M_1}$ ,  $(7, 1, 2, 13, 8, 15)_{M_1}$ ,  $(10, 23, 1, 13, 15, 25)_{M_1}$ ,
- $(30, 1, 3, 21, 13, 24)_{M_1}$ ,  $(11, 10, 3, 21, 20, 28)_{M_1}$ ,  $(9, 17, 5, 21, 15, 20)_{M_1}$ ,
- $(29, 7, 5, 17, 13, 21)_{M_1}$ ,  $(24, 28, 5, 17, 29, 14)_{M_1}$ ,  $(7, 10, 9, 28, 19, 4)_{M_1}$ ,
- $(24, 7, 11, 23, 31, 19)_{M_1}$ ,  $(17, 11, 6, 30, 0, 16)_{M_1}$ ,  $(25, 9, 13, 24, 30, 20)_{M_1}$ ,
- $(17, 3, 13, 25, 31, 1)_{M_1}$ ,  $(29, 9, 11, 30, 16, 27)_{M_1}$ ,

and

$(16, 8, 9, 3, 18, 25)_{M_2}$ ,  $(27, 5, 8, 6, 30, 0)_{M_2}$ ,  $(0, 7, 20, 27, 1, 24)_{M_2}$ ,  
 $(5, 3, 19, 20, 25, 1)_{M_2}$ ,  $(22, 14, 20, 5, 0, 9)_{M_2}$ ,  $(31, 20, 21, 18, 19, 24)_{M_2}$ ,  
 $(14, 8, 11, 25, 1, 4)_{M_2}$ ,  $(8, 17, 26, 19, 23, 0)_{M_2}$ ,  $(8, 18, 30, 19, 12, 2)_{M_2}$ ,  
 $(27, 11, 15, 2, 1, 22)_{M_2}$ ,  $(0, 13, 14, 15, 4, 28)_{M_2}$ ,  $(19, 10, 14, 2, 1, 8)_{M_2}$ ,  
 $(16, 7, 14, 30, 17, 0)_{M_2}$ ,  $(26, 9, 18, 14, 1, 15)_{M_2}$ ,  $(31, 14, 23, 17, 2, 9)_{M_2}$ ,  
 $(0, 21, 25, 14, 15, 29)_{M_2}$ ,  $(19, 0, 11, 31, 27, 2)_{M_2}$ ,  $(16, 10, 12, 0, 20, 1)_{M_2}$ ,  
 $(21, 2, 26, 0, 7, 12)_{M_2}$ ,  $(12, 5, 18, 0, 14, 6)_{M_2}$ ,  $(6, 15, 19, 21, 9, 1)_{M_2}$ ,  
 $(4, 13, 16, 19, 6, 25)_{M_2}$ ,  $(15, 16, 22, 31, 17, 4)_{M_2}$ ,  $(28, 16, 23, 11, 14, 26)_{M_2}$ ,  
 $(16, 21, 24, 3, 27, 29)_{M_2}$ ,  $(26, 5, 23, 15, 12, 20)_{M_2}$ ,  $(3, 15, 28, 18, 14, 27)_{M_2}$ ,  
 $(27, 12, 24, 15, 6, 23)_{M_2}$ ,  $(5, 4, 21, 27, 10, 30)_{M_2}$ ,  $(10, 27, 31, 25, 18, 4)_{M_2}$ ,  
 $(2, 17, 18, 27, 20, 24)_{M_2}$ ,  $(31, 4, 12, 9, 8, 29)_{M_2}$ ,  $(3, 7, 31, 26, 18, 23)_{M_2}$ ,  
 $(7, 6, 18, 31, 19, 4)_{M_2}$ ,  $(24, 18, 22, 13, 4, 8)_{M_2}$ ,  $(28, 13, 31, 5, 8, 21)_{M_2}$ ,  
 $(13, 2, 6, 28, 20, 8)_{M_2}$ ,  $(29, 1, 18, 11, 6, 20)_{M_2}$ ,  $(10, 6, 24, 26, 22, 8)_{M_2}$ ,  
 $(7, 22, 25, 26, 8, 15)_{M_2}$ ,  $(30, 4, 26, 11, 6, 16)_{M_2}$ ,  $(28, 19, 25, 12, 9, 27)_{M_2}$ ,  
 $(4, 22, 29, 19, 20, 23)_{M_2}$ ,  $(29, 2, 3, 4, 10, 26)_{M_2}$ ,  $(29, 23, 25, 2, 13, 21)_{M_2}$ ,  
 $(23, 21, 22, 6, 9, 19)_{M_2}$ ,  $(28, 22, 30, 2, 11, 20)_{M_2}$ ,  $(17, 12, 22, 3, 20, 10)_{M_2}$ ,  
 $(29, 12, 28, 1, 14, 24)_{M_2}$ ,  $(13, 1, 7, 2, 11, 12)_{M_2}$ ,  $(13, 10, 23, 1, 24, 30)_{M_2}$ ,  
 $(3, 1, 30, 21, 6, 0)_{M_2}$ ,  $(3, 10, 11, 21, 16, 26)_{M_2}$ ,  $(5, 9, 17, 21, 16, 2)_{M_2}$ ,  
 $(17, 7, 29, 5, 31, 1)_{M_2}$ ,  $(5, 24, 28, 17, 11, 22)_{M_2}$ ,  $(9, 7, 10, 28, 15, 20)_{M_2}$ ,  
 $(23, 7, 24, 11, 30, 12)_{M_2}$ ,  $(30, 11, 17, 6, 31, 15)_{M_2}$ ,  $(13, 9, 25, 24, 26, 27)_{M_2}$ ,  
 $(25, 3, 17, 13, 10, 15)_{M_2}$ ,  $(30, 9, 29, 11, 20, 25)_{M_2}$ .

## References

- [1] P. Adams, D.E. Bryant and M. Buchanan, A survey on the existence of  $G$ -designs, *J. Combin. Des.* **16** (2008), 373–410.
- [2] D.E. Bryant and T.A. McCourt, Existence results for  $G$ -designs, <http://wiki.smp.uq.edu.au/G-designs/>.
- [3] C.J. Colbourn and A. Rosa, *Triple Systems*, Clarendon Press, Oxford, 1999.
- [4] A.D. Forbes and T.S. Griggs, Designs for graphs with six vertices and nine edges, *Australas. J. Combin.* **70** (2018), 52–74.
- [5] Qing-de Kang, Lan-dang Yuan and Shu-xia Liu, Graph Designs for all Graphs with Six Vertices and Eight Edges, *Acta Mathematicae Applicatae Sinica, English Series* **21** (2005), 469–484.
- [6] D.R. Stinson, Hill-climbing algorithms for the construction of combinatorial designs, *Ann. Discrete Math.* **26** (1985), 321–334.

(Received 20 July 2017)