

Open Research Online

The Open University's repository of research publications and other research outputs

Climate warming will not decrease winter mortality

Journal Item

How to cite:

Staddon, Philip L.; Montgomery, Hugh E. and Depledge, Michael H. (2014). Climate warming will not decrease winter mortality. *Nature Climate Change*, 4(3) pp. 190–194.

For guidance on citations see [FAQs](#).

© 2014 Macmillan Publishers Limited.



<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Version: Accepted Manuscript

Link(s) to article on publisher's website:
<http://dx.doi.org/doi:10.1038/nclimate2121>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

1 **Climate warming will not decrease winter mortality**

2

3 Philip L Staddon^{1*}, Hugh E Montgomery², Michael H Depledge³

4

5 ¹European Centre for Environment & Human Health, University of Exeter Medical School,
6 Knowledge Spa, Royal Cornwall Hospital, Truro, Cornwall, TR1 3HD, UK

7 ²Institute for Human Health and Performance and NIHR University College London
8 Hospitals Biomedical Research Centre, University College London, Charterhouse Building,
9 Archway Campus, Highgate Hill, London, UK

10 ³European Centre for Environment & Human Health, University of Exeter Medical School,
11 St Luke's Campus, Magdalen Road, Exeter, UK

12

13

14 ***Corresponding author:**

15 Dr Philip L Staddon

16 Tel. +44 (0) 1872 258145

17 E-mail: P.L.Staddon@exeter.ac.uk

18

1 **It is widely assumed by policymakers and health professionals that the harmful health**
2 **impacts of anthropogenic climate change¹⁻³ will be partially offset by a decline in excess**
3 **winter deaths (EWDs) in temperate countries, as winters warm.⁴⁻⁶ Recent UK**
4 **government reports state that winter warming will decrease EWDs.⁷⁻⁸ Over the past few**
5 **decades however, the UK and other temperate countries have simultaneously**
6 **experienced better housing, improved health care, higher incomes, and greater**
7 **awareness of the risks of cold. The link between winter temperatures and EWDs may**
8 **therefore no longer be as strong as previously. Here we report on the key drivers which**
9 **underlie year-to-year variations in EWDs. We found that the association of year-to-year**
10 **variation in EWDs with the number of cold days in winter (< 5°C), evident until the mid**
11 **1970s, has disappeared, leaving only the incidence of influenza-like illnesses to explain**
12 **any of the year-to-year variation in EWDs in the last decade. Whilst excess winter**
13 **deaths evidently do exist, winter cold severity no longer predicts the numbers affected.**
14 **We conclude that no evidence exists that EWDs in England and Wales will fall if**
15 **winters warm with climate change. These findings have important implications for**
16 **climate change health adaptation policies.**

17 Seasonal variation in death rates in temperate countries has long been recognised.
18 Excess winter deaths (EWDs) in the UK are defined as the number of deaths from December
19 to March minus the average number of deaths in the preceding August to November, and the
20 following April to July.⁹ Despite fewer cases in northern than southern Europe,¹⁰ EWDs are
21 causally attributed to seasonal variations in temperature, with low temperatures thought to
22 cause death directly (e.g. through hypothermia or falls in icy conditions), and by altering
23 vulnerability to communicable or non-communicable diseases, such as influenza and
24 myocardial infarction, which are more common in winter.¹¹ We collated data from the past 60
25 years to identify key factors associated with the decreasing trend in EWDs in England and

1 Wales, and its year-to-year variation. We deliberately considered a very broad set of factors
2 to minimise the risk of erroneous conclusions. To clarify the thrust of this paper, we are
3 interested in explaining the year to year variation in EWDs not the daily variation, and we are
4 not saying that temperature does not play a role, if it didn't there would be no EWDs. What
5 we aim to demonstrate is that how harsh a winter is no longer predicts how many EWDs
6 there will be.

7 Figure 1 presents relative excess winter deaths (EWDs) and variables identified as
8 possible mediating or causal factors. Between 1951 and 2011, both absolute and year-to-year
9 variation in EWDs declined over time. Three distinct periods in EWDs changes were
10 apparent (Supplementary Fig. S1): (1) 1951-1970, where EWDs exhibited very high year-to-
11 year variation, and a strongly decreasing overall trend; (2) 1971-2000, where year-to-year
12 variation EWDs halved compared to the preceding period, and the decreasing trend continues,
13 albeit less strongly; and (3) 2001-2011, where year-to-year variation is very small and the
14 EWDs rate is flat.

15 Multi-factorial regressions were carried out on the whole dataset and by selected
16 periods (Table 1). Over the entire period, housing quality, heating costs, number of cold days
17 and influenza (flu) activity were highly significant in explaining the level of EWDs, and
18 together account for ca. 77% of its variation. However, factors associated with EWDs differ
19 over time when analysed in 20 year segments, and were: (1) for 1951-1971, housing quality
20 and number of cold days; (2) for 1971-1991, the number of cold days and flu activity; and (3)
21 for 1991-2011, flu activity. For the data split in 1976, which is the approximate time of
22 correlation breakdown between EWDs and number of cold days (see below), the following
23 factors were significant in explaining EWDs variation: (1) for 1951-1976, housing quality,
24 the number of cold days, and flu activity; and (2) for 1976-2011, flu activity. Other splits

1 show a similar pattern, with impacts of housing and cold days disappearing, leaving only the
2 impact of influenza to explain year-to-year variation in EWDs.

3 Multi-factorial regression analysis of smoothed data for the whole period, and for the
4 periods before and after 1986, add further insight (Table 1). Note that as the smoothed data
5 are for 10 year rolling periods, 1986 (i.e. 10 year average to 1986) was chosen rather than
6 1976 (as above), so that the second period only contained data after the correlation
7 breakdown between EWDs and number of cold days. This division resulted in two periods of
8 identical length. For the whole period, housing quality, heating costs, and policy initiatives
9 explain around 92% of the time trend variation. As for the previous analysis on split data,
10 there is a difference in the factors which are significant, depending on the period. Up to 1986,
11 housing quality, number of cold days, and number of cold days with a large drop in
12 temperature, were all significant; after 1986, it was housing quality, flu activity and policy
13 initiatives (marginally significant) that explain most of the variation over time. The difference
14 between these two periods is striking (Supplementary Fig. S2).

15 The analysis of the de-trended data and the causes of year-to-year variation in EWDs
16 are shown in Table 1. Over the whole period, the number of cold days and flu activity were
17 highly significant, explaining ca. 43% of the variation. Before 1976, most of the variation is
18 explained by the number of cold days, but with a proportion explained by flu activity. After
19 1976, only flu activity accounts for any of the year-to-year variation in EWDs. Figure 2
20 presents the de-trended data for EWDs in relation to (A) the number of cold days, and (B)
21 seasonal flu activity.

22 By performing rolling correlations between EWDs and factors exhibiting year-to-year
23 variation, namely the number of cold days, the number of cold days with a large temperature
24 drop and the magnitude of flu activity, it emerged that the correlation between EWDs and the
25 independent variable, when significant, was not stable over time. This is illustrated in Fig. 3

1 for the number of cold days (A) and flu activity (B). EWDs remained strongly correlated with
2 the number of cold days only up to the mid- to late-1970s, after which the correlation was
3 weak to non-existent. Similarly, for most of the period from 1951 to the mid 1990s EWDs
4 were correlated weakly with flu activity, while a strong correlation between the two has
5 emerged in recent years. For winter temperature volatility, the correlation with EWDs was
6 rarely strong and there was no established stable period (data not shown).

7 An extensive literature attests to the fact that changes in daily temperature influence
8 health outcomes at the local levels, and that EWDs are influenced by temperature. However,
9 our data suggest that year to year variation in EWDs is no longer explained by the year to
10 year variation in winter temperature: winter temperatures now contribute little to the yearly
11 variation in excess winter mortality so that milder winters resulting from climate change are
12 unlikely to offer a winter health dividend. Our research paints a clear picture of why EWDs
13 have been decreasing in the UK over the past six decades, and which factors explain most of
14 the year-to-year variation in EWDs. These include better housing, better standards of living,
15 increased help for vulnerable sections of the population, as well as better healthcare. This
16 confirms proposed mechanisms, presented in several recent studies in temperate countries,¹²⁻
17 ¹⁴ that explain the general decreasing trend in EWDs over the past century. Whilst the key
18 driver for year-to-year variation in EWDs was winter temperature until the mid 1970s, we
19 show that it has been superseded by the impact of influenza-like illnesses despite their
20 absolute impact being small; this is consistent with recent studies by the UK Office for
21 National Statistics.^{15,16} This time-dependency explains why there is so much confusion about
22 the link between winter temperatures *per se* and EWDs. Many of the papers that concluded
23 that climate change would lead to fewer EWDs, are not recent, and rely on relatively old data
24 (from *ca.* two decades ago).^{12,17-19} Based on the evidence available at the time, these early
25 papers¹⁷⁻¹⁹ predicting that climate change would cause a decrease in EWDs were correct.

1 However, more recent papers are either inconclusive,²⁰ or conclude that there will be little
2 impact of climate change on EWDs.²¹⁻²³ By analysing more recent data, and performing
3 rolling correlation analysis on time detrended data, we show unequivocally that the
4 correlation between the number of cold winter days per year and EWDs, which was strong
5 until the mid 1970s, no longer exists.

6 We used a ‘threshold model’ to identify a strong relationship between annual number
7 of cold days and EWDs before the mid-1970s, and to show that this relationship has since
8 disappeared. The relationship between mortality and local daily temperature is variable and
9 specific to local areas, and it is likely that the exposure-response relationship of daily
10 mortality to temperature will have changed over the past decades in response to improved
11 housing, health and wealth. A future avenue of research would be to explore our observations
12 further, by the use of exposure-response analysis of continuous and locally-applied data.

13 Our results should also be considered in light of the latest findings relating to the
14 influence of climate change on winter temperatures. The view that winter temperatures would
15 simply ‘ramp up’ has been replaced by recognition that extreme events, including cold spells
16 and storms, are likely to increase in frequency.^{5,24} Indeed, there is already evidence that
17 winter temperature volatility has increased in the UK over the past 20 years. For example, the
18 number of days per winter with a mean daily temperature $<5^{\circ}\text{C}$ and showing a 4°C drop from
19 the previous day (i.e. high variability), exhibits an increasing trend from 1990 to 2011 (R-
20 square 0.31; p 0.007). If this is exacerbated by climate change in the coming decades, then
21 winters will feel fundamentally different from now, being generally warmer, but with more
22 days of severe cold. The nefarious effects on EWDs could be substantial, with especially the
23 vulnerable being caught off guard by abrupt changes in temperature. This behaviour-
24 mediated impact of temperature variability on EWDs may be one of the reasons countries
25 with milder winters often exhibit higher levels of EWDs than countries with colder winters.¹¹

1 It is also possible that increased temperature volatility could increase influenza deaths,
2 although this is yet to be proved conclusively.

3 The fact that climate change will not reduce EWDs in the England and Wales, has
4 important implications for health policy. Probable increases in future winter temperature
5 volatility²⁵ mean that EWDs are more likely to rise than fall. Added to this, and irrespective
6 of whether climate change induced winter temperature volatility increases the risk of EWDs,
7 the absolute number of EWDs may increase in the coming decades simply because of a
8 growing and ageing population. The recent policy focus on protecting the public from
9 heatwaves should not be at the expense of preventing the much more numerous EWDs.
10 Energy efficiency regulations and government retrofitting initiatives to improve the thermal
11 efficiency of older homes, including double glazing, cavity wall and loft insulation, should
12 continue to capture co-benefits for both health and climate change mitigation.^{14,26} In view of
13 our findings, particular attention should also be paid to public health initiatives to reduce the
14 risk of infection with flu-like illnesses. Influenza vaccination, despite its decreasing
15 effectiveness in people over 70s, provides some protection.²⁷ Improving uptake in the over
16 65s would be very beneficial.²⁸

17 From a health perspective, managing risk uncertainty is a priority²⁹, and prevention is
18 better than cure. Urgently reducing greenhouse gas emissions to mitigate against climate and
19 weather change is therefore essential.⁵ This goes hand in hand with the need for a sound
20 strategy for health adaptation for an ageing population in a changing climate.

21

1 **Methods**

2 **Data sources and quality**

3 An initial search of the Web of Knowledge was performed to identify factors influencing
4 excess winter death rates. The search encompassed all years and excluded non-English
5 language articles. Multiple combinations of search terms were explored (Supplementary
6 Information Search Terms). Secondary searches were performed on references cited by
7 articles discussing EWDs and their causes.

8 Articles relating to the causes of EWDs were identified as targets for data retrieval.
9 These were supplemented by source data: temperature data were obtained from the UK's
10 Meteorological Office - Hadley Centre Central England Temperature dataset;³⁰ and economic,
11 social, population and mortality data from the Office for National Statistics (ONS). When the
12 required data were unavailable, such as those relating to housing quality or to government
13 initiatives to combat EWDs, a web search using Google was initiated, which was focussed on
14 information held in government departments, agencies and other organisations holding
15 specialist housing datasets (Supplementary Information Search Terms).

16 Data were collected for the period 1951 to 2011 and a full list of all data sources used
17 are provided in Supplementary Information (Table S1). Population (and a subset aged over 65
18 years), excess winter mortality and incidence of influenza-like illness in England and Wales
19 were documented. Daily mean temperatures for Central England were also obtained. These
20 data, being representative of a national geographical mean,³⁰ were sufficient for our study of
21 national trends. We therefore collected demographic, EWD, influenza incidence, and winter
22 temperature data representative of England and Wales. Certainly different regions will
23 experience different temperatures, but they are all highly correlated to this Central England
24 value. Also, EWDs are surprisingly stable across regions, with for example the EWDs in
25 Cornwall, the mildest part of England, being near identical that for England and Wales.

1 A range of extrinsic factors influencing seasonal, temperature-related mortality were
2 also documented. Specific factors were only excluded when their impacts or characteristics
3 were already represented within another factor (e.g. income level versus percentage
4 household expenditure on fuel). We recorded expenditure on heating as a percentage of
5 income, policy initiatives aimed at combating excess winter deaths (cold weather payment,
6 winter fuel allowance, ‘warm front’), and four key housing quality factors, each focussed on
7 a particular housing characteristic affecting health in winter (availability of inside toilet,
8 access to tapped hot water, central heating, and double glazing). Data was drawn from the
9 Domestic Energy fact file, the UK Housing Energy fact file, the English Housing Survey, and
10 the Halifax housing dataset, which contains a reliable description of the changing condition
11 of the UK housing stock. Housing quality improvements were assumed to be linear between
12 available years. This assumption was validated for double glazing, where data for the full
13 period were available. For statistical analyses, all four housing quality measures were
14 combined by simple averaging into a single measure.

15

16 **Statistical analyses**

17 Excess winter deaths (EWDs) were expressed as a function of population over 65 years (as *ca.*
18 90% of total EWDs occur in this age group⁹) to remove changing demographics as a factor.
19 The raw daily temperature data were transformed into two measures: number of days per
20 winter period below 5°C (a measure of winter cold intensity); and number of days per winter
21 below 5°C and showing a 4°C drop from the previous day (a measure of volatility within cold
22 spells, defined as one or more days below 5°C). Cold days are calculated for the same period
23 that EWDs are, namely 01 December to 31 March. Correlation analysis was used to
24 determine the interdependence of variables. Linear multi-factor regression analysis identified
25 those factors associated independently with excess winter deaths. This was achieved by

1 performing a series of regressions removing the least significant factor at each repeat until
2 only highly significant factors remained. We also ensured that the amount of variation
3 explained by the fitted model, (R-square), remained relatively stable. To explain the trend
4 over time in more detail, a moving average method was employed with a period of 10 years.
5 This allowed the smoothing of the data to eliminate most of the year-to-year variation. Linear
6 multi-factor regression analysis was applied to the smoothed data. To assess year-to-year
7 variation, the relevant data were de-trended by removal of the time component and analysed
8 by linear multi-factor regression. Using separate data sets, i.e. a smoothed data set and a
9 detrended data set, allows the elimination of confounding time dependent factors when
10 addressing the two specific questions of (1) what is causing the *long-term* trend in decreasing
11 EWDs, and (2) what is causing the *short term* year to year variation in EWDs. We also tested
12 for correlation breakdown between excess winter deaths and the factors with strong year-to-
13 year variation. To further elucidate factors associated with EWDs, data were split into subsets
14 characterised by key changes in factors, or the introduction of new factors (such as a policy
15 initiative).

16

17

1 **References**

- 2 1. Costello, A., Abbas, M., Allen, A., *et al.* Managing the health effects of climate
3 change: Lancet and University College London Institute for Global Health
4 Commission. *Lancet* 373, 1693-1733 (2009).
- 5 2. Patz, J.A., Campbell-Lendrum, D., Holloway, T. & Foley, J.A. Impact of regional
6 climate change on human health. *Nature* 438, 310-317 (2005).
- 7 3. Altizer, S., Ostfeld, R.S., Johnson, P.T.J., Kutz, S. & Harvell, C.D. Climate change
8 and infectious diseases: from evidence to a predictive framework. *Science* 341, 514-
9 519 (2013).
- 10 4. Stern, N. *The Economics of Climate Change. The Stern Review* (Cambridge
11 University Press, 2007).
- 12 5. McMichael, T., Montgomery, H. & Costello, A. Health risks, present and future, from
13 global climate change. *BMJ* 344, e1359 (2012).
- 14 6. Li, T., Horton, R.M. & Kinney, P.L. Projections of seasonal patterns in temperature-
15 related deaths for Manhattan, New York. *Nature Clim. Change* 3, 717-721.
- 16 7. HPA. *Health Effects of Climate Change in the UK 2012* (Health Protection Agency,
17 2012). <http://www.hpa.org.uk/hecc2012> (accessed 05/02/13).
- 18 8. CCRA. *The UK Climate Change Risk Assessment 2012*. (Department for Environment,
19 Food and Rural Affairs, 2012). [www.defra.gov.uk/environment/climate/government/
20 risk-assessment/#report](http://www.defra.gov.uk/environment/climate/government/risk-assessment/#report) (accessed 05/02/13).
- 21 9. ONS. *Excess winter mortality in England and Wales, 1950/51 to 2009/10* (Office for
22 National Statistics, 2010).

- 1 10. Analitis, A., Katsouyanni, K., Biggeri, A., *et al.* Effects of cold weather on mortality:
2 results form 15 European cities within PHEWE project. *Am. J. Epidemiol.* 168, 1397-
3 1408 (2008).
- 4 11. Keatinge, W.R., Donaldson, G.C., Bucher, K., *et al.* Cold exposure and winter
5 mortality from ischaemic heart disease, cerebrovascular disease, respiratory disease,
6 and all causes in warm and cold regions of Europe. *Lancet* 349, 1341-1346 (1997).
- 7 12. Carson, C., Hajat, S., Armstrong, B. & Wilkinson, P. Declining vulnerability to
8 temperature-related mortality in London over the 20th Century. *Am. J. Epidemiol.* 164,
9 77-84 (2006).
- 10 13. Healy, J.D. Excess winter mortality in Europe: a cross country analysis identifying
11 key risk factors. *J. Epidemiol. Comm. Health* 57, 784-789 (2003).
- 12 14. Howden-Chapman, P. & Chapman, R. Health co-benefits from housing-related
13 policies. *Curr. Opinion Environ. Sustain.* 4, 414-419 (2012).
- 14 15. Hicks, J. & Allen, G. A. *Century of Changes: Trends in UK Statistics since 1900.*
15 (1999). www.parliament.uk/documents/commons/lib/research/rp99/rp99-111.pdf
16 (accessed 17/01/13).
- 17 16. Brown, G., Fearn, V. & Wells, C. Exploratory analysis of seasonal mortality in
18 England and Wales, 1998-2007. *Health Statistics Quarterly* 48. (Office for National
19 Statistics, 2010).
- 20 17. Kalkstein, L.S. & Breene, J.S. An evaluation of climate/mortality relationships in
21 large U.S. cities and the possible impacts of a climate change. *Environ. Health*
22 *Perspect.* 105, 84-93 (1997).

- 1 18. Kovats, R.S., Haines, A., Stanwell-Smith, R., Martens, P., Menne, B. & Bertollini, R.
2 Climate change and human health in Europe. *BMJ* 318, 1682-1685 (1999).
- 3 19. Langford, I.H. & Bentham, G. The potential effects of climate-change on winter
4 mortality in England and Wales. *Int. J. Biometeorol.* 38, 141-147 (1995).
- 5 20. McGeehin, M.A. & Mirabelli, M. The potential impacts of climate variability and
6 change on temperature-related morbidity and mortality in the United States. *Environ.*
7 *Health Perspect.* 109, 185-189 (2001).
- 8 21. Davis, R.E., Knappenberger, P.C., Michaels, P.J. & Novicoff, W.M. Seasonality of
9 climate-human mortality relationships in US cities and impacts of climate change.
10 *Climate Res.* 26, 61-76 (2004).
- 11 22. Martin, S.L., Cakmak, S., Hebborn, C.A., Avramescu, M.L. & Tremblay, N. Climate
12 change and future temperature-related mortality in 15 Canadian cities. *Int. J.*
13 *Biometeorol.* 56, 605-619 (2012).
- 14 23. Morabito, M., Crisci, A., Moriondo, M., *et al.* Air temperature-related human health
15 outcomes: current impact and estimations of future risks in Central Italy. *Sci. Total*
16 *Environ.* 441, 28-40 (2012).
- 17 24. IPCC. *Managing the Risks of Extreme Events and Disasters to Advance Climate*
18 *Change Adaptation. A Special Report of Working Groups I and II of the*
19 *Intergovernmental Panel on Climate Change.* (Cambridge University Press, 2012.)
- 20 25. Coumou, D. & Rahmstorf, S. A decade of weather extremes. *Nature Clim. Change* 2,
21 491-496 (2012).

- 1 26. WHO. *Health in the Green Economy: Health Co-benefits of Climate Change*
2 *Mitigation - Housing Sector*. World Health Organization. (WHO Press, 2011).
- 3 27. Simonson, L., Taylor, R.J., Miller, M.A. & Jackson, L.A. Mortality benefits of
4 influenza vaccination in elderly people: an ongoing controversy. *Lancet Infect. Dis.* 7,
5 658-666 (2007).
- 6 28. Goddard, N.L., Joseph, C.A., Zambon, M., Nunn, M., Fleming, D. & Watson, J.M.
7 Influenza surveillance in England and Wales: October 1999 to May 2000. *Commun.*
8 *Dis. Public Health* 3, 261-266 (2000).
- 9 29. Kunreuther, H., Heal, G., Allen, M., Edenhofer, O., Field, C.B. & Yohe, G. Risk
10 management and climate change. *Nature Clim. Change* 3, 447-450 (2013).
- 11 30. Met Office. *Met Office Hadley Centre Central England Temperature Data*. (2012).
12 <http://www.metoffice.gov.uk/hadobs/hadcet/data/download.html> (accessed 24/10/12).
- 13

1 **Correspondence**

2 Correspondence and request for data should be addressed to PL Staddon.

3

4 **Acknowledgements**

5 We thank Charles McGilligan for comments and help identifying data sources. The European
6 Centre for Environment and Human Health is supported by the European Regional
7 Development Fund and the European Social Fund Convergence Programme for Cornwall and
8 the Isles of Scilly. The funder had no influence whatsoever on the research presented in this
9 article.

10

11 **Author contributors**

12 The idea for this work arose from a meeting between the three authors. PLS searched the
13 literature, collected the data, performed the analysis and wrote the first draft. All authors
14 contributed to the final version of the paper. PLS had full access to all the data in the study
15 and had final responsibility for the decision to publish this article.

16

17 **Competing financial interests**

18 The authors declare no competing financial interests.

19

20

Table 1| Multivariate regression analysis of the relationship between excess winter deaths and independent variables for selected periods between 1951 and 2011

		Significance level (<i>p</i>) and standardized coefficient (β)							
R-square		Total	HQ	HC	Pol	CD	TD	FA	
ON UNMODIFIED DATA									
1951-2011	0.78	<i>p</i>	0.000	0.002	0.032	0.312	0.000	0.544	0.000
		β	-0.597	-0.404	0.204	0.329	0.041	0.621	
1951-2011	0.77	<i>p</i>	0.000	0.000	0.000	X	0.000	X	0.000
		β	-0.523	-0.525	X	0.350	X	0.612	
1951-1971	0.72	<i>p</i>	0.000	0.025	NA	NA	0.002	0.461	0.178
		β	-0.377	NA	NA	0.606	-0.109	0.241	
1971-1991	0.61	<i>p</i>	0.022	0.844	0.622	0.437	0.023	0.592	0.003
		β	0.077	-0.171	-0.265	0.560	0.102	0.796	
1991-2011	0.72	<i>p</i>	0.003	0.141	0.299	0.290	0.622	0.455	0.000
		β	-0.873	-0.288	0.623	0.094	0.175	0.765	
1951-1976	0.75	<i>p</i>	0.000	0.011	NA	NA	0.001	0.654	0.011
		β	-0.323	NA	NA	0.510	-0.053	0.340	
1976-2011	0.65	<i>p</i>	0.000	0.170	0.089	0.932	0.092	0.318	0.000
		β	-0.498	-0.521	-0.031	0.221	0.125	0.681	
ON SMOOTHED ^a DATA									
1951-2011	0.92	<i>p</i>	0.000	0.000	0.016	0.040	0.534	0.633	0.712
		β	-1.542	-0.351	0.357	0.066	-0.028	-0.057	
1951-2011	0.92	<i>p</i>	0.000	0.000	0.010	0.009	X	X	X
		β	-1.517	-0.320	0.366	X	X	X	
1951-1985	0.97	<i>p</i>	0.000	0.000	NA	NA	0.002	0.010	0.241
		β	-0.952	NA	NA	0.154	0.130	-0.068	
1951-1985	0.97	<i>p</i>	0.000	0.000	NA	NA	0.000	0.000	X
		β	-0.907	NA	NA	0.176	0.161	X	
1986-2011	0.88	<i>p</i>	0.000	0.012	0.409	0.058	0.147	0.101	0.058
		β	-1.187	-0.196	0.677	0.412	-0.239	0.376	
1986-2011	0.85	<i>p</i>	0.000	0.000	X	0.057	X	X	0.007
		β	-1.202	X	0.695	X	X	0.489	
ON DETRENDED ^b DATA									
1951-2011	0.43	<i>p</i>	0.000	NA	NA	NA	0.000	0.982	0.000
		β	NA	NA	NA	0.411	-0.002	0.470	
1951-1976	0.62	<i>p</i>	0.000	NA	NA	NA	0.000	0.636	0.025
		β	NA	NA	NA	4.383	-0.480	2.404	
1976-2011	0.40	<i>p</i>	0.000	NA	NA	NA	0.207	0.938	0.000
		β	NA	NA	NA	0.181	-0.011	0.627	

HQ housing quality; HC heating costs; Pol policy initiatives; CD number of cold days; TD number of cold days with strong temperature drop; FA flu activity; NA not applicable; X removed; ^a analysis performed on smoothed data (10 year moving average) - to identify the variables behind the decreasing trend; ^b analysis performed on detrended data (time component removed) - to identify the variables behind the year-to-year variation.

1 **FIGURE LEGENDS**

2

3 **Figure 1| Relative excess winter mortality for England and Wales over the past 60 years**
4 **presented alongside key determinants**

5 An index is used to allow for easy comparison in trends and year-to-year variation. Policy
6 initiatives are cold weather payments (CWP), winter fuel payments (WFP), and warm front
7 (WF). Excess winter deaths are expressed relative to the size of the population over 65 years
8 old. Before indexation, activity of influenza like illness was categorised on a scale of 0 to 4,
9 with '0' as baseline, and '4' the level of the 1951 epidemic. Housing quality is based on four
10 parameters: inside toilet, hot water, central heating, and double glazing. Heating cost is
11 measured as relative to household expenditure.

12

13 **Figure 2| Detrended data showing the year-to-year variation in relative excess winter**
14 **mortality compared to the number of cold days and to the activity level of influenza like**
15 **illness**

16 Data was detrended by removing the time component. The year-to-year variation in excess
17 winter deaths is compared with that for (A) number of winter days $< 5^{\circ}\text{C}$, and (B) influenza
18 activity. An index is used to allow for easy comparison of comparison of peaks. Excess
19 winter deaths are expressed relative to the size of the population over 65 years old. Before
20 indexation, activity of influenza like illness was categorised on a scale of 0 to 4, with '0' as
21 baseline, and '4' the level of the 1951 epidemic.

22

23 **Figure 3| Rolling 10 year correlation between relative excess winter mortality and the**
24 **two main predictors of year-to-year variation: number of winter cold days and activity**
25 **of influenza like illness**

1 Correlation coefficients are from 10 year rolling correlations. Correlation above 0.50 is
2 deemed strong. (A) The relationship between excess winter deaths and number of cold days
3 exhibits a classic correlation breakdown in the late 1970s. (B) The relationship between
4 excess winter deaths and influenza activity stabilises and strengthens from the mid 1990s.