

The Open University's repository of research publications and other research outputs

Draft Genome Sequences of the Nitrate-Dependent Iron-Oxidizing Proteobacteria *Acidovorax* sp. Strain BoFeN1 and *Paracoccus pantotrophus* Strain KS1

Journal Item

How to cite:

Price, A.; Macey, M. C.; Miot, J. and Olsson-Francis, K. (2018). Draft Genome Sequences of the Nitrate-Dependent Iron-Oxidizing Proteobacteria *Acidovorax* sp. Strain BoFeN1 and *Paracoccus pantotrophus* Strain KS1. *Microbiology Resource Announcements*, 7(10), article no. e01050-18.

For guidance on citations see [FAQs](#).


© 2018 The Authors

Version: Version of Record

Link(s) to article on publisher's website:

<http://dx.doi.org/doi:10.1128/MRA.01050-18>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.



Draft Genome Sequences of the Nitrate-Dependent Iron-Oxidizing Proteobacteria *Acidovorax* sp. Strain BoFeN1 and *Paracoccus pantotrophus* Strain KS1

A. Price,^a M. C. Macey,^a J. Miot,^b K. Olsson-Francis^a

^aFaculty of Science, Technology, Engineering and Mathematics, The Open University, Milton Keynes, United Kingdom

^bInstitut de Minéralogie, de Physique des Matériaux et de Cosmochimie, Muséum National d'Histoire Naturelle, Université Pierre et Marie Curie–Sorbonne Universités, CNRS UMR 7590, IRD 206, Paris, France

ABSTRACT The draft genomes of the nitrate-dependent iron-oxidizing bacteria *Acidovorax* sp. strain BoFeN1 and *Paracoccus pantotrophus* strain KS1 are presented. These genomes supply supporting data to investigations of the mechanisms underlying this anaerobic form of microbial biogeochemical iron cycling.

Acidovorax sp. strain BoFeN1 and *Paracoccus pantotrophus* strain KS1 have both demonstrated the capability to perform nitrate-dependent Fe²⁺ oxidation (NDFO), whereby Fe³⁺ precipitates are formed during microbial nitrate reduction (1, 2). NDFO is a geomicrobiological process which has been recognized for the past 2 decades (3) and is responsible for the production of Fe³⁺-bearing minerals in anaerobic, circum-neutral, reducing environments under conditions similar to those that were prevalent on the early Earth and ancient Mars (1, 4–6). Despite this potential importance in the early history of life, the putative underlying mechanisms are largely unconfirmed experimentally (7), despite some advances (8, 9).

Acidovorax sp. strain BoFeN1 is a Gram-negative, rod-shaped bacterium belonging to the family *Comamonadaceae* in the order *Burkholderiales* of the class *Betaproteobacteria*. *Acidovorax* sp. strain BoFeN1 was isolated from littoral sediments of Lake Constance on the Germany-Switzerland border and assigned to the genus *Acidovorax* based on 16S rRNA gene sequence analysis (1). This isolate was donated by Jennyfer Miot of the Mineralogy, Material Physics and Cosmochemistry Institute (IMPMC) in Paris, France.

Paracoccus pantotrophus strain KS1 is a Gram-negative, motile coccoid bacterium belonging to the family *Rhodobacteraceae* in the order *Rhodobacterales*, in the class *Alphaproteobacteria*. *Paracoccus pantotrophus* strain KS1 was isolated from soil at Kew Gardens in the United Kingdom and was originally reported as a member of the genus *Thiobacillus* but was later reclassified to the genus *Paracoccus* based on the 16S rRNA gene sequence (10, 11). This strain was purchased as a live culture (DSM 11072) from the German Collection of Microorganisms and Cell Cultures (DSMZ) in Leibniz, Germany.

The two strains were maintained in pure culture in aerobic nutrient medium (5.0 g liter⁻¹ peptone and 3.0 g liter⁻¹ meat extract) at 30°C prior to DNA extraction and genome sequencing, which was performed by MicrobesNG (<https://microbesng.uk/>) using the Illumina MiSeq platform with 2 × 250-bp paired-end reads. Trimmed reads were produced using Trimmomatic version 0.30 (<http://www.usadellab.org/cms/?page=trimmomatic>) with a sliding window quality cutoff of Q15, and *de novo* assembly was performed using SPAdes version 3.7 (<http://cab.spbu.ru/software/spades/>) (12, 13). Coverage of 30× was achieved during sequencing, and genome annotation was

Received 30 July 2018 Accepted 15 August 2018 Published 13 September 2018

Citation Price A, Macey MC, Miot J, Olsson-Francis K. 2018. Draft genome sequences of the nitrate-dependent iron-oxidizing proteobacteria *Acidovorax* sp. strain BoFeN1 and *Paracoccus pantotrophus* strain KS1. *Microbiol Resour Announc* 7:e01050-18. <https://doi.org/10.1128/MRA.01050-18>.

Editor J. Cameron Thrash, Louisiana State University

Copyright © 2018 Price et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to A. Price, alex.price@open.ac.uk.

performed using the RAST annotation server version 2.0 (<http://rast.nmpdr.org/rast.cgi>) via the classic RAST pipeline (14). The draft genome of strain *Acidovorax* sp. strain BoFeN1 consisted of 4.06 Mb in 184 contigs ($N_{50} = 37,384$), with a GC content of 63.77%. A total of 3,794 protein-coding sequences were predicted from the annotated genome, with 50 tRNAs identified using the ARAGORN version 1.1 tRNA detection program (<http://mbio-serv2.mbioekol.lu.se/ARAGORN/>) (15). The closest match for the 16S rRNA gene was identified in the GenBank database, by using the BLASTN program (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>), as *Acidovorax defluvii* strain BSB411 (99% similarity; GenBank accession number [NR_026506](https://pubmed.ncbi.nlm.nih.gov/2026506/)). The draft genome of *Paracoccus pantotrophus* strain KS1 consisted of 4.16 Mb in 227 contigs ($N_{50} = 36,889$), with a GC content of 67.69%. A total of 3,941 protein-coding sequences and 54 tRNAs were predicted from the annotated genome. The nearest 16S rRNA gene sequence was identified using BLASTN, with a best match as *Paracoccus pantotrophus* (100%; accession number [AB098590](https://pubmed.ncbi.nlm.nih.gov/1098590/)).

Neither genome contained a dedicated ferroxidase, suggesting that NDFO in these strains is not a direct enzymatic process to acquire electrons. Full denitrification pathways were confirmed in both strains by the presence of nitrogen species reductase genes (*nar*, *nir*, *nor*, and *nos*), with additional genes for a periplasmic nitrate reductase (Nap) in the *Paracoccus pantotrophus* strain KS1 genome.

Data availability. DDBJ/ENA/GenBank accession numbers have been assigned for *Acidovorax* sp. strain BoFeN1 ([QOZT00000000](https://pubmed.ncbi.nlm.nih.gov/30000000/)) and *Paracoccus pantotrophus* strain KS1 ([QOZU00000000](https://pubmed.ncbi.nlm.nih.gov/30000000/)). Raw sequencing reads for *Acidovorax* sp. strain BoFeN1 ([SRP157586](https://pubmed.ncbi.nlm.nih.gov/157586/)) and *Paracoccus pantotrophus* strain KS1 ([SRP157588](https://pubmed.ncbi.nlm.nih.gov/157588/)) are available in the NCBI Sequence Read Archive.

ACKNOWLEDGMENTS

Genome sequencing was provided by MicrobesNG (<http://www.microbesng.uk>), which is supported by the BBSRC (grant number BB/L024209/1).

This work was supported by STFC training grant ST/N50421X/1.

REFERENCES

- Kappler A, Schink B, Newman DK. 2005. Fe(III) mineral formation and cell encrustation by the nitrate-dependent Fe(II)-oxidizer strain BoFeN1. *Geobiology* 3:235–245. <https://doi.org/10.1111/j.1472-4669.2006.00056.x>.
- Kumaraswamy R, Sjollem K, Kuenen G, Van Loosdrecht M, Muyzer G. 2006. Nitrate-dependent [Fe(II)EDTA]²⁻ oxidation by *Paracoccus ferrooxidans* sp. nov., isolated from a denitrifying bioreactor. *Syst Appl Microbiol* 29:276–286. <https://doi.org/10.1016/j.syapm.2005.08.001>.
- Straub KL, Benz M, Schink B, Widdel F. 1996. Anaerobic, nitrate-dependent microbial oxidation of ferrous iron. *Appl Environ Microbiol* 62:1458–1460.
- Miot J, Etique M. 2016. Formation and transformation of iron-bearing minerals by iron (II)-oxidizing and iron (III)-reducing bacteria, p. 53–98. *In* Faivre D (ed), *Iron oxides: from nature to applications*. Wiley-VCH, Weinheim, Germany.
- Ilbert M, Bonnefoy V. 2013. Insight into the evolution of the iron oxidation pathways. *Biochim Biophys Acta* 1827:161–175. <https://doi.org/10.1016/j.bbabi.2012.10.001>.
- Price A, Pearson VK, Schwenzler SP, Miot J, Olsson-Francis K. 2018. Nitrate-dependent iron oxidation: a potential mars metabolism. *Front Microbiol* 9:513. <https://doi.org/10.3389/fmicb.2018.00513>.
- Carlson H, Clark I, Melnyk R, Coates J. 2012. Toward a mechanistic understanding of anaerobic nitrate-dependent iron oxidation: balancing electron uptake and detoxification. *Front Microbiol* 3:57. <https://doi.org/10.3389/fmicb.2012.00057>.
- Carlson HK, Clark IC, Blazewicz SJ, Iavarone AT, Coates JD. 2013. Fe(II) oxidation is an innate capability of nitrate-reducing bacteria that involves abiotic and biotic reactions. *J Bacteriol* 195:3260–3268. <https://doi.org/10.1128/JB.00058-13>.
- Miot J, Remusat L, Duprat E, Gonzalez A, Pont S, Poinot M. 2015. Fe biomineralization mirrors individual metabolic activity in a nitrate-dependent Fe(II)-oxidizer. *Front Microbiol* 6:879. <https://doi.org/10.3389/fmicb.2015.00879>.
- Jordan SL, Kraczkiewicz-Dowjat AJ, Kelly DP, Wood AP. 1995. Novel eubacteria able to grow on carbon disulfide. *Arch Microbiol* 163:131–137.
- Jordan SL, McDonald IR, Kraczkiewicz-Dowjat AJ, Kelly DP, Rainey FA, Murrell JC, Wood AP. 1997. Autotrophic growth on carbon disulfide is a property of novel strains of *Paracoccus denitrificans*. *Arch Microbiol* 168:225–236. <https://doi.org/10.1007/s002030050492>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
- Laslett D, Canback B. 2004. ARAGORN, a program to detect tRNA genes and tmRNA genes in nucleotide sequences. *Nucleic Acids Res* 32:11–16. <https://doi.org/10.1093/nar/gkh152>.