

Open Research Online

The Open University's repository of research publications and other research outputs

Metal Impactor Fragment found in Lunar Regolith Breccia Meteorite North West Africa 10989

Conference or Workshop Item

How to cite:

Morland, Zoe; Joy, Katherine; Gholinia, A and Degli Alessandrini, Giulia (2019). Metal Impactor Fragment found in Lunar Regolith Breccia Meteorite North West Africa 10989. In: 7th European Lunar Symposium, 21-23 May 2019, Manchester, UK.

For guidance on citations see [FAQs](#).

© [not recorded]



<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Version: Version of Record

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

METAL IMPACTOR FRAGMENT FOUND IN LUNAR REGOLITH BRECCIA METEORITE NORTH WEST AFRICA 10989. Z. S. Morland^{1,2}, K. H. Joy², A. Gholinia³ and G. Degli Alessandrini¹. ¹School of Physical Sciences, Open University, Milton Keynes, UK., ²School of Earth and Environmental Sciences, University of Manchester, UK. ³School of Materials, University of Manchester, UK.
Corresponding email: zoe.morland@open.ac.uk

Introduction: The composition of material delivered to the inner Solar System throughout history remains a mystery [1,2]. Lunar meteorites, unlike samples returned from Apollo and Luna missions, provide a rich resource for understanding the whole surface of the Moon [3]. Here we analyse a recently discovered lunar regolith breccia meteorites, North West Africa (NWA) 10989 (Fig. 1 A), to see whether it holds preserved impactor material. We consider the implications of its existence under the context of regolith evolution and *in-situ* resource utilisation.

Methods: At the University of Manchester, the petrology of NWA 10989 was analysed using the following methods: (1) an ESEM to obtain whole sample merger and close-up BSE images, qualitative EDX point spectra, and whole sample EDS elemental maps; (2) a Raman spectrometer to identify Fe-oxide phases; (3) an EPMA to quantitatively measure major element concentrations; (4) a Broad-ion beam (Argon) mill to polish; and (5) an Oxford Instruments EBSD with AZtec software to study crystallographic structure.

Methods: The 1 cm sized fragment of NWA 10989 we investigated specifically contains highland granulites, glassy impact melt breccias (GIMBs), clast-rich impact melt breccias (CIMBs), monomineralic pyroxene and rare mare basalt fragments. These clasts are held within a glassy matrix that includes impact melt spherules and Fe-Ni metal. The Fe-Ni metal is dominantly contained within a single 1.90×0.85 mm grain (Metal_1) along with several other smaller grains and abundant fine particles in the matrix (Figs. 1 A & B).

Results: The metal grains are composed of predominantly kamacite (low-Ni), enriched in K (0.16-0.22 wt%). Metal_1 also contains minor amounts of taenitic (high-Ni) material and K-rich schreibersite. Comparison between the metals' compositions and the meteoritic field suggest they are similar asteroid-derived meteorite groups, and thus could indicate and exogenous origin [4]. However, silicates in contact with and occasionally embedded within Metal_1 (Fig. 1 B) reflect native lunar compositions.

Metal_1's internal microstructure consists of an aggregate of 50-150 μm sized variably orientated grains (Fig. 2 A & B). The colour gradients within grains indicate the presence of lattice bending, which in places advanced to sub-grain development.

Discussion: The compositional dichotomy between the native lunar silicates and the sizeable, apparently exogenous, Fe-Ni metal suggests either (i) an almost intact fragment of meteorite or (i) a precipitate

from an impact melt sheet composed of a vaporised impactor and lunar target rock mixture. Metal_1's complex internal structure, indicative of crystal plastic deformation, could be attributed to strain during regolith reprocessing.

This study advances current knowledge in ISRU through evidence of impactor survival and modification on the surface of the Moon, plus it aids in understanding what material was delivered to the inner solar system during its history.




Figure 1: (A) Whole sample BSE image, (B) Metal_1 close-up BSE image with EBSD areas shown




Figure 2: Inverse pole figures of parts of Metal_1, indicated in Fig. 1 B, where colours represent lattice orientation corresponding to the triangle key.

Acknowledgements: Thanks to Martin Goff, for donating the sample.

References: [1] Hartmann W. et al. (2000) In: R. Canup and K. Righter, eds. Tuscon: University of Arizona Press, 805-826. [2] Joy K. H. (2012) *Science*, 336(6087), 1426-1429. [3] Jolliff B. L. et al. (2000) *JGR*, 105, 4197-4216. [4] Goldstein J. I. and Yakowitz H. (1971) *P. Lunar. Plant. Sci.*, 2, 177-191.