

# Open Research Online

---

The Open University's repository of research publications and other research outputs

## Limit measures for affine cellular automata II

### Journal Item

How to cite:

Pivato, Marcus and Yassawi, Reem (2004). Limit measures for affine cellular automata II. *Ergodic Theory and Dynamical Systems*, 24(6) pp. 1961–1980.

For guidance on citations see [FAQs](#).

© 2004 Cambridge University Press



<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Version: Accepted Manuscript

Link(s) to article on publisher's website:

<http://dx.doi.org/doi:10.1017/S0143385704000288>

---

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

---

[oro.open.ac.uk](http://oro.open.ac.uk)

# Limit Measures for Affine Cellular Automata II

Marcus Pivato

*Department of Mathematics  
University of Houston*

email: `pivato@math.toronto.edu`

and

Reem Yassawi\*

*Department of Mathematics,  
Trent University,*

email: `ryassawi@trentu.ca`

February 21, 2019

## Abstract

If  $\mathbb{M}$  is a monoid, and  $\mathcal{A}$  is an abelian group, then  $\mathcal{A}^{\mathbb{M}}$  is a compact abelian group; a *linear cellular automaton* (LCA) is a continuous endomorphism  $\mathfrak{F} : \mathcal{A}^{\mathbb{M}} \rightarrow \mathcal{A}^{\mathbb{M}}$  that commutes with all shift maps. If  $\mathfrak{F}$  is **diffusive**, and  $\mu$  is a **harmonically mixing** (HM) probability measure on  $\mathcal{A}^{\mathbb{M}}$ , then the sequence  $\{\mathfrak{F}^N \mu\}_{N=1}^{\infty}$  weak\*-converges to the Haar measure on  $\mathcal{A}^{\mathbb{M}}$ , in density. Fully supported Markov measures on  $\mathcal{A}^{\mathbb{Z}}$  are HM, and nontrivial LCA on  $\mathcal{A}^{(\mathbb{Z}^D)}$  are diffusive when  $\mathcal{A} = \mathbb{Z}/p$  is a prime cyclic group.

In the present work, we provide sufficient conditions for diffusion of LCA on  $\mathcal{A}^{(\mathbb{Z}^D)}$  when  $\mathcal{A} = \mathbb{Z}/n$  is any cyclic group or when  $\mathcal{A} = (\mathbb{Z}/p^r)^J$  ( $p$  prime). We show that any fully supported Markov random field  $\mathcal{A}^{(\mathbb{Z}^D)}$  is HM (where  $\mathcal{A}$  is any abelian group). We also provide examples of HM Markov measures *not* having full support, and measures on  $\mathcal{A}^{\mathbb{Z}}$  which have the Kolmogorov property but which are not HM.

---

\*This research partially supported by NSERC Canada.

# 1 Introduction

Let  $\mathcal{A}$  be a finite abelian group, with discrete topology. If  $\mathbb{M}$  is any set, then  $\mathcal{A}^{\mathbb{M}}$  is a compact abelian group when endowed with the Tychonoff product topology and componentwise addition. If  $\mathbb{M}$  is a monoid (for example, a lattice:  $\mathbb{Z}^D \times \mathbb{N}^E$ ), then the action of  $\mathbb{M}$  on itself by translation induces a natural **shift action** of  $\mathbb{M}$  on configuration space: for all  $\mathbf{e} \in \mathbb{M}$ , and  $\mathbf{a} \in \mathcal{A}^{\mathbb{M}}$ , define  $\sigma^{\mathbf{e}}[\mathbf{a}] = [b_{\mathbf{m}}]_{\mathbf{m} \in \mathbb{M}}$  where,  $\forall \mathbf{m} \in \mathbb{M}$ ,  $b_{\mathbf{m}} = a_{\mathbf{e} + \mathbf{m}}$ . Here “ $\cdot$ ” is the monoid operator (“ $+$ ” for  $\mathbb{M} = \mathbb{Z}^D \times \mathbb{N}^E$ ).

A **linear cellular automaton** (LCA) is a continuous endomorphism  $\mathfrak{F} : \mathcal{A}^{\mathbb{M}} \rightarrow \mathcal{A}^{\mathbb{M}}$  which commutes with all shift maps. If  $\mu$  is a measure on  $\mathcal{A}^{\mathbb{M}}$ , it is natural to consider the sequence of measures  $\{\mathfrak{F}^n \mu|_{n \in \mathbb{N}}\}$ , and ask whether this sequence converges in the weak\* topology on the space  $\mathcal{M}_{EAS}[\mathcal{A}^{\mathbb{M}}]$  of Borel probability measures on  $\mathcal{A}^{\mathbb{M}}$ . If  $\{\mathfrak{F}^n \mu|_{n \in \mathbb{N}}\}$  does not itself converge, we might hope at least for convergence in density (that is, convergence of a subsequence  $\{\mathfrak{F}^j \mu|_{j \in \mathbb{J}}\}$ , where  $\mathbb{J} \subset \mathbb{N}$  is a subset of Cesàro density 1), or

convergence of the Cesàro average  $\frac{1}{N} \sum_{n=1}^N \mathfrak{F}^n \mu$ .

Let  $\mathcal{H}^{aar}$  denote the Haar measure on  $\mathcal{A}^{\mathbb{M}}$ . Since  $\mathcal{H}^{aar}$  is invariant under the algebraic operations of  $\mathcal{A}^{\mathbb{M}}$ , it seems like a natural limit point for  $\{\mathfrak{F}^n \mu|_{n \in \mathbb{N}}\}$ . Indeed, D. Lind showed [4] that, if  $\mathcal{A} = \mathbb{Z}/2$ , and  $\mathfrak{F}$  is the automaton defined:  $\mathfrak{F}(\mathbf{a})_0 = a_{(-1)} + a_1$ , and  $\mu$  is any Bernoulli measure, then

$\mathbf{wk}^* \text{-} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \mathfrak{F}^n \mu = \mathcal{H}^{aar}$ . Lind also showed that  $\{\mathfrak{F}^n \mu|_{n \in \mathbb{N}}\}$  does *not* converge to  $\mathcal{H}^{aar}$ ; convergence fails along the subsequence  $\{\mathfrak{F}^{(2^n)} \mu|_{n \in \mathbb{N}}\}$ .

Later, Ferrari, Maass, Martinez, and Ney showed similar Cesàro convergence results in a variety of special cases [7, 6, 1]. Recently, Pivato and Yassawi [5] developed broad sufficient conditions for convergence. The concepts of **harmonic mixing** for measures and **diffusion** for LCA were introduced; if  $\mu$  is a harmonically mixing probability measure and  $\mathfrak{F}$  a diffusive LCA, then  $\{\mathfrak{F}^n \mu|_{n \in \mathbb{N}}\}$  weak\* converged to  $\mathcal{H}^{aar}$  in density, and thus, also in Cesàro mean.

This paper is a continuation of [5]. First we will extend the results on diffusion of LCA to a broader class of abelian groups: in §3, to the case when  $\mathcal{A} = \mathbb{Z}/n$ , for any  $n \in \mathbb{N}$ , and then in §4, to the case when  $\mathcal{A} = (\mathbb{Z}/p^r)^J$  ( $p$  prime,  $J, r \in \mathbb{N}$ ). Next, we extend the theory of harmonic mixing. In §5, we demonstrate harmonic mixing for any Markov random field on  $\mathcal{A}^{(\mathbb{Z}^D)}$

with full support. In §6, we show that full support is not necessary for harmonic mixing, by demonstrating harmonic mixing for a Markov measure on  $(\mathbb{Z}/2)^{\mathbb{Z}}$  *not* having full support. On the other hand, in §7, we construct a measure which, when seen as an ergodic dynamical system under the shift, is actually a  $\mathbf{K}$ -automorphism, but which nonetheless is *not* harmonically mixing.

## 2 Preliminaries

We recommend that the reader consult [5] before reading the present work; we will depend heavily upon results introduced there. We will now briefly review the relevant concepts; all theorems in this section are proved in [5].

### 2.1 Characters and Harmonic Mixing

Let  $\mathbb{T}^1$  be the unit circle group. A **character** of  $\mathcal{A}^{\mathbb{M}}$  is a continuous group homomorphism  $\phi : \mathcal{A}^{\mathbb{M}} \rightarrow \mathbb{T}^1$ . The set of all characters of  $\mathcal{A}^{\mathbb{M}}$  forms a group, denoted  $\widehat{\mathcal{A}^{\mathbb{M}}}$ .

If  $[\chi_m]_{m \in \mathbb{M}}$  is a sequence of characters of  $\mathcal{A}$ , with all but finitely many elements equal to the constant 1-function (denoted “ $\mathbf{1}$ ”), then define  $\chi = \bigotimes_{m \in \mathbb{M}} \chi_m : \mathcal{A}^{\mathbb{M}} \rightarrow \mathbb{T}^1$ ; thus, if  $\mathbf{a} = [a_m]_{m \in \mathbb{M}}$  is an element of  $\mathcal{A}^{\mathbb{M}}$ , then

$\chi(\mathbf{a}) = \prod_{m \in \mathbb{M}} \chi_m(a_m)$ . All elements of  $\widehat{\mathcal{A}^{\mathbb{M}}}$  arise in this manner. The **rank** of the character  $\chi$  is the number of nontrivial entries in the **coefficient system**  $[\chi_m]_{m \in \mathbb{M}}$ .

When  $\mathcal{A} = \mathbb{Z}/n$ , elements of  $\widehat{\mathcal{A}}$  are maps of the form  $\chi(a) = \exp\left(\frac{2\pi i}{n} c \cdot a\right)$ ,

where  $c \in \mathbb{Z}/n$  is some constant. Elements of  $\widehat{\mathcal{A}^{\mathbb{M}}}$  are then products of the form  $\chi(\mathbf{a}) = \prod_{m \in \mathbb{M}} \chi_m(a_m)$ , where,  $\forall m \in \mathbb{M}$ ,  $\chi_m : a \mapsto \exp\left(\frac{2\pi i}{n} c_m \cdot a\right)$  for some  $c_m \in \mathcal{A}$ , with all but finitely many  $c_m$  are equal to 0. In this case, we will use the term **coefficient system** also to describe the sequence  $[c_m]_{m \in \mathbb{M}}$ .

If  $\mu$  is a measure on  $\mathcal{A}^{\mathbb{M}}$ , then the **Fourier coefficients** of  $\mu$  are defined:

$\widehat{\mu}[\chi] = \langle \chi, \mu \rangle = \int_{\mathcal{A}^{\mathbb{M}}} \chi d\mu$ , for every  $\chi \in \widehat{\mathcal{A}^{\mathbb{M}}}$ . The measure  $\mu$  is called **harmonically mixing** if, for all  $\epsilon > 0$ , there is some  $R > 0$  so that, for all  $\chi \in \widehat{\mathcal{A}^{\mathbb{M}}}$ ,  $\left(\text{rank}[\chi] \geq R\right) \implies \left(|\widehat{\mu}[\chi]| < \epsilon\right)$ .

Let  $\mathcal{M}_{\text{EAS}} [\mathcal{A}^{\mathbb{M}}; \mathbb{C}]$  be the space of complex-valued Borel measures on  $\mathcal{A}^{\mathbb{M}}$ , treated as a Banach algebra under the total variation norm, with operations of addition and convolution. Let  $\mathcal{H} \subset \mathcal{M}_{\text{EAS}} [\mathcal{A}^{\mathbb{M}}; \mathbb{C}]$  be the set of harmonically mixing measures.

**Proposition 1:** *Let  $\mathcal{A}$  be any finite abelian group.*

1.  $\mathcal{H}$  is an ideal of  $\mathcal{M}_{\text{EAS}} [\mathcal{A}^{\mathbb{M}}; \mathbb{C}]$ .
2.  $\mathcal{H}$  is closed under the total variation norm and dense in the weak\* topology.
3.  $\mathcal{H}$  contains all Bernoulli measures  $\beta^{\otimes \mathbb{M}}$ , where  $\beta$  is a measure on  $\mathcal{A}$  such that, for any subgroup  $\mathcal{G} \subset \mathcal{A}$ , the support of  $\mu$  extends over more than one coset of  $\mathcal{G}$ .
4. If  $\mathbb{M} = \mathbb{Z}$ , then, for any  $N > 0$ ,  $\mathcal{H}$  contains all  $N$ -step Markov measures on  $\mathcal{A}^{\mathbb{Z}}$  giving nonzero probability to all elements of  $\mathcal{A}^{[0..N]}$ .
5.  $\mathcal{H}$  contains any measure absolutely continuous with respect to the aforementioned Bernoulli or Markov measures. \_\_\_\_\_□

## 2.2 Linear Cellular Automata

A **cellular automaton** (CA) is a continuous map  $\mathfrak{F} : \mathcal{A}^{\mathbb{M}} \rightarrow \mathcal{A}^{\mathbb{M}}$  that commutes with all shift maps. The Curtis-Hedlund-Lyndon Theorem [3] states that any CA is determined by a **local map**  $\mathfrak{f} : \mathcal{A}^{\mathbb{U}} \rightarrow \mathcal{A}$ , where  $\mathbb{U} \subset \mathbb{M}$  is some finite subset (a “neighbourhood of the identity”).  $\mathfrak{F}$  is an LCA if and only if  $\mathfrak{f}$  is a homomorphism from the product group  $\mathcal{A}^{\mathbb{U}}$  into  $\mathcal{A}$ .

The set  $\mathbf{End} [\mathcal{A}]$  of group endomorphisms from  $\mathcal{A}$  to itself is a ring under composition and pointwise addition: if  $\mathfrak{f}, \mathfrak{g}$  are in  $\mathbf{End} [\mathcal{A}]$ , then so are  $\mathfrak{f} \circ \mathfrak{g}$  and  $\mathfrak{f} + \mathfrak{g}$ . If  $\mathbf{Hom} [\mathcal{A}^{\mathbb{U}}; \mathcal{A}]$  is the set of group homomorphisms from  $\mathcal{A}^{\mathbb{U}}$  into  $\mathcal{A}$ , then there is a natural bijection between  $(\mathbf{End} [\mathcal{A}])^{\mathbb{U}}$  and  $\mathbf{Hom} [\mathcal{A}^{\mathbb{U}}; \mathcal{A}]$ , as follows: For each  $\mathbf{u} \in \mathbb{U}$ , suppose that  $\mathfrak{f}_{\mathbf{u}} \in \mathbf{End} [\mathcal{A}]$ . Define  $\mathfrak{f} : \mathcal{A}^{\mathbb{U}} \rightarrow \mathcal{A}$  by  $\mathfrak{f}[a_{\mathbf{u}}|_{\mathbf{u} \in \mathbb{U}}] = \sum_{\mathbf{u} \in \mathbb{U}} \mathfrak{f}_{\mathbf{u}}(a_{\mathbf{u}})$ . Then  $\mathfrak{f}$  is a group homomorphism, and every element of  $\mathbf{Hom} [\mathcal{A}^{\mathbb{U}}; \mathcal{A}]$  arises in this manner.

Thus, if  $\mathfrak{F}$  is an LCA, then there is some set of **coefficients**  $\{\mathfrak{f}_{\mathbf{u}}; \mathbf{u} \in \mathbb{U}\}$  so that, for any  $\mathbf{a} \in \mathcal{A}^{\mathbb{M}}$ ,  $\mathfrak{F}(\mathbf{a}) = \mathbf{b}$ , where for all  $\mathbf{m} \in \mathbb{M}$ ,  $b_{\mathbf{m}} = \sum_{\mathbf{u} \in \mathbb{U}} \mathfrak{f}_{\mathbf{u}}(a_{(\mathbf{m}, \mathbf{u})}) = \sum_{\mathbf{u} \in \mathbb{U}} \mathfrak{f}_{\mathbf{u}}(\sigma^{\mathbf{u}}(\mathbf{a})_{\mathbf{m}})$ .

For any  $\mathbf{u} \in \mathbb{U}$ , treat  $f_{\mathbf{u}}$  as an endomorphism on  $\mathcal{A}^{\mathbb{M}}$  by letting it act componentwise on elements of  $\mathcal{A}^{\mathbb{M}}$ . Then  $\forall \mathbf{a} \in \mathcal{A}^{\mathbb{M}}$ ,  $\mathfrak{F}(\mathbf{a}) = \sum_{\mathbf{u} \in \mathbb{U}} f_{\mathbf{u}} \circ \sigma^{\mathbf{u}}(\mathbf{a})$ . Thus,  $\mathfrak{F}$  can be written as a formal “polynomial of shift maps”:

$$\mathfrak{F} = \sum_{\mathbf{u} \in \mathbb{U}} f_{\mathbf{u}} \circ \sigma^{\mathbf{u}}.$$

If  $\mathcal{A} = \mathbb{Z}/n$  ( $n \in \mathbb{N}$ ), then the elements of  $\mathbf{End}[\mathcal{A}]$  are all maps of the form  $f([a]_n) = [f \cdot a]_n$ , where “[ $\bullet$ ] <sub>$n$</sub> ” refers to a mod- $n$  congruence class, and  $f \in \mathbb{Z}/n$  is a constant, with multiplication via the natural ring structure on  $\mathbb{Z}/n$ . In this case, we can write  $\mathfrak{F} = \sum_{\mathbf{u} \in \mathbb{U}} f_{\mathbf{u}} \cdot \sigma^{\mathbf{u}}$ , a polynomial with coefficients in  $\mathbb{Z}/n$ . For example, if  $\mathbb{M} = \mathbb{Z}$  and  $\mathfrak{F} = \sigma^{-1} + 3 \circ \sigma^1 + 5 \sigma^2$ , then this means that  $\mathfrak{F}(\mathbf{a})_k = [a_{(k-1)} + 3 \cdot a_{(k+1)} + 5 \cdot a_{(k+2)}]_n$ .

### 2.3 Diffusion

If  $\mathfrak{F} : \mathcal{A}^{\mathbb{M}} \curvearrowright$  is an LCA and  $\chi$  is a character of  $\mathcal{A}^{\mathbb{M}}$ , then  $\chi \circ \mathfrak{F}$  is also a character.  $\mathfrak{F}$  is called **diffusive**<sup>1</sup> if, for every nontrivial  $\chi \in \widehat{\mathcal{A}^{\mathbb{M}}}$ , there is some subset  $\mathbb{J}_{\chi} \subset \mathbb{N}$  of density 1 so that  $\lim_{\substack{j \rightarrow \infty \\ j \in \mathbb{J}_{\chi}}} \text{rank}[\chi \circ \mathfrak{F}^j] = \infty$ . We will abbreviate this to “rank  $[\chi \circ \mathfrak{F}^N] \xrightarrow[N \rightarrow \infty]{\text{dense}} \infty$ ”.

**Theorem 2:** *Let  $p$  be a prime number, and  $\mathcal{A} = \mathbb{Z}/p$ . Let  $D \geq 1$ . Then any nontrivial LCA on  $\mathcal{A}^{(\mathbb{Z}^D)}$  is diffusive. □*

By **nontrivial** we mean that  $\mathfrak{F}$ , as a polynomial of shift maps, has more than one nontrivial coefficient. The significance of diffusion and harmonic mixing is the following:

**Theorem 3:** *Let  $\mathcal{A}$  be a finite abelian group, and  $\mathbb{M}$  a countable monoid. Suppose that  $\mathfrak{F} : \mathcal{A}^{\mathbb{M}} \curvearrowright$  is an LCA, and that  $\mu$  is a harmonically mixing measure on  $\mathcal{A}^{\mathbb{M}}$ . If  $\mathfrak{F}$  is diffusive, then there is a set  $\mathbb{J} \subset \mathbb{N}$  of Cesàro density 1 so that  $\mathbf{wk}^* \text{-} \lim_{\substack{j \rightarrow \infty \\ j \in \mathbb{J}}} \mathfrak{F}^j \mu = \mathcal{H}^{\text{avr}}$ . Thus,  $\mathbf{wk}^* \text{-} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \mathfrak{F}^n \mu = \mathcal{H}^{\text{avr}}$ . □*

<sup>1</sup>In [5], this was called **diffusion in density**. Since diffusion in density is the only kind we will encounter in this paper, we have opted for more concise terminology.

For example,  $\mathfrak{F}^n \mu$  weak\*-converges to Haar measure in density whenever  $\mu$  is one of the aforementioned Bernoulli or  $N$ -step Markov measures.

### 3 Diffusion on other cyclic groups

Suppose  $n = p_1^{r_1} \dots p_J^{r_J}$ , where  $p_1, \dots, p_J$  are distinct primes and  $r_1, \dots, r_J \in \mathbb{N}$ . Let  $\mathcal{A} = \mathbb{Z}/n$ , and  $\mathcal{A}_j = \mathbb{Z}/q_j$ , with  $q_j = p_j^{r_j}$ , for  $j \in [1..J]$ . Then  $\mathcal{A} \cong$

$\bigoplus_{j=1}^J \mathcal{A}_j$ , and thus,  $\widehat{\mathcal{A}} \cong \bigoplus_{j=1}^J \widehat{\mathcal{A}}_j$ . There is then a canonical identification:

$\mathcal{A}^{\mathbb{M}} \cong \bigoplus_{j=1}^J \mathcal{A}_j^{\mathbb{M}}$ , and thus,  $\widehat{\mathcal{A}^{\mathbb{M}}} \cong \bigoplus_{j=1}^J \widehat{\mathcal{A}}_j^{\mathbb{M}}$ . Concretely: if  $\chi \in \widehat{\mathcal{A}^{\mathbb{M}}}$  has

coefficient sequence  $[c_{\mathbf{m}}]_{\mathbf{m} \in \mathbb{M}}$ , then,  $\chi \cong \bigoplus_{j=1}^J \chi^{[j]}$ , where for each  $j \in [1..J]$ ,  $\chi^{[j]} \in \widehat{\mathcal{A}}_j^{\mathbb{M}}$  has coefficient sequence  $[c_{\mathbf{m}}^{[j]}]_{\mathbf{m} \in \mathbb{M}}$ , with  $c_{\mathbf{m}}^{[j]} = [c_{\mathbf{m}}]_{q_j}$  for all  $\mathbf{m} \in \mathbb{M}$ .

$$\begin{aligned} \text{Also, } \mathbf{End}[\mathcal{A}] &\cong \bigoplus_{i,j=1}^J \mathbf{Hom}[\mathcal{A}_i, \mathcal{A}_j] = \bigoplus_{j=1}^J \mathbf{End}[\mathcal{A}_j] \text{ (cross-terms are trivial)} \\ &\cong \bigoplus_{j=1}^J \mathbb{Z}/q_j. \end{aligned}$$

Concretely, if  $f \in \mathbb{N}$ , and  $\mathfrak{f} : \mathcal{A} \curvearrowright$  is the map  $[a]_n \mapsto [f \cdot a]_n$ , then  $\mathfrak{f} = \mathfrak{f}^{[1]} \oplus \dots \oplus \mathfrak{f}^{[J]}$ , where, for each  $j$ ,  $\mathfrak{f}^{[j]} : \mathcal{A}_j \curvearrowright$  is the map  $[a]_{q_j} \mapsto [f_j \cdot a]_{q_j}$ , and where  $f \equiv f_j \pmod{p}$ . In particular, if  $q_j$  divides  $f$ , then  $\mathfrak{f}^{[j]}$  is trivial.

Thus, if  $\mathfrak{F} : \mathcal{A}^{\mathbb{M}} \curvearrowright$  is the LCA  $\sum_{\mathbf{u} \in \mathbb{U}} \mathfrak{f}_{\mathbf{u}} \circ \sigma^{\mathbf{u}}$ , with  $\mathfrak{f}_{\mathbf{u}} \in \mathbf{End}[\mathcal{A}]$ , then,

$\forall \mathbf{u} \in \mathbb{U}$ , we can write  $\mathfrak{f}_{\mathbf{u}} = \mathfrak{f}_{\mathbf{u}}^{[1]} \oplus \dots \oplus \mathfrak{f}_{\mathbf{u}}^{[J]}$ , with  $\mathfrak{f}_{\mathbf{u}}^{[j]} \in \mathbf{End}[\mathcal{A}_j]$  a scalar-multiplication map determined by some  $f_{\mathbf{u}}^{[j]} \in \mathbb{Z}/q_j$ , and then write  $\mathfrak{F} = \bigoplus_{j=1}^J \mathfrak{F}^{[j]}$ , where,  $\forall j \in [1..J]$ ,  $\mathfrak{F}^{[j]} : \mathcal{A}_j^{\mathbb{M}} \curvearrowright$  is the LCA given by  $\sum_{\mathbf{u} \in \mathbb{U}} f_{\mathbf{u}}^{[j]} \circ \sigma^{\mathbf{u}}$ .

Note also that, if  $\chi = \bigoplus_{j=1}^J \chi_j \in \widehat{\mathcal{A}^{\mathbb{M}}}$ , then  $\chi \circ \mathfrak{F} = \bigoplus_{j=1}^J (\chi_j \circ \mathfrak{F}^{[j]})$ .

**Lemma 4:** If  $\mathfrak{F} = \bigoplus_{j=1}^J \mathfrak{F}^{[j]}$  is an LCA on  $\bigoplus_{j=1}^J \mathcal{A}_j^{\mathbb{M}}$ , then

$$\left( \mathfrak{F} \text{ is diffusive} \right) \iff \left( \forall j \in [1..J], \mathfrak{F}^{[j]} \text{ is diffusive.} \right)$$

**Proof:**

**Proof of “ $\Leftarrow$ ”:** Let  $\chi \in \widehat{\mathcal{A}^{\mathbb{M}}}$  be nontrivial. Thus,  $\chi = \chi^{[1]} \oplus \dots \oplus \chi^{[J]}$ , where at least one of  $\chi^{[j]} \in \widehat{\mathcal{A}_j^{\mathbb{M}}}$ , is nontrivial; suppose it is  $\chi^{[j_0]}$ . Since  $\mathfrak{F}^{[j_0]}$  is diffusive, we conclude:  $\text{rank} [\chi \circ \mathfrak{F}^N] \geq \text{rank} [\chi^{[j_0]} \circ (\mathfrak{F}^{j_0})^N] \xrightarrow[n \rightarrow \infty]{\text{dense}} \infty$ .

**Proof of “ $\Rightarrow$ ”:** Suppose that  $\mathfrak{F}_{j_0}$  is not diffusive. Let  $\chi_{j_0}$  be some character on  $\mathcal{A}_{j_0}^{\mathbb{M}}$  so that  $\text{rank} [\chi_{j_0} \circ \mathfrak{F}_{j_0}] \xrightarrow[n \rightarrow \infty]{\text{dense}} \infty$ , and let  $\chi = \bigoplus_{j=1}^J \chi_j$ , where  $\chi_j = \mathbf{1}$  for all  $j \neq j_0$ . Then  $\text{rank} [\chi \circ \mathfrak{F}] = \text{rank} [\chi_{j_0} \circ \mathfrak{F}_{j_0}] \xrightarrow[n \rightarrow \infty]{\text{dense}} \infty$ , so  $\mathfrak{F}$  is not diffusive.  $\underline{\hspace{10em}} \square$

Hence, we have reduced the proof of diffusion to the prime power case.

Suppose  $\mathcal{A} = \mathbb{Z}/8$ , and let  $\mathfrak{F} = \mathbf{Id} + 2\sigma^1$  act on  $\mathcal{A}^{\mathbb{Z}}$ . Then  $\mathfrak{F}^{4 \cdot N} = \mathbf{Id}$  for all  $N \in \mathbb{N}$ , so  $\mathfrak{F}$  cannot be diffusive. This motivates the conditions of the following theorem.

**Lemma 5:** Suppose  $\mathcal{A} = \mathbb{Z}/q$ , where  $q = p^r$ , with  $p$  prime and  $r \in \mathbb{N}$ . Let  $\mathbb{M} = \mathbb{Z}^D$ , and  $\mathfrak{F} = \sum_{\mathbf{u} \in \mathbb{U}} f_{\mathbf{u}} \circ \sigma^{\mathbf{u}}$ . If  $f_{\mathbf{u}} \in [0..q)$  are relatively prime to  $p$  for at least two  $\mathbf{u} \in \mathbb{U}$ , then  $\mathfrak{F}$  is diffusive.

**Proof:** Let  $\chi \in \widehat{\mathcal{A}^{\mathbb{M}}}$  have coefficient sequence  $[c_{\mathbf{v}}]_{\mathbf{v} \in \mathbb{V}}$ , where  $c_{\mathbf{v}} \in \mathbb{Z}/q$ , for all  $\mathbf{v} \in \mathbb{V}$ , with  $\mathbb{V} \subset \mathbb{M}$  some finite subset. Thus,  $\chi^{[N]} = \chi \circ \mathfrak{F}^N$  has coefficient sequence  $[c_{\mathbf{m}}^{[N]}]_{\mathbf{m} \in \mathbb{M}}$ , where, for all  $\mathbf{m} \in \mathbb{M}$ ,

$$c_{\mathbf{m}}^{[N]} = \sum_{\mathbf{v} \in \mathbb{V}} \sum_{\substack{\mathbf{u}_1, \dots, \mathbf{u}_N \in \mathbb{U} \\ \mathbf{v} + \mathbf{u}_1 + \dots + \mathbf{u}_N = \mathbf{m}}} c_{\mathbf{v}} \cdot f_{\mathbf{u}_1} \cdot \dots \cdot f_{\mathbf{u}_N} \quad (1)$$

Thus,  $\text{rank} [\chi \circ \mathfrak{F}^N]$  is the number of these coefficients that are nonzero, mod  $q$ .

**Case 1:** One of the coefficients  $\{c_{\mathbf{v}}\}_{\mathbf{v} \in \mathbb{V}}$  is nonzero, mod  $p$ .



Consider the character  $\chi_{/p}$  and the (nontrivial) LCA  $\mathfrak{F}_{/p}$  on  $\mathbb{Z}_{/p}^{\mathbb{M}}$  induced by the coefficients  $[c_v|_{v \in \mathbb{V}}]$  and  $[f_u|_{u \in \mathbb{U}}]$  respectively, and, for all  $N \in \mathbb{N}$ , the character  $\chi_{/p}^{[N]}$  induced by  $[c_m^{[N]}|_{m \in \mathbb{M}}]$ .

First, note that  $\forall N \in \mathbb{N}$ ,  $\chi_{/p}^{[N]} = \chi_{/p} \circ \mathfrak{F}_{/p}^N$  (simply consider equation (1), only mod  $p$  instead). Notice that, for any  $m$  and  $N$ , if the expression in (1) is nonzero mod  $p$ , then it must be nonzero mod  $q$ . Thus  $\text{rank} [\chi^{[N]}] \geq \text{rank} [\chi_{/p}^{[N]}] = \text{rank} [\chi_{/p} \circ \mathfrak{F}_{/p}^N]$ . Hence, it suffices to show that  $\text{rank} [\chi_{/p} \circ \mathfrak{F}_{/p}^N] \xrightarrow[N \rightarrow \infty]{\text{dense}} \infty$ .

But one of  $\{c_v|_{v \in \mathbb{V}}\}$  is nonzero, mod  $p$ , so  $\chi_{/p}$  is nontrivial as a character on  $\mathbb{Z}_{/p}^{\mathbb{M}}$ . Thus, by Theorem 2,  $\text{rank} [\chi_{/p} \circ \mathfrak{F}_{/p}^N] \xrightarrow[N \rightarrow \infty]{\text{dense}} \infty$ .

**Case 2:** All the coefficients  $\{c_v|_{v \in \mathbb{V}}\}$  are divisible by  $p$ .

Let  $p^s$  be the greatest power of  $p$  that divides all elements of  $\{c_v|_{v \in \mathbb{V}}\}$ ; clearly  $s < r$ . Let  $\tilde{r} = r - s$  and  $\tilde{q} = p^{\tilde{r}}$ , and let  $\tilde{\mathcal{A}} = \mathbb{Z}_{/\tilde{q}}$ . We will reduce the problem to consideration of an LCA on  $\mathbb{Z}_{/\tilde{q}}$ , and then apply **Case 1**.

For all  $v \in \mathbb{V}$ , let  $\tilde{c}_v = c_v/p^s$ , and let  $\tilde{\chi} \in \widehat{\tilde{\mathcal{A}}^{\mathbb{M}}}$  be the corresponding character. Let  $\tilde{\mathfrak{F}}$  be the LCA on  $\tilde{\mathcal{A}}^{\mathbb{M}}$  having the same coefficients as  $\mathfrak{F}$ ; thus,  $\tilde{\chi}^{[N]} = \tilde{\chi} \circ \tilde{\mathfrak{F}}^N$  has coefficient sequence  $[\tilde{c}_m^{[N]}|_{m \in \mathbb{M}}]$ , where, for all  $m \in \mathbb{M}$ ,  $\tilde{c}_m^{[N]} = \sum_{v \in \mathbb{V}} \sum_{\substack{u_1, \dots, u_N \in \mathbb{U} \\ v + u_1 + \dots + u_N = m}} \tilde{c}_v \cdot f_{u_1} \cdot \dots \cdot f_{u_N}$ .

Clearly, for all  $N \in \mathbb{N}$  and  $m \in \mathbb{M}$ ,  $c_m^{[N]} = p^s \cdot \tilde{c}_m^{[N]}$ , so if  $\tilde{c}_m^{[N]} \not\equiv 0 \pmod{\tilde{q}}$ , then  $c_m^{[N]} \not\equiv 0 \pmod{q}$ . Thus,  $\text{rank} [\chi^{[N]}] \geq \text{rank} [\tilde{\chi}^{[N]}]$ . But by construction, at least one coefficient of  $\tilde{\chi}$  is nonzero, mod  $p$ . Thus, by **Case 1**, we have:  $\text{rank} [\tilde{\chi} \circ \tilde{\mathfrak{F}}^N] \xrightarrow[N \rightarrow \infty]{\text{dense}} \infty$ .  $\square$

**Theorem 6:** Let  $n \in \mathbb{N}$ , and  $\mathcal{A} = \mathbb{Z}_{/n}$ . Let  $D \geq 1$ , and let  $\mathfrak{F} : \mathcal{A}^{(\mathbb{Z}^D)} \curvearrowright$  be an LCA such that, for each prime divisor  $p$  of  $n$ , at least two coefficients of  $\mathfrak{F}$  are relatively prime to  $p$ . Then  $\mathfrak{F}$  is diffusive.

**Proof:** Write  $n = p_1^{r_1} \cdot \dots \cdot p_J^{r_J}$ ,  $\mathcal{A} = \mathcal{A}_1 \oplus \dots \oplus \mathcal{A}_J$ ,  $\mathfrak{F} = \mathfrak{F}_1 \oplus \dots \oplus \mathfrak{F}_J$  as before. By Lemma 4, it suffices to show that each of  $\mathfrak{F}_1, \dots, \mathfrak{F}_J$  is diffusive. By Lemma 5 and the hypothesis, this is the case.  $\square$

## 4 Diffusion on finite abelian groups

Now suppose  $\mathcal{A}$  is an arbitrary finite abelian group. Then  $\mathcal{A}$  has a canonical decomposition:  $\mathcal{A} = \bigoplus_{k=1}^K \bigoplus_{j=1}^{J_k} \mathcal{A}_{(k,j)}$ , with  $\mathcal{A}_{(k,j)} = \mathbb{Z}/q_{(k,j)}$ ;  $q_{(k,j)} = p_k^{r_{(k,j)}}$ , where  $p_1, \dots, p_K$  are distinct primes with  $r_{(k,1)}, r_{(k,2)}, \dots, r_{(k,J_k)}$  natural numbers for each  $k \in [1..K]$ .

We will assume that  $\mathcal{A}$  is of the special form where, for all  $k \in [1..K]$ ,  $r_{k,1} = \dots = r_{k,J_k} = r_k$ . In other words,  $\mathcal{A} = \bigoplus_{k=1}^K \mathcal{A}_k$ , with  $\mathcal{A}_k = (\mathbb{Z}/q_k)^{J_k}$ , where  $p_1, \dots, p_K$  are distinct primes, with  $q_k = p_k^{r_k}$ , and  $r_k, J_k \in \mathbb{N}$ . Thus, as before,  $\mathbf{End}[\mathcal{A}] = \bigoplus_{j,k=1}^K \mathbf{Hom}[\mathcal{A}_j, \mathcal{A}_k] = \bigoplus_{k=1}^K \mathbf{End}[\mathcal{A}_k]$ , (cross-terms are trivial), and  $\mathcal{A}^{\mathbb{M}} \cong \mathcal{A}_1^{\mathbb{M}} \oplus \dots \oplus \mathcal{A}_K^{\mathbb{M}}$ , so we can write any LCA  $\mathfrak{F} : \mathcal{A}^{\mathbb{M}} \curvearrowright$  as a direct sum  $\mathfrak{F} = \mathfrak{F}_1 \oplus \dots \oplus \mathfrak{F}_K$ , where  $\mathfrak{F}_k : \mathcal{A}_k^{\mathbb{M}} \curvearrowright$ .

By Lemma 4, to prove  $\mathfrak{F}$  is diffusive, it suffices to show that each of  $\mathfrak{F}_1, \dots, \mathfrak{F}_K$  is diffusive. Hence, we will assume from now on that  $\mathcal{A} = (\mathbb{Z}/q)^J$ , where  $p$  is prime,  $q = p^r$ , and  $J \in \mathbb{N}$ . Elements of  $\mathcal{A}$  are thought of as  $J$ -tuples of  $\mathbb{Z}/q$ -elements.  $\mathcal{A}$  is a  $J$ -dimensional module over the commutative ring<sup>2</sup>  $\mathbb{Z}/q$ . The endomorphisms of  $\mathcal{A}$  as an abelian group are just the  $\mathbb{Z}/q$ -linear endomorphisms of this  $\mathbb{Z}/q$ -module, and are described by  $J \times J$  matrices of elements in  $\mathbb{Z}/q$ .

**Lemma 7:** *Let  $\mathcal{A} = (\mathbb{Z}/q)^J$ , where  $p$  is prime and  $q = p^r$ .*

1. Any  $\chi \in \widehat{\mathcal{A}}$  is of the form:  $\chi(\mathbf{a}) = \exp\left(\frac{2\pi i}{q} \cdot \langle \mathbf{c}, \mathbf{a} \rangle\right)$ , where  $\mathbf{c} = (c_1, \dots, c_J) \in (\mathbb{Z}/q)^J$ , and for any  $\mathbf{a} = (a_1, \dots, a_J) \in (\mathbb{Z}/q)^J$ , we define  $\langle \mathbf{c}, \mathbf{a} \rangle = c_1 a_1 + \dots + c_J a_J$ . Thus,  $\chi$  is nontrivial if and only if  $\mathbf{c} \neq \mathbf{0}$ .
2. If  $\mathfrak{f} \in \mathbf{End}[\mathcal{A}]$  has matrix  $\mathbf{F}$  with adjoint  $\dagger \mathbf{F}$ , then  $\chi \circ \mathfrak{f}$  is the character  $\mathbf{a} \mapsto \exp\left(\frac{2\pi i}{q} \cdot \langle \mathbf{c}', \mathbf{a} \rangle\right)$ , where  $\mathbf{c}' = \dagger \mathbf{F} \cdot \mathbf{c}$ .

*In particular,  $\chi \circ \mathfrak{f}$  is nontrivial if and only if  $\mathbf{c}$  is not in  $\ker[\dagger \mathbf{F}]$ .*

3. Let  $\mathfrak{f} \in \mathbf{Aut}[\mathcal{A}]$ . If  $\chi \in \widehat{\mathcal{A}}$  is nontrivial then  $\chi \circ \mathfrak{f}$  is also nontrivial.  $\square$

<sup>2</sup>If  $r = 1$  then  $q = p$  is prime,  $\mathbb{Z}/q$  is a field, and  $\mathcal{A}$  is a  $\mathbb{Z}/q$ -vector space. It may be helpful to keep this case in mind in what follows.

Let  $\mathbb{V} \subset \mathbb{M}$  be a subset not containing 0. If  $\mathfrak{G} = \sum_{m \in \mathbb{M}} \mathfrak{g}_m \sigma^m$  is an LCA on  $\mathcal{A}^{\mathbb{M}}$ , then a subset  $\mathbb{W} \subset \mathbb{M}$  is called  **$\mathbb{V}$ -separating** for  $\mathfrak{G}$  if, for every  $w \in \mathbb{W}$ ,  $\mathfrak{g}_w \in \mathbf{Aut}[\mathcal{A}]$ , but for all  $v \in \mathbb{V}$ ,  $\mathfrak{g}_{(w-v)} = 0$ . Intuitively,  $\mathbb{W}$  indexes a set of nontrivial (indeed, automorphic) coefficients of  $\mathfrak{G}$ , separated from one another by  $\mathbb{V}$ -shaped ‘‘gaps’’. If  $\mathbb{U} = \mathbb{V} \sqcup \{0\}$ , and  $\chi = \bigotimes_{u \in \mathbb{U}} \chi_u$  is a character, then we will show that these gaps ensure that  $(\chi \circ \mathfrak{G})_w$  is nontrivial, for all  $w \in \mathbb{W}$ . We will then construct  $\mathbb{V}$ -separating sets for  $\mathfrak{G} = \mathfrak{F}^N$ . This argument was already used implicitly to prove Theorem 15 in [5].

**Proposition 8:** *Let  $\mathbb{M} = \mathbb{Z}^D$ . An LCA  $\mathfrak{F} : \mathcal{A}^{\mathbb{M}} \curvearrowright$  is diffusive if, for every finite subset  $\mathbb{V} \subset \mathbb{M}$  not containing zero, and every  $R \in \mathbb{N}$ , there is a set  $\mathbb{J}_{(\mathbb{V}; R)} \subset \mathbb{N}$  of density 1 so that, for all  $j \in \mathbb{J}_{(\mathbb{V}; R)}$  there is a  $\mathbb{V}$ -separating set  $\mathbb{W}_j \subset \mathbb{M}$  for  $\mathfrak{F}^j$  with  $\mathcal{C}_{\text{ord}}[\mathbb{W}_j] > R$ .*

**Proof:** Suppose  $\mathfrak{F}$  is not diffusive; thus, there is some character  $\chi = \prod_{u \in \mathbb{U}} \chi_u$

so that  $\text{rank}[\chi \circ \mathfrak{F}^N] \xrightarrow[N \rightarrow \infty]{\text{dense}} \infty$ ; hence, there is some subset  $\mathbb{B} \subset \mathbb{N}$  of nonzero upper density and some bound  $R$  so that  $\text{rank}[\chi \circ \mathfrak{F}^N] < R$  for all  $N \in \mathbb{B}$ .

Fix  $u_0 \in \mathbb{U}$  and let  $\mathbb{V} = \{u - u_0; u \in \mathbb{U} \setminus \{u_0\}\}$ ; let  $\mathbb{J}_{(\mathbb{V}; R)}$  be the set described by the hypothesis. The set  $\mathbb{B} \subset \mathbb{N}$  has nonzero upper density, so  $\mathbb{B} \cap \mathbb{J}_{(\mathbb{V}; R)} \neq \emptyset$ ; let  $j \in \mathbb{B} \cap \mathbb{J}_{(\mathbb{V}; R)}$ , and let  $\mathbb{W}_j \subset \mathbb{M}$  be the  $\mathbb{V}$ -separating set for  $\mathfrak{F}^j$ .

Write  $\mathfrak{F}^j = \sum_{m \in \mathbb{M}} f_m^{[j]} \sigma^m$ , and then write  $\chi \circ \mathfrak{F}^j = \prod_{m \in \mathbb{M}} \chi_m^{[j]}$ , where  $\chi_m^{[j]} = \prod_{u \in \mathbb{U}} (\chi_u \circ f_{m-u}^{[j]})$ . Then  $\forall w \in \mathbb{W}_j$ ,  $\chi_{(w+u_0)}^{[j]} = \prod_{u \in \mathbb{U}} (\chi_u \circ f_{(w+u_0)-u}^{[j]}) = (\chi_{u_0} \circ f_w^{[j]}) \cdot \prod_{v \in \mathbb{V}} (\chi_{(v+u_0)} \circ f_{(w-v)}^{[j]}) = (\chi_{u_0} \circ f_w^{[j]}) \cdot \left( \prod_{v \in \mathbb{V}} \mathbf{1} \right) = (\chi_{u_0} \circ f_w^{[j]})$ ,

which is nontrivial by Lemma 7, because  $f_w^{[j]}$  is an automorphism. Thus,  $\chi_w^{[j]} \neq \mathbf{1}$  for all  $w \in \mathbb{W} + u_0$ , a set of cardinality greater than  $R$ , contradicting the hypothesis that  $\text{rank}[\chi \circ \mathfrak{F}^j] < R$ .  $\square$

Applying Proposition 8 often involves tracking binomial coefficients, mod  $p$ , via Lucas’ Theorem [5]. For a fixed prime  $p$ , and any  $n \in \mathbb{N}$ , let  $\mathbb{P}(n) \in$

$[0..p]^{\mathbb{N}}$  be the  $p$ -ary expansion of  $n$  (conventionally written with digits in reversed order). Thus, for example, if  $p = 3$ , then  $\mathbb{P}(34) = \dots 0000001021$ .

If  $n, N \in \mathbb{N}$ , with  $\mathbb{P}(n) = [n^{[i]}]_{i=0}^{\infty}$  and  $\mathbb{P}(N) = [N^{[i]}]_{i=0}^{\infty}$  then we write “ $n \ll N$ ” if  $n^{[i]} \leq N^{[i]}$  for all  $i \in \mathbb{N}$ . Lucas’ Theorem then implies:

$$\left( \begin{bmatrix} N \\ n \end{bmatrix}_p \neq 0 \right) \iff (n \ll N)$$

A **commuting automorphism linear cellular automaton** is an LCA of the form  $\mathfrak{F} = \sum_{u \in \mathbb{U}} f_u \circ \sigma^u$ , where  $\{f_u|_{u \in \mathbb{U}}\} \subset \mathbf{Aut}[\mathcal{A}]$  is a commuting collection of automorphisms of  $\mathcal{A}$ . For example:

- $\{f_u|_{u \in \mathbb{U}}\}$  are simultaneously diagonalizable automorphisms. In other words, there is some  $\mathbb{Z}/q$ -basis  $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_J\}$  for  $\mathcal{A}$ , so that the elements of  $\mathcal{B}$  are eigenvectors for every element of  $\{f_u|_{u \in \mathbb{U}}\}$ , and all eigenvalues are relatively prime to  $p$ .
- There is some  $f \in \mathbf{Aut}[\mathcal{A}]$  so that  $\forall u \in \mathbb{U}$ ,  $f_u = f^{n_u}$  for some  $n_u \in \mathbb{Z}$ .

**Theorem 9:** *If  $\mathfrak{G} : \mathcal{A}^{(\mathbb{Z}^D)} \rightrightarrows$  is a commuting automorphism LCA with two or more nontrivial coefficients, then  $\mathfrak{G}$  is diffusive.*

**Proof:** We will use Proposition 8; the argument is basically identical to the proof of Theorem 15 in [5], so we will only sketch it here.

Suppose  $\mathfrak{G} = \mathfrak{g}_0 \sigma^{n_0} + \mathfrak{g}_1 \sigma^{n_1} + \dots + \mathfrak{g}_U \sigma^{n_U}$ , where  $\mathfrak{g}_0, \mathfrak{g}_1, \dots, \mathfrak{g}_U \in \mathbf{Aut}[\mathcal{G}]$  commute, and where  $n_0, n_1, \dots, n_U \in \mathbb{Z}^D$ . We can rewrite:  $\mathfrak{G} = \mathfrak{g}_0 \circ (\mathfrak{F} \circ \sigma^{n_0})$ , where:

$$\mathfrak{F} = \mathbf{Id} + f_1 \sigma^{m_1} \left( \mathbf{Id} + f_2 \sigma^{m_2} \left[ \dots (\mathbf{Id} + f_{U-1} \sigma^{m_{U-1}} [\mathbf{Id} + f_U \sigma^{m_U}]) \dots \right] \right),$$

and, for all  $u \in [1..U]$ ,  $m_u = n_u - n_{u-1}$ , and  $f_u = \mathfrak{g}_{u-1}^{-1} \circ \mathfrak{g}_u$ . We can do this because  $\mathfrak{g}_0, \mathfrak{g}_1, \dots, \mathfrak{g}_U$  are automorphisms, and thus, invertible. It suffices to show that  $\mathfrak{F}$  is diffusive.

Let  $J \in \mathbb{N}$ . The coefficients of  $\mathfrak{F}$  commute, so we can employ the Binomial Theorem—and thus, Lucas’ Theorem—to compute the coefficients of  $\mathfrak{F}^J$ , mod  $p$ .

Let  $\mathcal{L}^U(J) = \{[k_1, k_2, \dots, k_U] \in \mathbb{N}^U ; k_U \ll k_{U-1} \ll \dots k_2 \ll k_1 \ll J\}$ .

Then  $\mathfrak{F}^J = \sum_{\mathbf{n} \in \mathbb{Z}^D} \mathfrak{f}_{\mathbf{n}}^{[J]} \circ \sigma^{\mathbf{n}}$ , where  $\mathfrak{f}_{\mathbf{n}}^{[J]} = \sum_{\substack{\mathbf{k} \in \mathcal{L}^U(J) \\ (k_1 \mathbf{m}_1 + \dots + k_U \mathbf{m}_U) = \mathbf{n}}} \mathfrak{f}_{(\mathbf{k})}^{[J]}$ , and, for

any  $\mathbf{k} = [k_1, k_2, \dots, k_U] \in \mathbb{N}^U$ , we define

$$\mathfrak{f}_{(\mathbf{k})}^{[J]} := \begin{bmatrix} J \\ k_1 \end{bmatrix}_p \begin{bmatrix} k_1 \\ k_2 \end{bmatrix}_p \cdots \begin{bmatrix} k_{U-1} \\ k_U \end{bmatrix}_p \mathfrak{f}_1^{k_1} \circ \mathfrak{f}_2^{k_2} \circ \dots \circ \mathfrak{f}_U^{k_U}.$$

(See [5] for details.)

Fix a finite subset  $\mathbb{V} \subset \mathbb{Z}^D$  not containing 0, and let  $R > 0$ ; we want to build a  $\mathbb{V}$ -separating set for  $\mathfrak{F}^J$  of cardinality  $R$ . To do this, note that there is some  $\Gamma \in \mathbb{N}$  such that, if  $J \in \mathbb{N}$  and  $\mathbb{P}(J)$  contains at least  $R$  “gaps” of size at least  $\Gamma$  (ie. sequences of  $\Gamma$  successive zeros, delimited by nonzero entries), then we can construct a set  $\mathbb{W}_J \subset \mathbb{Z}^D$  with  $\mathcal{C}_{\text{ad}}[\mathbb{W}_J] \geq R$ , so that:

1. For every  $\mathbf{w} \in \mathbb{W}_J$ , there is a unique  $\mathbf{k} \in \mathcal{L}^U(J)$  so that  $(k_1 \mathbf{m}_1 + \dots + k_U \mathbf{m}_U) = \mathbf{w}$ ; thus  $\mathfrak{f}_{\mathbf{w}}^{[J]} = \mathfrak{f}_{(\mathbf{k})}^{[J]} \in \mathbf{Aut}[\mathcal{A}]$ .
2. For all  $\mathbf{v} \in \mathbb{V}$ , there are no  $\mathbf{k} \in \mathcal{L}^U(J)$  with  $(k_1 \mathbf{m}_1 + \dots + k_U \mathbf{m}_U) = \mathbf{w} - \mathbf{v}$ ; thus  $\mathfrak{f}_{(\mathbf{w}-\mathbf{v})}^{[J]} = 0$ .

Thus,  $\mathbb{W}_J$  is  $\mathbb{V}$ -separating for  $\mathfrak{F}^J$ . By Birkhoff’s Ergodic Theorem, the set  $\mathbb{J}_{(\Gamma; R)}$  of  $J \in \mathbb{N}$  with  $R$  such  $\Gamma$ -gaps is a set of Cesàro density one. Thus, we satisfy the conditions of Proposition 8.  $\square$

To apply Proposition 8 it is clearly sufficient to construct sets  $\mathbb{J}_{(\mathbb{V}; R)}$  for some increasing sequence of numbers  $R_1, R_2, \dots \rightarrow \infty$ , along with a sequence  $\mathbb{V}_1, \mathbb{V}_2, \dots$  so that, for any finite  $\mathbb{V} \subset \mathbb{M}$  we have  $\mathbb{V} \subset \mathbb{V}_k + \mathbf{m}$  for some  $\mathbf{m} \in \mathbb{M}$  and  $k \in \mathbb{N}$ . Also, it suffices to prove that the LCA  $\mathfrak{F}^K$  is diffusive for some power  $K > 0$ : for any  $\chi \in \widehat{\mathcal{A}^{\mathbb{M}}}$ , and any  $k \in [0 \dots K)$ ,  $\chi \circ \mathfrak{F}^k$  is also a character; if  $\mathfrak{F}^K$  is diffusive, then  $\text{rank}[\chi \circ \mathfrak{F}^k \circ \mathfrak{F}^{n-K}] \xrightarrow[n \rightarrow \infty]{\text{dense}} \infty$  for every  $k \in [0 \dots K)$ , which in turn implies that  $\text{rank}[\chi \circ \mathfrak{F}^n] \xrightarrow[n \rightarrow \infty]{\text{dense}} \infty$ .

Proposition 8 can be applied even when the coefficients of  $\mathfrak{F}$  do not commute. For example:

**Example 10:** Let  $\mathcal{A} = (\mathbb{Z}/p)^2$ , and let  $\mathfrak{F} : \mathcal{A}^{\mathbb{Z}} \rightrightarrows \mathcal{A}^{\mathbb{Z}}$  have local map  $\mathfrak{f} : \mathcal{A}^{\{0,1\}} \rightarrow \mathcal{A}$  given:

$$\mathfrak{f} \left( \begin{bmatrix} x_0 \\ y_0 \end{bmatrix}, \begin{bmatrix} x_1 \\ y_1 \end{bmatrix} \right) = (y_0, x_0 + y_1) = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \cdot \begin{bmatrix} x_0 \\ y_0 \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ y_1 \end{bmatrix}$$

$j$	:	...	*	1	0	*	...	*	1	0	*	...	*	0	$q$	1	*	...	*	*	*	...	*	*	*	
$w$	:	...	0	0	0	0	...	0	0	1	0	...	0	0	1	0	0	...	0	0	0	...	0	0	0	
$v$	:	...	0	0	0	0	...	0	0	0	0	...	0	0	0	0	0	...	0	0	*	...	*	*	*	
$j+w$	:	...	*	1	0	*	...	*	1	1	*	...	*	1	0	1	*	...	*	*	*	...	*	*	*	
$j+w-1$	:	...	*	1	0	*	...	*	1	1	*	...	*	1	0	*	*	...	*	*	*	...	*	*	*	
$j+w-v$	:	...	*	1	0	*	...	*	1	1	*	...	*	1	0	*	*	...	*	*	*	...	*	*	*	
$j+w-v-1$	:	...	*	1	0	*	...	*	1	1	*	...	*	1	0	*	*	...	*	*	*	...	*	*	*	
$2w$	:	...	0	0	0	0	...	0	1	0	0	...	0	1	0	0	0	...	0	0	0	...	0	0	0	
$2v$	:	...	0	0	0	0	...	0	0	0	0	...	0	0	0	0	0	...	0	*	*	...	*	*	*	
$2w-2v$	:	...	0	0	0	0	...	0	1	0	0	...	0	0	$q$	$q$	$q$	...	$q$	*	*	...	*	*	*	
$2w-2v-1$	:	...	0	0	0	0	...	0	1	0	0	...	0	0	$q$	$q$	$q$	...	$q$	*	*	...	*	*	*	
Digit #		.....		$i_2$	.....		$i_1$	.....		$i_0$	.....	$L_v$	...	2	1	0										

Figure 1: The  $p$ -ary expansions of  $j$ ,  $w$ ,  $v$ , etc. Here, “\*” represents any digit in in  $[0\dots q]$ .

This invertible LCA was studied in [1], where it was shown to take fully supported Markov measures to Haar measure in the weak\* Cesàro limit. Proposition 3.1 of [1] can be reformulated as:

$$\mathfrak{F}^N = \sum_{m=0}^N \mathfrak{f}_m^{[N]} \sigma^m, \quad \text{where } \mathfrak{f}_m^{[N]} = \begin{bmatrix} \varphi_m^{(N-2)} & \varphi_m^{(N-1)} \\ \varphi_m^{(N-1)} & \varphi_m^{(N)} \end{bmatrix},$$

$$\text{with } \varphi_m^{(N)} = \begin{cases} \left[ \binom{\frac{N+m}{2}}{m} \right]_p & \text{if } m \equiv N \pmod{2} \\ 0 & \text{if } m \not\equiv N \pmod{2} \end{cases}.$$

Thus, if  $m \equiv (N-1) \pmod{2}$ , then the matrix  $\mathfrak{f}_m^{[N]}$  is antidiagonal, and an automorphism iff  $\varphi_m^{(N-1)} = \left[ \binom{\frac{N-1+m}{2}}{m} \right]_p \neq 0$ , which, by Lucas’ Theorem, occurs only when  $m \ll \frac{N-1+m}{2}$ . If  $m \equiv N \equiv (N-2) \pmod{2}$ , then matrix  $\mathfrak{f}_m^{[N]}$  is diagonal, and an automorphism iff  $m \ll \frac{N+m}{2}$  and  $m \ll \frac{N-2+m}{2}$ .

As noted earlier, it suffices to prove that  $\mathfrak{F}^2$  is diffusive. So, fix  $\mathbb{V} = (0 \dots 2V] \subset \mathbb{Z}$  and  $R > 0$ ; we will find a set  $\mathbb{J}_{(\mathbb{V}; R)}$  and, for all  $j \in \mathbb{J}_{(\mathbb{V}; R)}$  some  $\mathbb{W}_j \subset \mathbb{Z}$  with  $\mathcal{C}_{ad}[\mathbb{W}_j] > R$ , so that  $2\mathbb{W}_j$  is  $\mathbb{V}$ -separating for  $\mathfrak{F}^{2j}$ . In other words,  $\forall w \in \mathbb{W}_j$ ,  $\mathfrak{f}_{2w}^{[2j]} \in \mathbf{Aut}[\mathcal{A}]$ , but  $\forall v \in \mathbb{V}$ ,  $\mathfrak{f}_{(2w-v)}^{[2j]} = 0$ . This is equivalent to:

$\forall w \in \mathbb{W}_j$ ,  $\varphi_{2w}^{(2j)} \neq 0 \neq \varphi_{2w}^{(2j-2)}$ , but for all even  $v = 2u \in \mathbb{V}$ ,  $\varphi_{(2w-v)}^{(2j)} = 0 = \varphi_{(2w-v)}^{(2j-2)}$ , and for all odd  $v = 2u + 1 \in \mathbb{V}$ ,  $\varphi_{(2w-v)}^{(2j-1)} = 0$ . This, in turn, is equivalent to:

For all  $w \in \mathbb{W}_j$ ,

$$2w \ll j + w \quad \text{and} \quad 2w \ll j + w - 1, \quad (2)$$

but for all  $u \in (0 \dots V]$ ,

$$\begin{aligned} 2w - 2u &\not\ll j + w - u, & 2w - 2u &\not\ll j + w - u - 1, \\ \text{and } 2w - 2u - 1 &\not\ll j + w - u - 1. \end{aligned} \quad (3)$$

So, let  $q = p - 1$ ,  $L_V = \lceil \log_p(V) \rceil + 1$  and  $L_R = \lceil \log_2(R) \rceil$ , and let  $\mathbb{J}_{(\mathbb{V};R)}$  be the set of all  $j \in \mathbb{N}$  such that  $\mathbb{P}(j)$  contains the word “0q1” somewhere after the first  $L_V$  digits, and contains at least  $L_R$  separate instances of the word “10” after the “0q1”. By Birkhoff’s Ergodic Theorem,  $\mathbb{J}_{(\mathbb{V};R)} \subset \mathbb{N}$  has density 1.

Suppose  $j \in \mathbb{J}_{(\mathbb{V};R)}$ ; and suppose that “0q1” occurs at position  $i_0 > L_v$ , while “10” occurs at positions  $i_{(L_R)} > \dots > i_2 > i_1$ . Let  $w$  to be a number so that  $\mathbb{P}(w)$  contains the word “010” at  $i_0$ , and contains either “01” or “00” at each of  $i_1, i_2, \dots, i_{(L_R)}$ , with zeros everywhere else. Clearly, we can construct  $2^{L_R} > R$  distinct numbers  $w$  of this kind; let  $\mathbb{W}_j$  be the set of all such numbers.

For example, if  $w$  has “01” at  $i_1$  and “00” at  $i_2$ , and  $v \in [0 \dots V]$ , then the  $p$ -ary expansions of the relevant numbers are depicted in Figure 1. By inspection, one can see that equations (2) and (3) are satisfied. Clearly, this will be true for any choice of  $w \in \mathbb{W}_j$  and  $v \in \mathbb{V}$ . 

---

## 5 Harmonic Mixing of Markov Random Fields

**Notation:** Suppose  $\mathbf{a} = \mathcal{A}^{\mathbb{M}}$ , with  $\mathbf{a} = [a_m]_{m \in \mathbb{M}}$ . If  $\mathbb{V} \subset \mathbb{M}$ , then  $\mathbf{a}|_{\mathbb{V}} = [a_v]_{v \in \mathbb{V}} \in \mathcal{A}^{\mathbb{V}}$ . This determines a continuous map  $\mathbf{pr}_{\mathbb{V}} : \mathcal{A}^{\mathbb{M}} \ni \mathbf{a} \mapsto \mathbf{a}|_{\mathbb{V}} \in \mathcal{A}^{\mathbb{V}}$ ; if  $\mu \in \mathcal{M}_{\mathcal{EAS}}[\mathcal{A}^{\mathbb{M}}]$ , then let  $\mathbf{pr}_{\mathbb{V}}^*(\mu)$  be the  $\mathbb{V}$ -marginal projection of  $\mu$  (so that, for any  $\mathcal{U} \subset \mathcal{A}^{\mathbb{V}}$ ,  $\mathbf{pr}_{\mathbb{V}}^*(\mu)[\mathcal{U}] = \mu[\mathcal{U} \times \mathcal{A}^{\mathbb{M} \setminus \mathbb{V}}]$ ).

If  $\mathbf{b} \in \mathcal{A}^{\mathbb{V}}$ , then  $\langle \mathbf{b} \rangle = \left\{ \mathbf{a} \in \mathcal{A}^{\mathbb{M}} ; \mathbf{a}|_{\mathbb{V}} = \mathbf{b} \right\}$  is the associated **cylinder set**, and, if  $\mu \in \mathcal{M}_{\mathcal{EAS}}[\mathcal{A}^{\mathbb{M}}]$ , then  $\mu[\mathbf{b}]$  is the measure of this cylinder set. If  $\mathbb{W} \subset \mathbb{M}$  is disjoint from  $\mathbb{V}$ , and  $\mathbf{c} \in \mathcal{A}^{\mathbb{W}}$ , then  $\mathbf{b\_c} \in \mathcal{A}^{\mathbb{V} \sqcup \mathbb{W}}$  is defined so that  $(\mathbf{b\_c})|_{\mathbb{V}} = \mathbf{b}$  and  $(\mathbf{b\_c})|_{\mathbb{W}} = \mathbf{c}$ ; thus,  $\langle \mathbf{b\_c} \rangle = \langle \mathbf{b} \rangle \cap \langle \mathbf{c} \rangle$ .

Let  $\mathcal{B}(\mathbb{V})$  be the sigma-subalgebra of  $\mathcal{A}^{\mathbb{M}}$  generated by coordinates in  $\mathbb{V}$ ; if  $\phi \in \mathbf{L}^1(\mathcal{A}^{\mathbb{M}}, \mu)$ , let  $\mathbf{E}_{\mathbb{V}}[\phi] \in \mathbf{L}^1(\mathcal{A}^{\mathbb{V}}, \mu)$  be the **conditional expectation** of  $\phi$  given  $\mathcal{B}(\mathbb{V})$ , which we regard as a function on  $\mathcal{A}^{\mathbb{V}}$ . If  $\mathbf{b} \in \mathcal{A}^{\mathbb{V}}$ , then the **conditional probability measure** of  $\mu$ , **given  $\mathbf{b}$** , is the unique measure  $\mu_{\mathbf{b}} \in \mathcal{M}_{\mathcal{EAS}}[\mathcal{A}^{\mathbb{M}}]$  such that  $\langle \phi, \mu_{\mathbf{b}} \rangle = \mathbf{E}_{\mathbb{V}}[\phi](\mathbf{b})$  for every  $\phi \in \mathbf{L}^1(\mathcal{A}^{\mathbb{M}}, \mu)$ . The map  $\mathcal{A}^{\mathbb{V}} \ni \mathbf{b} \mapsto \mu_{\mathbf{b}} \in \mathcal{M}_{\mathcal{EAS}}[\mathcal{A}^{\mathbb{M}}]$  is measurable, and, if  $\mu_{\mathbb{V}} = \mathbf{pr}_{\mathbb{V}}^*(\mu)$  is the marginal projection of  $\mu$  onto  $\mathcal{A}^{\mathbb{V}}$ , then  $\mu$  has the **disintegration** [11, 2]:  $\mu = \int_{\mathcal{A}^{\mathbb{V}}} \mu_{\mathbf{b}} d\mu_{\mathbb{V}}[\mathbf{b}]$ . Note that  $\mathbf{pr}_{\mathbb{V}}(\mu_{\mathbf{b}}) = \delta_{\mathbf{b}}$ , the point mass at  $\mathbf{b}$ . If  $\mathbb{W} \subset \mathbb{M} \setminus \mathbb{V}$  and  $\mathbf{c} \in \mathcal{A}^{\mathbb{W}}$ , we will sometimes write  $\mu_{\mathbf{b}}[\mathbf{c}]$  as “ $\mu[\mathbf{c} \parallel \mathbf{b}]$ ”, or, if  $\mathbf{a} \in \mathcal{A}^{\mathbb{M}}$  is a  $\mu$ -random configuration, as “ $\mu \left[ \frac{\mathbf{a}|_{\mathbb{W}} = \mathbf{c}}{\mathbf{a}|_{\mathbb{V}} = \mathbf{b}} \right]$ ”

## 5.1 Markov Processes

Let  $(\mathbf{X}, \mathcal{X})$  be a measurable space, and let  $\mu \in \mathcal{M}_{\mathcal{EAS}}[\mathbf{X}^{\mathbb{Z}}]$  be a probability measure. Let  $\mathbb{U} = [0..U) \subset \mathbb{Z}$ . If  $n \in \mathbb{Z}$  and  $\mathbf{x} \in \mathbf{X}^{(\mathbb{U}+n)}$ , then  $\langle \mathbf{x} \rangle = \left\{ \mathbf{y} \in \mathbf{X}^{\mathbb{Z}} ; \mathbf{y}|_{(\mathbb{U}+n)} = \mathbf{x} \right\}$ , and  $\mu_{\mathbf{x}}$  is the conditional probability measure of  $\mu$ , given  $\mathbf{x}$ .

$\mu$  is the **path distribution** of a ( $\mathbf{X}$ -valued,  $U$ -step, nonstationary) **Markov process** if, for any  $n \in \mathbb{Z}$  and  $\mathbf{x} \in \mathbf{X}^{(\mathbb{U}+n)}$ , events occurring after time  $n + U$  are independent of those occurring before time  $n$ , relative to  $\mu_{\mathbf{x}}$ : for any  $\mathbb{V}_p \subset (-\infty..n)$ ,  $\mathbb{V}_f \subset [U + n.. \infty)$ , and  $\mathbf{y}_p \in \mathbf{X}^{\mathbb{V}_p}$  and  $\mathbf{y}_f \in \mathbf{X}^{\mathbb{V}_f}$ , we have  $\mu_{\mathbf{x}}[\mathbf{y}_p, \mathbf{y}_f] = \mu_{\mathbf{x}}[\mathbf{y}_p] \cdot \mu_{\mathbf{x}}[\mathbf{y}_f]$ .

Any  $U$ -step Markov process is entirely described by its  $(U+1)$ -dimensional marginals  $\mu_{[n..U+n]} = \mathbf{pr}_{[n..U+n]}^*[\mu]$  for all  $n \in \mathbb{Z}$ , which are called the  $(U$ -step) **transition probabilities** of  $\mu$ . If  $\mathbf{X}$  is finite, then  $\mathcal{M}_{\mathcal{EAS}}[\mathbf{X}; \mathbb{R}] \cong \mathbb{R}^{\mathbf{X}}$ ; if  $U = 1$ , then the transition probabilities  $\{\mu_{\{n, n+1\}}\}_{n \in \mathbb{Z}}$  can be encoded by a sequence of **transition probability matrices**  $\{\mathbf{Q}^{(n)} \in \mathbb{R}^{\mathbf{X} \times \mathbf{X}} ; n \in \mathbb{Z}\}$  and **state distributions**  $\{\eta_n \in \mathbb{R}^{\mathbf{X}} ; n \in \mathbb{Z}\}$  so that, for any  $n \in \mathbb{Z}$ ,  $\eta_{(n+1)} = \mathbf{Q}^{(n)} \cdot \eta_n$ , and, for any  $x_n, x_{(n+1)} \in \mathbf{X}$ ,  $\mu_{\{n, n+1\}}[x_n, x_{(n+1)}] = \mathbf{Q}_{(x_{(n+1)}; x_n)}^{(n)} \cdot \eta_n(x_n)$ .

If  $\mu_{[n..U+n]} = \mu_{[0..U]}$  for all  $n \in \mathbb{Z}$ , then  $\mu$  is **stationary**. If  $\mathbf{X}$  is finite and  $U = 1$ , this means there is some  $\mathbf{Q} \in \mathbb{R}^{\mathbf{X} \times \mathbf{X}}$  and  $\eta \in \mathcal{M}_{\mathcal{EAS}}[\mathbf{X}]$  (with  $\mathbf{Q} \cdot \eta = \eta$ ) so that  $\mathbf{Q}^{(n)} = \mathbf{Q}$  and  $\eta_n = \eta$  for all  $n \in \mathbb{Z}$ . We call  $\eta$  the **stationary state distribution**.

If  $\mathfrak{M} \subset \mathcal{M}_{\mathcal{EAS}}[\mathbf{X}^{[0..n]}]$  is a finite family of transition probabilities, we say  $\mu$  is  **$\mathfrak{M}$ -semistationary** if  $\mu_{[n..U+n]} \in \mathfrak{M}$  for all  $n \in \mathbb{Z}$ . When  $\mathbf{X}$  is



finite and  $U = 1$ , this means that there are some finite families  $\mathfrak{Q}$  and  $\mathfrak{H}$  of transition probability matrices and state distributions, respectively, for  $\mu$  so that, for any  $\eta \in \mathfrak{H}$  and  $\mathbf{Q} \in \mathfrak{Q}$ ,  $\mathbf{Q} \cdot \eta \in \mathfrak{H}$ ; we say  $\mathfrak{Q}$ -semistationary.

If  $\mu$  is  $\mathfrak{M}$ -semistationary, then  $\mu$  has **full support** if every element of  $\mathfrak{M}$  has full support on  $\mathbf{X}^{[0 \dots U]}$ ; as a consequence,  $\mu$  assigns nonzero probability to every finite cylinder set. If  $\mathbf{X}$  is finite and  $U = 1$ , this means that every entry of every transition probability matrix in  $\mathfrak{Q}$  is nonzero.

If  $\mu \in \mathcal{M}_{\text{EAS}}[\mathbf{X}^{\mathbb{Z}}]$  is a Markov process,  $u, w \in \mathbf{X}$ , and  $n \in \mathbb{Z}$ , then the **sandwich measure**  ${}_n\mu_u^w \in \mathcal{M}_{\text{EAS}}[\mathbf{X}]$  is defined so that, if  $\mathbf{x} = [x_n |_{n \in \mathbb{Z}}]$  is a  $\mu$ -random sequence, then for any  $\mathcal{V} \subset \mathbf{X}$ ,  ${}_n\mu_u^w(\mathcal{V}) = \mu \left[ \frac{x_{n+1} \in \mathcal{V}}{(x_n = u) \& (x_{n+2} = w)} \right]$ .

## 5.2 Exponential Harmonic Mixing

If  $\mathcal{A}$  is a finite abelian group, and  $\mu \in \mathcal{M}_{\text{EAS}}[\mathcal{A}^{\mathbb{M}}, \mathbb{C}]$ , we will say  $\mu$  is **exponentially harmonically mixing** with **decay parameter**  $\lambda > 0$  (or “ $\lambda$ -EHM”) if, for all  $\chi \in \widehat{\mathcal{A}^{\mathbb{M}}}$  with  $\text{rank}[\chi] \geq R$ , we have  $|\langle \chi, \mu \rangle| < e^{-\lambda \cdot R}$ . It is straightforward to verify the following

**Lemma 11:** *Suppose  $(\mathbf{X}, \rho)$  is a probability space, and  $\mathbf{X} \ni \mathbf{x} \mapsto \nu_{\mathbf{x}} \in \mathcal{M}_{\text{EAS}}[\mathcal{A}^{\mathbb{M}}; \mathbb{C}]$  is a measurable function so that  $\nu_{\mathbf{x}}$  is  $\lambda$ -EHM for all  $\mathbf{x} \in \mathbf{X}$ . If  $\phi : \mathbf{X} \rightarrow \mathbb{C}$  is measurable and  $\|\phi\|_{\infty} = 1$ , then  $\int_{\mathbf{X}} \phi(x) \cdot \nu_{\mathbf{x}} \, d\rho[\mathbf{x}]$  is also  $\lambda$ -EHM.* □

If  $\mu$  is a stationary, fully supported  $U$ -step Markov measure on  $\mathcal{A}^{\mathbb{Z}}$ , then  $\mu$  is harmonically mixing ( **Part 4** of Proposition 1 in this paper, or Corollary 10 of [5]). The same method easily generalizes to show:

**Proposition 12:** *Suppose  $\mathcal{A}$  is a finite abelian group, and that  $\mathfrak{M} \subset \mathcal{M}_{\text{EAS}}[\mathcal{A}^{[0 \dots n]}]$  is a finite family of fully supported transition probabilities.*

1. *There is a constant  $\lambda > 0$  determined by  $\mathfrak{M}$ , so that, if  $\mu$  is any  $\mathfrak{M}$ -semistationary Markov process on  $\mathcal{A}^{\mathbb{Z}}$ , then  $\mu$  is  $\lambda$ -EHM.*
2. *In particular, if  $\mu$  is a 1-step  $\mathfrak{Q}$ -semistationary Markov process with full support, then  $-\lambda = \frac{1}{2} \cdot \sup_{\substack{\xi, \chi \in \hat{\mathcal{A}} \\ \chi \neq \mathbf{1}}} \sup_{\mathbf{Q}, \mathbf{P} \in \mathfrak{Q}} \log \left\| \xi_{\bullet} \cdot \dagger \mathbf{Q} \cdot \chi_{\bullet} \cdot \dagger \mathbf{P} \right\|_{\infty}$ ,*

*where  $\xi_{\bullet}$  is the diagonal matrix with elements of  $\xi$  along the diagonal (so that, for any  $\phi \in \mathbb{C}^{\mathcal{A}}$ ,  $\xi_{\bullet} \phi$  is the result of multiplying  $\xi$  and  $\phi$  componentwise), and where  $\|\bullet\|_{\infty}$  is the uniform operator norm.*

**Proof:** (Sketch) Proposition 8 in [5] showed that a stationary 1-step Markov matrix was harmonically mixing; in fact, the proof showed that  $|\langle \chi, \mu \rangle| < e^{-\lambda R}$  for all  $\chi \in \widehat{\mathcal{A}^{\mathbb{Z}}}$  with  $\text{rank}[\chi] = R$ , where  $-\lambda := \frac{1}{2} \cdot \sup_{\substack{\xi, \chi \in \widehat{\mathcal{A}} \\ \chi \neq \mathbf{1}}} \log \left\| \xi \cdot \dagger \mathbf{Q} \cdot \chi \cdot \dagger \mathbf{Q} \right\|_{\infty}$ . The same argument works for a semistationary 1-step process; this yields **Part 2**.

The proof of Corollary 10 in [5] showed how any fully supported  $U$ -step process could be “recoded” as a fully supported 1-step process; harmonic mixing of the latter implied harmonic mixing of the former. Corollary 10 thus followed from Proposition 8. By an identical argument **Part 1** follows from **Part 2**. □

### 5.3 Markov Random Fields

Let  $\mathbb{U} \subset \mathbb{M}$  be a finite “neighbourhood of 0” (e.g.  $\mathbb{M} = \mathbb{Z}^D$  and  $\mathbb{U} = [-1\dots 1]^D$ ). For any subset  $\mathbb{V} \subset \mathbb{M}$ , let  $cl(\mathbb{V}) := \mathbb{V} + \mathbb{U}$ , and let  $\partial(\mathbb{V}) := cl(\mathbb{V}) \setminus \mathbb{V}$  (see Figure 2).




Figure 2:  $\mathbb{U}$ ,  $\mathbb{V}$ , and  $\partial\mathbb{V}$

$\mu \in \mathcal{M}_{EAS}[\mathcal{A}^{\mathbb{M}}]$  is a **(nonstationary) Markov random field** [10] with **interaction range**  $\mathbb{U}$  (or “ $\mathbb{U}$ -MRF”) if, for any  $\mathbb{W} \subset \mathbb{M}$ , and any  $\mathbf{a} \in \mathcal{A}^{\partial(\mathbb{W})}$ , events occurring “inside”  $\mathbb{W}$  are independent of those occurring “outside”, relative to the conditional measure  $\mu_{\mathbf{a}}$ . In other words, for any  $\mathbb{V}_{in} \subset \mathbb{W}$ ,  $\mathbb{V}_{out} \subset \mathbb{M} \setminus cl(\mathbb{W})$ , and  $\mathbf{b}_{in} \in \mathcal{A}^{\mathbb{V}_{in}}$ ,  $\mathbf{b}_{out} \in \mathcal{A}^{\mathbb{V}_{out}}$ , we have:  $\mu_{\mathbf{a}}[\mathbf{b}_{in}, \mathbf{b}_{out}] = \mu_{\mathbf{a}}[\mathbf{b}_{in}] \cdot \mu_{\mathbf{a}}[\mathbf{b}_{out}]$ .

For example, if  $\mathbb{M} = \mathbb{Z}$ , then the  $U$ -step Markov processes on  $\mathcal{A}^{\mathbb{M}}$  are exactly the Markov random fields with interaction range  $\mathbb{U} = (-U\dots U)$ .

$\mu$  is **stationary** if it is invariant under translation by  $\mathbb{M}$ . In this case,  $\mu_{(\mathbb{U}+m)} = \mu_{\mathbb{U}}$  for every  $m \in \mathbb{M}$ , and  $\mu_{\mathbb{U}} = \mathbf{pr}_{\mathbb{U}}^*(\mu)$  is called the **local interaction** for  $\mu$ .

If  $\mathfrak{J} \subset \mathcal{M}_{\mathcal{EAS}}[\mathcal{A}^{\mathbb{U}}]$  is finite, then  $\mu$  is  **$\mathfrak{J}$ -semistationary** if  $\mu_{(\mathbb{U}+m)} \in \mathfrak{J}$  for every  $m \in \mathbb{M}$ .  $\mathfrak{J}$  is called the set of **local interactions**. We say  $\mu$  has **full support** if all elements of  $\mathfrak{J}$  have full support on  $\mathcal{A}^{\mathbb{U}}$ .

**Lamination Processes:** Suppose  $\tilde{\mathbb{U}} \subset \tilde{\mathbb{M}} = \mathbb{M} \times \mathbb{Z}$  and  $\mu \in \mathcal{M}_{\mathcal{EAS}}[\mathcal{A}^{\tilde{\mathbb{M}}}]$  is a  $\tilde{\mathbb{U}}$ -MRF. By a suitable recoding, we can assume  $\tilde{\mathbb{U}} = \mathbb{U} \times \{-1, 0, 1\}$  for some  $\mathbb{U} \subset \mathbb{M}$ . We can then realize  $\mu$  via an  $\mathcal{A}^{\mathbb{M}}$ -valued, 1-step Markov process, called the **lamination process**. Intuitively, we imagine this Markov process as constructing a  $\mu$ -random configuration in  $\mathcal{A}^{\tilde{\mathbb{M}}}$  by laying down successive random “ $\mathbb{M}$ -layers”, with each  $\mathbb{M}$ -layer conditional on the previous one.




Figure 3:  $\tilde{\mathbb{M}} = \mathbb{M} \times \mathbb{Z}$  and  $\tilde{\mathbb{U}} = \mathbb{U} \times \{-1, 0, 1\}$ ;  $\tilde{\mathbb{V}}_k$  and  $\partial \tilde{\mathbb{V}}_k$

To see that this is a Markov process on  $\mathcal{A}^{\mathbb{M}}$ , fix  $k$ , and let  $\tilde{\mathbb{V}}_k = \mathbb{M} \times (-\infty\dots k)$  (the “past”). Then  $\partial(\tilde{\mathbb{V}}_k) = \mathbb{M} \times \{k\}$  (the “present”) and  $\tilde{\mathbb{M}} \setminus \text{cl}(\tilde{\mathbb{V}}_k) = \mathbb{M} \times (k\dots\infty)$  (the “future”); the Markov field condition of  $\mu$  implies that events in the past are independent of those in the future, given complete information about the present (see Figure 3). The original field measure  $\mu \in \mathcal{M}_{\mathcal{EAS}}[\mathcal{A}^{\mathbb{M} \times \mathbb{Z}}]$  is also the path distribution (as a measure on  $(\mathcal{A}^{\mathbb{M}})^{\mathbb{Z}}$ ) for the lamination process.

**Sandwich Measures:** Again assume  $\tilde{\mathbb{M}} = \mathbb{M} \times \mathbb{Z}$  and  $\tilde{\mathbb{U}} = \mathbb{U} \times \{-1, 0, 1\}$ . If  $\mathbf{a} \in \mathcal{A}^{\mathbb{M} \times \{k-1\}}$  and  $\mathbf{c} \in \mathcal{A}^{\mathbb{M} \times \{k+1\}}$  (see Figure 4), then the **sandwich measure** determined by  $\mathbf{a}$  and  $\mathbf{c}$  is the sandwich measure  ${}_{(k-1)}\mu_{\mathbf{a}}^{\mathbf{c}} \in \mathcal{M}_{\text{EAS}}[\mathcal{A}^{\mathbb{M}}]$  of the lamination process; since  $k$  is implicit in the definition of  $\mathbf{a}$  and  $\mathbf{c}$ , we will suppress it, and denote the sandwich measure as “ $\mu_{\mathbf{a}}^{\mathbf{c}}$ ”. In other words,  $\mu_{\mathbf{a}}^{\mathbf{c}}$  is the conditional measure  $\mu_{\mathbf{a}, \mathbf{c}}$ , projected onto  $\mathcal{A}^{\mathbb{M} \times \{k\}}$ . The following is easy to verify:




Figure 4:

**Lemma 13:**

1.  $\mu_{\mathbf{a}}^{\mathbf{c}}$  is a Markov random field on  $\mathcal{A}^{\mathbb{M}}$ , with interaction range  $\mathbb{U}$ .
2. If  $\tilde{\mathcal{J}} \subset \mathcal{M}_{\text{EAS}}[\mathcal{A}^{\tilde{\mathbb{U}}}]$  and  $\mu$  is  $\tilde{\mathcal{J}}$ -semistationary, then there is some finite  $\mathcal{J} \subset \mathcal{M}_{\text{EAS}}[\mathcal{A}^{\mathbb{U}}]$  so that all sandwich measures of  $\mu$  are  $\mathcal{J}$ -semistationary.
3. If  $\mu$  has full support, then so does every sandwich measure of  $\mu$ .  $\square$

The harmonic mixing of an MRF depends on the the harmonic mixing of its sandwich measures:

**Proposition 14:** If  $\mathcal{A}$  is a finite abelian group, and  $\mu$  is a semistationary MRF on  $\mathcal{A}^{\mathbb{M} \times \mathbb{Z}}$  and all sandwich measures of  $\mu$  are  $\lambda$ -EHM, then  $\mu$  is  $\lambda'$ -EHM, where  $\lambda' = \lambda/2$ .

**Proof:** See §5.5.  $\square$

From this follows our main result:

**Theorem 15:** Suppose  $\mathcal{A}$  is a finite abelian group,  $\mathbb{U} \subset \mathbb{Z}^D$ , and let  $\tilde{\mathcal{J}} \subset \mathcal{M}_{\mathcal{EAS}}[\mathcal{A}^{\mathbb{U}}]$  be a finite set of local interactions with full support. Then  $\exists \lambda > 0$  so that if  $\mu$  is any  $\tilde{\mathcal{J}}$ -semistationary MRF on  $\mathcal{A}^{\mathbb{Z}^D}$ , then  $\mu$  is  $\lambda$ -EHM.

**Proof:** (by induction on  $D$ ) If  $D = 1$ , this is just Proposition 12.

Suppose inductively that the claim is true for MRFs on  $\mathbb{Z}^{D-1}$ , and let  $\mu \in \mathcal{A}^{\mathbb{Z}^D}$ . By Lemma 13, all sandwich measures of  $\mu$  are  $\mathcal{J}$ -semistationary MRFs on  $\mathcal{A}^{\mathbb{Z}^{D-1}}$ , where  $\mathcal{J}$  is some finite set of local interactions with full support. Thus, by induction hypothesis, all these sandwich measures are  $\lambda$ -EHM for some  $\lambda > 0$ . Thus, by Proposition 14,  $\mu$  is  $\lambda'$ -EHM, with  $\lambda' = -\lambda/2$ . □

## 5.4 Markov Operators

When  $\mathbf{X}$  is finite, a 1-step  $\mathbf{X}$ -valued Markov process can be defined by a series of with transition probability matrices  $\{\mathbf{Q}^{(n)}\}_{n \in \mathbb{Z}}$ . These matrices define linear operators on the space  $\mathcal{M}_{\mathcal{EAS}}[\mathbf{X}; \mathbb{R}] \cong \mathbb{R}^{\mathbf{X}}$ , so that, if  $\eta_n \in \mathcal{M}_{\mathcal{EAS}}[\mathbf{X}]$  is the state distribution at time  $n$ , then  $\mathbf{Q}^{(n)} \cdot \eta_n = \eta_{n+1}$  is the state distribution at time  $n + 1$ .

When  $\mathbf{X}$  is an arbitrary measurable space (with sigma-algebra  $\mathcal{X}$ ), transition probabilities are described by linear operators on the vector space  $\mathcal{M}_{\mathcal{EAS}}[\mathbf{X}; \mathbb{R}]$  (which, for technical reasons, we will treat as linear operators on  $\mathcal{M}_{\mathcal{EAS}}[\mathbf{X}; \mathbb{C}]$ ).

**Idea:** Informally speaking, a **Markov operator** is linear operator  $\mathbf{Q} : \mathcal{M}_{\mathcal{EAS}}[\mathbf{X}; \mathbb{C}] \rightarrow \mathcal{M}_{\mathcal{EAS}}[\mathbf{X}; \mathbb{C}]$  mapping the set  $\mathcal{M}_{\mathcal{EAS}}[\mathbf{X}]$  of probability measures into itself. Suppose  $(y_0, y_1) \in \mathbf{X}^{\{0,1\}}$  is a random couple, and  $\mathbf{Q}$  is the transition probability operator from time 0 to time 1. If  $x \in \mathbf{X}$ , and  $\delta_x \in \mathcal{M}_{\mathcal{EAS}}[\mathbf{X}]$  is the point mass at  $x$ , then the probability measure  $\mathbf{q}_x := \mathbf{Q}(\delta_x)$  is the conditional state distribution of  $y_1$  given that  $y_0 = x$ : for all  $\mathcal{U} \subset \mathbf{X}$ ,  $\mathbf{q}_x[\mathcal{U}] = \mathbf{P}_{nb} \left[ \frac{y_1 \in \mathcal{U}}{y_0 = x} \right]$ . When  $\mathbf{X}$  is finite, measures on  $\mathbf{X}$  are vectors and  $\mathbf{Q}$  is a matrix, and  $\mathbf{q}_x$  is just the  $x$ th column of this matrix.

Suppose  $y_0, y_1$  have distributions  $\eta_0, \eta_1 \in \mathcal{M}_{\mathcal{EAS}}[\mathbf{X}]$  respectively, with  $\eta_1 = \mathbf{Q}(\eta_0)$ . If  $\phi : \mathbf{X} \rightarrow \mathbb{C}$  is a measurable function, then the expected value of  $\phi(y_1)$  is given by  $\langle \phi, \eta_1 \rangle = \langle \phi, \mathbf{Q}(\eta_0) \rangle = \langle \dagger \mathbf{Q}(\phi), \eta_0 \rangle$ , where  $\dagger \mathbf{Q}$  is the **adjoint** of  $\mathbf{Q}$ .

For any measurable  $\mathcal{U} \subset \mathbf{X}$ , let  $\dagger \mathbf{q}_{\mathcal{U}} := \dagger \mathbf{Q}(\mathbf{1}_{\mathcal{U}})$ . Thus, for any  $x \in \mathbf{X}$ ,  $\dagger \mathbf{q}_{\mathcal{U}}(x) = \dagger \mathbf{Q}(\mathbf{1}_{\mathcal{U}})(x) = \langle \dagger \mathbf{Q}(\mathbf{1}_{\mathcal{U}}), \delta_x \rangle = \langle \mathbf{1}_{\mathcal{U}}, \mathbf{Q}(\delta_x) \rangle = \langle \mathbf{1}_{\mathcal{U}}, \mathbf{q}_x \rangle =$

$\mathbf{q}_x[\mathcal{U}]$ . When  $\mathbf{X}$  is finite and  $\mathbf{Q}$  is a matrix and  $\mathcal{U} = \{u\}$  is a singleton set, then  $\dagger\mathbf{q}_{\mathcal{U}}$  is just the  $u$ th row of  $\mathbf{Q}$  (or the  $u$ th column of  $\dagger\mathbf{Q}$ ).

We need to develop some technology to make these ideas well-defined.

**Formalism:** If  $\Phi : \mathbf{X} \rightarrow \mathbb{C}$  is measurable, then let  $\|\Phi\|_{\infty} = \sup_{x \in \mathbf{X}} |\Phi(x)|$ , and consider the Banach space  $\mathcal{M}_{\infty}(\mathbf{X}, \mathcal{X}) = \{\Phi : \mathbf{X} \rightarrow \mathbb{C}; \Phi \text{ measurable, } \|\Phi\|_{\infty} < \infty\}$  and its unit ball,  $\mathcal{B}_1 = \{\Phi \in \mathcal{M}_{\infty}; \|\Phi\|_{\infty} \leq 1\}$ . Now,  $\mathcal{M}_{\mathcal{EAS}}[\mathbf{X}; \mathbb{C}]$  embeds into the dual space  $\mathcal{M}_{\infty}^*$  in a natural way; endow it with the appropriate weak\* topology. The following results are straightforward:

**Lemma 16:** *The simple functions of the form  $\Phi = \sum_n \phi_n \mathbb{1}_{\mathcal{U}_n}$  are dense in  $\mathcal{M}_{\infty}$  (where  $\phi_n \in \mathbb{C}$  and  $\mathcal{U}_n \subset \mathbf{X}$  are measurable).*

*The weak\* topology on  $\mathcal{M}_{\mathcal{EAS}}[\mathbf{X}; \mathbb{C}]$  is determined by convergence on measurable sets. Thus, a sequence  $\{\mu_n\}_{n=1}^{\infty}$  converges to  $\mu \in \mathcal{M}_{\mathcal{EAS}}[\mathbf{X}; \mathbb{C}]$  if and only if  $\mu_n[\mathcal{U}] \xrightarrow{n \rightarrow \infty} \mu[\mathcal{U}]$  for all measurable  $\mathcal{U} \subset \mathbf{X}$ .  $\square$*

If a function  $\mathbf{X} \ni x \mapsto \mu_x \in \mathcal{M}_{\mathcal{EAS}}[\mathbf{X}; \mathbb{C}]$  is measurable relative to the weak\* Borel algebra of  $\mathcal{M}_{\mathcal{EAS}}[\mathbf{X}; \mathbb{C}]$ , and  $\nu$  is some other measure on  $\mathbf{X}$ , then  $\mu = \int_{\mathbf{X}} \mu_x d\nu[x]$  is the measure so that, for all  $\mathcal{U} \subset \mathbf{X}$ ,  $\mu[\mathcal{U}] = \int_{\mathbf{X}} \mu_x[\mathcal{U}] d\nu[x]$ ; by Lemma 16, this well-defines the action of  $\mu$  on  $\mathcal{M}_{\infty}$ .

If  $\mathbf{Q} : \mathcal{M}_{\mathcal{EAS}}[\mathbf{X}; \mathbb{C}] \xrightarrow{\leftarrow} \mathbb{C}$ , then define  $\|\mathbf{Q}\| := \sup_{x \in \mathbf{X}} \|\mathbf{q}_x\|_{var}$  (note: this is *not* the operator norm of  $\mathbf{Q}$ ). Say that  $\mathbf{Q}$  is **smooth** if  $\mathbf{Q}$  is linear, measurable relative to the weak\* Borel sigma algebra, and  $\|\mathbf{Q}\| < \infty$ .

**Lemma 17:**

1. *If  $\mathbf{Q}$  is smooth, then its adjoint  $\dagger\mathbf{Q} : \mathcal{M}_{\infty} \xrightarrow{\leftarrow} \mathbb{C}$  is a well-defined, bounded linear operator, and  $\|\dagger\mathbf{Q}\|_{\infty} \leq \|\mathbf{Q}\|$ .*
2. *If  $\mathbf{X} \ni x \mapsto \mathbf{q}_x \in \mathcal{M}_{\mathcal{EAS}}[\mathbf{X}; \mathbb{C}]$  is a measurable function and  $M = \sup_{x \in \mathbf{X}} \|\mathbf{q}_x\|_{var} < \infty$ , then the function  $\mathbf{Q} : \mathcal{M}_{\mathcal{EAS}}[\mathbf{X}; \mathbb{C}] \xrightarrow{\leftarrow} \mathbb{C}$  defined:  $\mathbf{Q}(\mu) = \int_{\mathbf{X}} \mathbf{q}_x d\mu[x]$  is smooth and continuous, and  $\|\mathbf{Q}\| = M$ .*

**Proof:**

**Proof of Part 1:** For any  $\phi \in \mathcal{M}_{\infty}$ , and any  $x \in \mathbf{X}$ , define  $(\dagger\mathbf{Q}\phi)(x) = \langle \phi, \mathbf{q}_x \rangle$ . Then  $\dagger\mathbf{Q}(\phi)$  is measurable (the function  $\mathbf{X} \ni x \mapsto \delta_x \in \mathcal{M}_{\mathcal{EAS}}[\mathbf{X}; \mathbb{C}]$  is measurable; hence, so is the function  $(x \mapsto \mathbf{q}_x)$ ; thus, so is  $\dagger\mathbf{Q}(\phi)$ ). Also,  $\left\| \dagger\mathbf{Q}(\phi) \right\|_{\infty} \leq \|\phi\|_{\infty} \cdot \sup_{x \in \mathbf{X}} \|\mathbf{q}_x\|_{var}$ .

**Proof of Part 2:** Clearly,  $\mathbf{Q}$  is well-defined and linear, and  $\|\mathbf{Q}\| = M$ .

To see that  $\mathbf{Q}$  is continuous, let  $\mathcal{U} \subset \mathbf{X}$ ; then  $\mathbf{Q}(\mu)[\mathcal{U}] = \int_{\mathbf{X}} \mathbf{q}_x[\mathcal{U}] d\mu[x]$ . The function  $\mathbf{X} \ni x \mapsto \mathbf{q}_x[\mathcal{U}] \in \mathbb{C}$  is measurable; thus, if  $\mu_n \xrightarrow{n \rightarrow \infty} \mu$  in the weak\* topology, then  $\mathbf{Q}(\mu_n)[\mathcal{U}] = \int_{\mathbf{X}} \mathbf{q}_x[\mathcal{U}] d\mu_n[x] \xrightarrow{n \rightarrow \infty} \int_{\mathbf{X}} \mathbf{q}_x[\mathcal{U}] d\mu[x] = \mathbf{Q}(\mu)[\mathcal{U}]$ .  $\square$

We define a **Markov operator** to be a smooth linear operator on  $\mathcal{M}_{\text{EAS}}[\mathbf{X}; \mathbb{C}]$  that maps  $\mathcal{M}_{\text{EAS}}[\mathbf{X}]$  into itself. By Lemma 17, it suffices to define a measurable collection  $(x \mapsto \mathbf{q}_x)$  of transition probability measures. We will be concerned with the following case:

**Example 18:** Suppose  $\tilde{\mathbb{M}} = \mathbb{M} \times \mathbb{Z}$  and  $\mu \in \mathcal{M}_{\text{EAS}}[\mathcal{A}^{\tilde{\mathbb{M}}}]$  is an MRF and consider the **lamination process**; we claim the transition probabilities are determined by a sequence  $\{\mathbf{Q}^{(n)}\}_{n \in \mathbb{Z}}$  of Markov operators.

Let  $\mathbb{M}_k := \mathbb{M} \times \{k\}$  for  $k = n$  or  $n + 1$ . If  $\mathbf{c} \in \mathcal{A}^{\tilde{\mathbb{M}}}$  is a  $\mu$ -random configuration, then for each  $\mathbf{a} \in \mathcal{A}^{\mathbb{M}}$ ,  $\mathbf{q}_{\mathbf{a}}^{(n)}$  is the conditional distribution of  $\mathbf{c}|_{\mathbb{M}_{(n+1)}}$  given that  $\mathbf{c}|_{\mathbb{M}_n} = \mathbf{a}$ . Formally, if  $\mu_{\mathbf{a}} \in \mathcal{M}_{\text{EAS}}[\mathcal{A}^{\mathbb{M}}]$  is the conditional distribution given  $\mathbf{a}$ , then  $\mathbf{q}_{\mathbf{a}}^{(n)} = \mathbf{pr}_{\mathbb{M}_{(n+1)}}^*(\mu_{\mathbf{a}})$ . The map  $\mathcal{A}^{\mathbb{M}} \ni \mathbf{a} \mapsto \mathbf{q}_{\mathbf{a}}^{(n)} \in \mathcal{M}_{\text{EAS}}[\mathcal{A}^{\mathbb{M}}]$  is measurable because the map  $\mathcal{A}^{\mathbb{M}_n} \ni \mathbf{a} \mapsto \mu_{\mathbf{a}} \in \mathcal{M}_{\text{EAS}}[\mathcal{A}^{\mathbb{M}}]$  is measurable [11], while  $\mathbf{pr}_{\mathbb{M}_{(n+1)}} : \mathcal{M}_{\text{EAS}}[\mathcal{A}^{\mathbb{M}}] \rightarrow \mathcal{M}_{\text{EAS}}[\mathcal{A}^{\mathbb{M}_{(n+1)}}]$  is continuous.  $\square$

If  $\chi \in \mathcal{M}_{\infty}$ , then let  $\chi_{\bullet} : \mathcal{M}_{\infty} \leftarrow$  be the bounded linear operator induced by multiplication with  $\chi$ : for any  $\phi \in \mathcal{M}_{\infty}$  and  $x \in \mathbf{X}$ ,  $(\chi_{\bullet}\phi)(x) = \chi(x) \cdot \phi(x)$ . To establish that Markov processes on  $\mathcal{A}^{\mathbb{Z}}$  were EHM (Proposition 12), we bounded the norm of operators of the form  $\xi_{\bullet} \circ \mathbf{Q} \circ \chi_{\bullet} \circ \mathbf{P}$ , where  $\xi, \chi \in \widehat{\mathcal{A}^{\mathbb{M}}}$ . We will employ a similar strategy to show that MRFs are EHM; this will require the following result:

**Lemma 19:** Let  $\mathbf{Q}, \mathbf{P} : \mathcal{M}_{\text{EAS}}[\mathbf{X}] \leftarrow$  be Markov operators. For any  $\phi \in \mathcal{M}_{\infty}$ , define  $\mu_x^{\phi} \in \mathcal{M}_{\text{EAS}}[\mathbf{X}; \mathbb{C}]$  by:  $d\mu_x^{\phi} = \dagger\mathbf{P}(\phi) \cdot d\mathbf{q}_x$ .

1. For any  $\chi \in \mathcal{M}_{\infty}$ ,  $\left\| \dagger\mathbf{Q} \circ \chi_{\bullet} \circ \dagger\mathbf{P} \right\|_{\infty} = \sup_{\phi \in \mathcal{B}_1} \sup_{x \in \mathbf{X}} \left| \left\langle \chi, \mu_x^{\phi} \right\rangle \right|$ .

2. Suppose that  $\mu \in \mathcal{M}_{\text{EAS}}[\mathbf{X}^{\mathbb{Z}}]$  is a Markov process and  $\mathbf{Q}$  and  $\mathbf{P}$  are the transition probability operators at time 0 and 1, respectively. For any  $u, w \in \mathbf{X}$ , let  $\mu_u = \mathbf{P} \circ \mathbf{Q}(\delta_u)$  be the conditional probability measure on  $\mathbf{X}$  at time 2 induced by state  $u \in \mathbf{X}$  at time 0, and let  $\mu_u^w$  be the sandwich

measure on  $\mathbf{X}$  induced by  $u \in \mathbf{X}$  at time 0 and  $w \in \mathbf{X}$  at time 2. Then for any  $\Phi \in \mathcal{M}_\infty$ ,  $\mu_u^\Phi = \int_{\mathbf{X}} \Phi(w) \cdot \mu_u^w d\mu_u[w]$ .

**Proof:**

**Proof of Part 1:** For any  $\phi \in \mathcal{B}_1$  and  $x \in \mathbf{X}$ ,  $\dagger\mathbf{Q} \circ \chi_\bullet \circ \dagger\mathbf{P}(\phi)(x) = \langle \dagger\mathbf{Q} \circ \chi_\bullet \circ \dagger\mathbf{P}(\phi), \delta_x \rangle = \langle \chi_\bullet \circ \dagger\mathbf{P}(\phi), \mathbf{Q}(\delta_x) \rangle = \langle \chi \cdot \dagger\mathbf{P}(\phi), \mathbf{q}_x \rangle = \langle \chi, \mu_x^\phi \rangle$ . Thus,  $\left\| \dagger\mathbf{Q} \circ \chi_\bullet \circ \dagger\mathbf{P} \right\|_\infty = \sup_{\phi \in \mathcal{B}_1} \left\| \dagger\mathbf{Q} \circ \chi_\bullet \circ \dagger\mathbf{P}(\phi) \right\|_\infty = \sup_{\phi \in \mathcal{B}_1} \sup_{x \in \mathbf{X}} \left| \dagger\mathbf{Q} \circ \chi_\bullet \circ \dagger\mathbf{P}(\phi)(x) \right| = \sup_{\phi \in \mathcal{B}_1} \sup_{x \in \mathbf{X}} \left| \langle \chi, \mu_x^\phi \rangle \right|$ .

**Proof of Part 2:** We want to show that for any  $\mathbf{V} \subset \mathbf{X}$ ,  $\mu_u^\Phi(\mathbf{V}) = \int_{\mathbf{X}} \Phi(w) \cdot \mu_u^w(\mathbf{V}) d\mu_u[w]$ . First, suppose that  $\mathbf{W} \subset \mathbf{X}$  and  $\Phi = \mathbf{1}_\mathbf{W}$ . Let  $\mu_x^{\mathbf{X}\mathbf{W}} := \mu_x^{\mathbf{1}\mathbf{W}}$ ; thus,  $d\mu_x^{\mathbf{W}} = \dagger\mathbf{p}_\mathbf{W} d\mathbf{q}_x$ . Then:

$$\begin{aligned} \mu_u^{\mathbf{W}}(\mathbf{V}) &= \int_{\mathbf{V}} d\mu_u^{\mathbf{W}}[v] = \int_{\mathbf{V}} \dagger\mathbf{p}_\mathbf{W}(v) d\mathbf{q}_u[v] \\ &= \int_{\mathbf{V}} \mathbf{p}_v[\mathbf{W}] d\mathbf{q}_u[v] = \int_{\mathbf{V}} \mu \left[ \frac{x_2 \in \mathbf{W}}{x_1 = v} \right] d\mathbf{q}_u[v] \\ &= \int_{\mathbf{V}} \mu \left[ \frac{x_2 \in \mathbf{W}}{(x_1 = v) \& (x_0 = u)} \right] d\mathbf{q}_u[v] \quad (\text{by the Markov property}) \\ &\stackrel{(1)}{=} \int_{\mathbf{W}} \mu_u^w(\mathbf{V}) d\mu_u[w] = \int_{\mathbf{X}} \mathbf{1}_\mathbf{W}(w) \cdot \mu_u^w(\mathbf{V}) d\mu_u[w]. \end{aligned}$$

To see (1), let  $\mathcal{X}_k$  be the sigma-subalgebra of  $\mathbf{X}^\mathbb{Z}$  generated by coordinate  $x_k$ , and let  $\mathbf{E}_k[\bullet]$  (resp.  $\mathbf{E}_{k,j}[\bullet]$ ) be the conditional expectation with respect to  $\mathcal{X}_k$  (resp.  $\mathcal{X}_k \vee \mathcal{X}_j$ ). Let  $\mathbf{W}_2 = \{\mathbf{x} \in \mathbf{X}^\mathbb{Z}; x_2 \in \mathbf{W}\}$  and  $\mathbf{V}_1 = \{\mathbf{x} \in \mathbf{X}^\mathbb{Z}; x_1 \in \mathbf{V}\}$ . Then

$$\begin{aligned} \int_{\mathbf{V}} \mu \left[ \frac{x_2 \in \mathbf{W}}{(x_1 = v) \& (x_0 = u)} \right] d\mathbf{q}_u[v] &= \int_{\mathbf{V}} \mathbf{E}_{0,1}[\mathbf{1}_{\mathbf{W}_2}](u, v) d\mathbf{q}_u[v] \\ &= \int_{\mathbf{X}} \mathbf{1}_\mathbf{V} \cdot \mathbf{E}_{0,1}[\mathbf{1}_{\mathbf{W}_2}](u, v) d\mathbf{q}_u[v] = \mathbf{E}_0[\mathbf{1}_{\mathbf{V}_1} \cdot \mathbf{E}_{0,1}[\mathbf{1}_{\mathbf{W}_2}]](u) \\ &= \mathbf{E}_0[\mathbf{E}_{0,1}[\mathbf{1}_{\mathbf{V}_1} \cdot \mathbf{1}_{\mathbf{W}_2}]](u) = \mathbf{E}_0[\mathbf{1}_{\mathbf{V}_1} \cdot \mathbf{1}_{\mathbf{W}_2}](u) \\ &= \mathbf{E}_0[\mathbf{E}_{0,2}[\mathbf{1}_{\mathbf{V}_1} \cdot \mathbf{1}_{\mathbf{W}_2}]](u) = \mathbf{E}_0[\mathbf{1}_{\mathbf{W}_2} \cdot \mathbf{E}_{0,2}[\mathbf{1}_{\mathbf{V}_1}]](u) \\ &= \int_{\mathbf{W}} \mu \left[ \frac{x_1 \in \mathbf{V}}{(x_2 = w) \& (x_0 = u)} \right] d\mu_u[w] = \int_{\mathbf{W}} \mu_u^w(\mathbf{V}) d\mu_u[w]. \end{aligned}$$



Next, if  $\Phi = \sum_n \phi_n \mathbf{1}_{\mathbf{W}_n}$  is a simple function, then  $\dagger\mathbf{P}(\Phi) = \sum_n \phi_n \dagger\mathbf{P}_{\mathbf{W}_n}$ , so that  $d\mu_x^\Phi = \dagger\mathbf{P}(\Phi) \cdot d\mathbf{q}_x = \sum_n \phi_n \dagger\mathbf{P}_{\mathbf{W}_n} d\mathbf{q}_x = \sum_n \phi_n d\mu_x^{\mathbf{W}_n}$ . Thus,  $\mu_x^\Phi(\mathbf{V}) = \int_{\mathbf{V}} d\mu_x^\Phi = \sum_n \phi_n \cdot \int_{\mathbf{V}} d\mu_x^{\mathbf{W}_n} = \sum_n \phi_n \cdot \mu_x^{\mathbf{W}_n}[\mathbf{V}] = \sum_n \phi_n \cdot \int_{\mathbf{X}} \mathbf{1}_{\mathbf{W}_n}(w) \mu_u^w(\mathbf{V}) d\mu_u[w] = \int_{\mathbf{X}} \left( \sum_n \phi_n \mathbf{1}_{\mathbf{W}_n}(w) \right) \cdot \mu_u^w(\mathbf{V}) d\mu_u[w] = \int_{\mathbf{X}} \Phi(w) \cdot \mu_u^w(\mathbf{V}) d\mu_u[w]$ , as desired.

Finally, if  $\{\Phi_n\}_{n=1}^\infty$  is a bounded sequence of simple functions so that  $\Phi_n \xrightarrow{n \rightarrow \infty} \Phi$  in  $\mathcal{M}_\infty$ , then  $\dagger\mathbf{P}(\Phi_n) \xrightarrow{n \rightarrow \infty} \dagger\mathbf{P}(\Phi)$  in  $\mathcal{M}_\infty$ , so that  $\mu_x^{\Phi_n} \xrightarrow{n \rightarrow \infty} \mu_x^\Phi$  in the weak\* topology on  $\mathcal{M}_{\text{EAS}}[\mathbf{X}]$ . But by dominated convergence, we also know that  $\mu_x^{\Phi_n}(\mathbf{V}) = \int_{\mathbf{X}} \Phi_n(w) \cdot \mu_u^w(\mathbf{V}) d\mu_u[w] \xrightarrow{n \rightarrow \infty} \int_{\mathbf{X}} \Phi(w) \cdot \mu_u^w(\mathbf{V}) d\mu_u[w]$ ; hence,  $\mu_x^\Phi(\mathbf{V}) = \int_{\mathbf{X}} \Phi(w) \cdot \mu_u^w(\mathbf{V}) d\mu_u[w]$ , for all  $\mathbf{V} \subset \mathbf{X}$ , as desired.  $\square$

## 5.5 Uniform Harmonic Mixing

Now, let  $\mathbf{X} = \mathcal{A}^{\mathbb{M}}$ , and consider a 1-step Markov process on  $\mathcal{A}^{\mathbb{M}}$  determined by a sequence of Markov operators  $\{\mathbf{Q}^{(n)}; n \in \mathbb{Z}\}$ . For every  $n \in \mathbb{Z}$ ,  $\mathbf{a} \in \mathcal{A}^{\mathbb{M}}$  and  $\phi \in \mathcal{M}_\infty$ , define  ${}_n\mu_{\mathbf{a}}^\phi \in \mathcal{M}_{\text{EAS}}[\mathcal{A}^{\mathbb{M}}; \mathbb{C}]$  so that  $d{}_n\mu_{\mathbf{a}}^\phi = \dagger\mathbf{Q}^{(n+1)}(\phi) d\mathbf{q}_{\mathbf{a}}^{(n)}$  as in Lemma 19.

If  $\lambda > 0$  then the sequence  $\{\mathbf{Q}^{(n)}; n \in \mathbb{Z}\}$  is **uniformly harmonically mixing** with **decay parameter**  $\lambda$  (or “ $\lambda$ -UHM”) if, for every  $\mathbf{a} \in \mathcal{A}^{\mathbb{M}}$  and measurable  $\phi \in \mathcal{B}_1$ , and every  $n \in \mathbb{Z}$ , the measure  ${}_n\mu_{\mathbf{a}}^\phi$  is  $\lambda$ -EHM. Thus, applying **Part 1** of Lemma 19, we have:  $\left\| \dagger\mathbf{Q}^{(n+1)} \circ \chi_\bullet \circ \dagger\mathbf{Q}^{(n)} \right\|_\infty \leq e^{-\lambda \cdot R}$  for any  $\chi \in \widehat{\mathcal{A}^{\mathbb{M}}}$  with  $\text{rank}[\chi] \geq R$ .

**Proposition 20:** *Let  $\widetilde{\mathbf{U}} = \mathbb{U} \times \{-1, 0, 1\}$  and  $\mu \in \mathcal{M}_{\text{EAS}}[\mathcal{A}^{\mathbb{M} \times \mathbb{Z}}]$  be a  $\widetilde{\mathbf{U}}$ -MRF such that  $\forall n \in \mathbb{Z}$ ,  $\mathbf{a} \in \mathcal{A}^{\mathbb{M} \times \{n\}}$ , and  $\mathbf{c} \in \mathcal{A}^{\mathbb{M} \times \{n+2\}}$ , the sandwich measure  $\mu_{\mathbf{a}}^{\mathbf{c}}$  is  $\lambda$ -EHM. Then  $\{\mathbf{Q}^{(n)}; n \in \mathbb{Z}\}$  is  $\lambda$ -UHM.*

**Proof:** Let  $\phi \in \mathcal{B}_1$ . By **Part 2** of Lemma 19,  ${}_n\mu_{\mathbf{a}}^\phi = \int_{\mathcal{A}^{\mathbb{M}}} \phi(\mathbf{c}) \cdot \mu_{\mathbf{a}}^{\mathbf{c}} d\mu_{\mathbf{a}}[\mathbf{c}]$ , where  $\mu_{\mathbf{a}} = \mathbf{Q}^{(n+1)} \circ \mathbf{Q}^{(n)}(\delta_{\mathbf{a}})$ . By hypothesis,  $\mu_{\mathbf{a}}^{\mathbf{c}}$  is  $\lambda$ -EHM for all  $\mathbf{c} \in \mathcal{A}^{\mathbb{M}}$ ;

apply Lemma 11 to conclude that  ${}_n\mu_{\mathbf{a}}^{\phi}$  is also  $\lambda$ -EHM.  $\square$

**Proposition 21:** *Let  $\mu \in \mathcal{M}_{\mathcal{EAS}}[\mathcal{A}^{\mathbb{M} \times \mathbb{Z}}]$  be a the path distribution of a Markov process determined by Markov operators  $\{\mathbf{Q}^{(n)}; n \in \mathbb{Z}\}$ .*

*If  $\{\mathbf{Q}^{(n)}; n \in \mathbb{Z}\}$  is  $\lambda$ -UHM, then  $\mu$  is  $\lambda'$ -HM, where  $\lambda' = \lambda/2$ .*

**Proof:** Let  $\chi \in \widehat{\mathcal{A}^{\mathbb{M} \times \mathbb{Z}}}$  with  $\text{rank}[\chi] = 2R$ , and suppose that  $\chi = \bigotimes_{k=0}^{2K} \chi^{(k)}$ , where, for all  $k \in [0..2K]$ ,  $\chi^{(k)}$  is a character on  $\mathcal{A}^{\mathbb{M} \times \{n_k\}}$ , with  $\text{rank}[\chi^{(k)}] = R_k$ , for some  $n_0 < n_1 < \dots < n_{2K}$ . For any  $k \in [1..2K]$ , define  $\dagger\mathbf{Q}_k = \dagger\mathbf{Q}^{(n_k)} \circ \dagger\mathbf{Q}^{(n_{k-1})} \circ \dots \circ \dagger\mathbf{Q}^{(n_{(k-1)+2})} \circ \dagger\mathbf{Q}^{(n_{(k-1)+1})}$ .

If  $\{\eta_n; n \in \mathbb{Z}\} \subset \mathcal{M}_{\mathcal{EAS}}[\mathcal{A}^{\mathbb{M}}]$  are the state distributions of the process, then it is not hard to show:

$$\begin{aligned} \langle \chi, \mu \rangle &= \\ &\left\langle \dagger\mathbf{Q}^{(n_{(2K)+1})} \circ \chi_{\bullet}^{(2K)} \circ \dagger\mathbf{Q}_{2K} \circ \chi_{\bullet}^{(2K-1)} \circ \dagger\mathbf{Q}_{(2K-1)} \circ \dots \circ \dagger\mathbf{Q}_2 \circ \chi_{\bullet}^{(1)} \circ \dagger\mathbf{Q}_1 \left( \chi^{(0)} \right), \right. \\ &\quad \left. \eta_{(n_{(2K)+1})} \right\rangle, \end{aligned}$$

(see e.g. **Claim 1** of Proposition 8 in [5]). Thus,

$$\begin{aligned} &|\langle \chi, \mu \rangle| \\ &\leq_{(1)} \left\| \dagger\mathbf{Q}^{(n_{(2K)+1})} \circ \chi_{\bullet}^{(2K)} \circ \dagger\mathbf{Q}_{2K} \circ \chi_{\bullet}^{(2K-1)} \circ \dots \circ \dagger\mathbf{Q}_2 \circ \chi_{\bullet}^{(1)} \circ \dagger\mathbf{Q}_1 \left( \chi^{(0)} \right) \right\|_{\infty} \\ &\leq_{(2)} \left\| \dagger\mathbf{Q}^{(n_{(2K)+1})} \circ \chi_{\bullet}^{(2K)} \circ \dagger\mathbf{Q}_{2K} \circ \chi_{\bullet}^{(2K-1)} \circ \dots \circ \dagger\mathbf{Q}_2 \circ \chi_{\bullet}^{(1)} \circ \dagger\mathbf{Q}_1 \right\|_{\infty} \\ &\leq_{(3)} \left\| \dagger\mathbf{Q}^{(n_{(2K)+1})} \circ \chi_{\bullet}^{(2K)} \circ \dagger\mathbf{Q}_{2K} \right\|_{\infty} \cdot \prod_{k=1}^{K-1} \left\| \dagger\mathbf{Q}_{(2k+1)} \circ \chi_{\bullet}^{(2k)} \circ \dagger\mathbf{Q}_{2k} \right\|_{\infty} \cdot \prod_{k=0}^{K-1} \left\| \chi_{\bullet}^{(2k+1)} \right\|_{\infty} \\ &\leq_{(4)} \prod_{k=1}^K \left\| \dagger\mathbf{Q}^{(n_{(2k)+1})} \circ \chi_{\bullet}^{(2k)} \circ \dagger\mathbf{Q}^{(n_{(2k)})} \right\|_{\infty} \\ &\leq_{(5)} \prod_{k=1}^K \exp[-\lambda \cdot R_{2k}] = \exp \left[ \sum_{k=1}^K -\lambda \cdot R_{2k} \right] = \exp \left[ -\lambda \cdot \sum_{k=1}^K R_{2k} \right] \end{aligned}$$

(1) Because  $\eta_{(n_{(2K)+1})}$  is a probability measure.

(2) Because  $\left\| \chi^{(0)} \right\|_{\infty} = 1$ .

(3) Separating out  $\chi_{\bullet}^{(k)}$  for all odd  $k$ .

(4) Dropping  $\chi_{\bullet}^{(k)}$  for all odd  $k$ , and  $\dagger\mathbf{Q}^{(n_{(2k)}-1)}$ ,  $\dagger\mathbf{Q}^{(n_{(2k)}-2)}$ ,  $\dots$ ,  $\dagger\mathbf{Q}^{(n_{(2k-1)}+2)}$  for every  $k$  (because  $\|\dagger\mathbf{Q}^{(n)}\| \leq 1$  for every  $n \in \mathbb{Z}$ ).

(5) By UHM hypothesis and **Part 1** of Lemma 19.

$$\text{By the same logic, } |\langle \chi, \mu \rangle| \leq \exp \left[ -\lambda \cdot \sum_{k=1}^K R_{(2k-1)} \right].$$

Now, clearly, one of  $\left( \sum_{k=1}^K R_{2k} \right)$  and  $\left( \sum_{k=1}^K R_{(2k-1)} \right)$  must equal or exceed

$$R, \text{ since together, they sum to } \text{rank}[\chi] = 2R. \text{ Thus either } -\lambda \cdot \sum_{k=1}^K R_{2k} \leq -\lambda \cdot R \text{ or } -\lambda \cdot \sum_{k=1}^K R_{(2k-1)} \leq -\lambda \cdot R. \text{ Hence } |\langle \chi, \mu \rangle| \leq -\lambda \cdot R. \quad \square$$

**Proof of Proposition 14:** If  $\mu$  is a MRF and all sandwich measures of  $\mu$  are  $\lambda$ -EHM, then, by Proposition 20, the sequence  $\{\mathbf{Q}^{(n)}; n \in \mathbb{Z}\}$  is  $\lambda$ -UHM. Then, by Proposition 21,  $\mu$  is  $\lambda'$ -HM, where  $\lambda' = \lambda/2$ .  $\square$

## 6 Harmonic Mixing on the Golden Mean Shift

In [5] and in §5 of the present paper, we have demonstrated harmonic mixing for measures with “full support”, in the sense that every finite cylinder set has nonzero measure. Is full support necessary for harmonic mixing? Is full support of  $\mu$  necessary for the iterates  $\mathfrak{F}^N \mu$  to converge to  $\mathcal{H}^{\text{avr}}$  in Cesàro average? We will answer both these questions in the negative, by proving the following:

**Proposition 22:** *Let  $\mathcal{A} = \mathbb{Z}/2$ . The Markov measure on  $\mathcal{A}^{\mathbb{Z}}$  with transition probability matrix  $\mathbf{Q} = \begin{bmatrix} 1/2 & 1 \\ 1/2 & 0 \end{bmatrix}$  is harmonically mixing.*

**Proof:** Let  $\eta \in \mathcal{M}_{\text{EAS}}[\mathcal{A}]$  be the Perron probability measure for  $\mathbf{Q}$ , so that  $\mathbf{Q}(\eta) = \eta$ , and let  $\mu \in \mathcal{M}_{\text{EAS}}[\mathcal{A}^{\mathbb{Z}}]$  be the Markov measure induced by  $\mathbf{Q}$  and  $\eta$ .

Let  $\mathcal{Q} : \mathbb{R}^2 \curvearrowright$  be the linear operator with matrix  $\dagger\mathbf{Q} = \begin{bmatrix} 1/2 & 1/2 \\ 1 & 0 \end{bmatrix}$ .

Recall that  $\mathbb{Z}/2$  has two characters,  $\mathbf{1}$  and  $\chi$ , where  $\chi(a) := (-1)^a$ . We will use the notation of **Proposition 8** in [5]. In particular  $\mathcal{M}_\chi : \mathbb{R}^2 \curvearrowright$  is the operator with matrix  $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ , while  $\mathcal{M}_\mathbf{1}$  is just the identity operator. Let  $\mathcal{P} = \mathcal{M}_\chi \circ \mathcal{Q}$ .

**Claim 1:**  $\mu$  is harmonically mixing if:

1. For all  $n \geq 1$  and  $\ell \in \{1, 2\}$ ,  $\|\mathcal{Q}^n \circ \mathcal{P}^\ell\| \leq 3/4$ .
2. For all  $n \geq 1$ ,  $\|\mathcal{Q}^n\| \leq 3/2$ .
3.  $\|\mathcal{P}\| = 3/2$  and  $\|\mathcal{P}^2\| = 3/4$ .

**Proof:** Let  $\Xi = \bigotimes_{n=0}^N \xi_n$  be a character of  $(\mathbb{Z}/2)^{\mathbb{Z}}$ , where  $\xi_n \in \{\mathbf{1}, \chi\}$  for all  $n \in [0..N]$ . Let  $\text{rank}[\Xi] = R$ . As in **Claim 1** of **Proposition 8** in [5], we have:

$$\begin{aligned} \widehat{\mu}[\Xi] &= \langle \Xi, \mu \rangle = \langle \mathcal{M}_{\xi_0} \circ \mathcal{Q} \circ \mathcal{M}_{\xi_1} \circ \mathcal{Q} \circ \dots \circ \mathcal{M}_{\xi_{N-1}} \circ \mathcal{Q}[\xi_N], \eta \rangle \\ \text{so that } |\widehat{\mu}[\Xi]| &\leq \|\mathcal{M}_{\xi_0} \circ \mathcal{Q} \circ \mathcal{M}_{\xi_1} \circ \mathcal{Q} \circ \dots \circ \mathcal{M}_{\xi_{N-1}} \circ \mathcal{Q}[\xi_N]\|_\infty \\ &\leq \|\mathcal{M}_{\xi_0} \circ \mathcal{Q} \circ \mathcal{M}_{\xi_1} \circ \mathcal{Q} \circ \dots \circ \mathcal{M}_{\xi_{N-1}} \circ \mathcal{Q}[\xi_N]\|_1 \\ &\leq \|\mathcal{M}_{\xi_0} \circ \mathcal{Q} \circ \mathcal{M}_{\xi_1} \circ \mathcal{Q} \circ \dots \circ \mathcal{M}_{\xi_{N-1}} \circ \mathcal{Q}\| \cdot \|\xi_N\|_1, \end{aligned}$$

where  $\|\xi_N\|_1 = |\xi_N(0)| + |\xi_N(1)| = 2$ , and, for any operator  $\mathcal{R} : \mathbb{R}^2 \curvearrowright$ ,  $\|\mathcal{R}\|$  the operator norm of  $\mathcal{R}$  relative to the norm  $\|\bullet\|_1$  on  $\mathbb{R}^2$ .

To bound  $\|\mathcal{M}_{\xi_0} \circ \mathcal{Q} \circ \mathcal{M}_{\xi_1} \circ \mathcal{Q} \circ \dots \circ \mathcal{M}_{\xi_{N-1}} \circ \mathcal{Q}\|$ , write

$$\begin{aligned} &\mathcal{M}_{\xi_0} \circ \mathcal{Q} \circ \mathcal{M}_{\xi_1} \circ \mathcal{Q} \circ \dots \circ \mathcal{M}_{\xi_{N-1}} \circ \mathcal{Q} \\ &= \mathcal{P}^{\ell_0} \circ \mathcal{P}^{2m_0} \circ (\mathcal{Q}^{n_1} \circ \mathcal{P}^{\ell_1}) \circ \mathcal{P}^{2m_1} \circ (\mathcal{Q}^{n_2} \circ \mathcal{P}^{\ell_2}) \circ \mathcal{P}^{2m_2} \circ \dots \\ &\quad \dots \circ (\mathcal{Q}^{n_{K-1}} \circ \mathcal{P}^{\ell_{K-1}}) \circ \mathcal{P}^{2m_{K-1}} \circ \mathcal{Q}^{n_K} \end{aligned}$$

where,  $\ell_0 \in \{0, 1\}$  and  $\forall k \in (0..K]$ ,  $\ell_k \in \{1, 2\}$ , and  $\forall k \in [0..K]$ ,  $m_k, n_k \geq 0$ , with  $n_k \neq 0$  if  $k \neq K$ . Thus

$$\begin{aligned} &\|\mathcal{M}_{\xi_0} \circ \mathcal{Q} \circ \mathcal{M}_{\xi_1} \circ \mathcal{Q} \circ \dots \circ \mathcal{M}_{\xi_{N-1}} \circ \mathcal{Q}\| \\ &\leq \|\mathcal{P}^{\ell_0}\| \cdot \|\mathcal{P}^{2m_0}\| \cdot \|\mathcal{Q}^{n_1} \circ \mathcal{P}^{\ell_1}\| \cdot \|\mathcal{P}^{2m_1}\| \cdot \|\mathcal{Q}^{n_2} \circ \mathcal{P}^{\ell_2}\| \cdot \|\mathcal{P}^{2m_2}\| \cdot \dots \\ &\quad \dots \|\mathcal{Q}^{n_{K-1}} \circ \mathcal{P}^{\ell_{K-1}}\| \cdot \|\mathcal{P}^{2m_K}\| \cdot \|\mathcal{Q}^{n_K}\| \\ &\leq \left(\frac{3}{2}\right) \cdot \left(\frac{3}{4}\right)^{m_0} \cdot \left(\frac{3}{4}\right) \cdot \left(\frac{3}{4}\right)^{m_1} \cdot \dots \cdot \left(\frac{3}{4}\right) \cdot \left(\frac{3}{4}\right)^{m_K} \cdot \left(\frac{3}{2}\right) \\ &= \left(\frac{3}{2}\right)^2 \cdot \left(\frac{3}{4}\right)^{K+(m_0+\dots+m_K)} \stackrel{(1)}{\leq} \left(\frac{3}{2}\right)^2 \cdot \left(\frac{3}{4}\right)^{R/2} \xrightarrow{R \rightarrow \infty} 0, \end{aligned}$$

(1) Because  $R = \sum_{k=0}^{K-1} \ell_k + \sum_{k=0}^{K-1} 2m_k \leq 2K+2 \cdot \left( \sum_{k=0}^{K-1} m_k \right)$ .  $\square$  [Claim 1]

To verify these operator norms, let  $\mathbb{B}_1 = \{\mathbf{x} \in \mathbb{R}^2; \|\mathbf{x}\|_1 = 1\}$ ; if  $\mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$  and  $\mathbf{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ , then the extremal set of  $\mathbb{B}_1$  is  $\{\pm\mathbf{e}_1, \pm\mathbf{e}_2\}$ . Thus, for any operator  $\mathcal{R} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ , with matrix  $\mathbf{R} = \begin{bmatrix} r_{11} & r_{12} \\ r_{21} & r_{22} \end{bmatrix}$ , it is clear that

$$\|\mathcal{R}\| = \sup_{\mathbf{x} \in \mathbb{B}_1} \|\mathcal{R}(\mathbf{x})\|_1 = \sup_{\mathbf{x} \in \{\pm\mathbf{e}_1, \pm\mathbf{e}_2\}} \|\mathcal{R}(\mathbf{x})\|_1 = \max \left\{ \left\| \begin{bmatrix} r_{11} \\ r_{21} \end{bmatrix} \right\|_1, \left\| \begin{bmatrix} r_{12} \\ r_{22} \end{bmatrix} \right\|_1 \right\}.$$

At this point, hypothesis 3 of **Claim 1** can be verified immediately. The other two hypotheses can be proved by induction. For example: **Claim 2:**  $\forall n \geq 1$  and  $\ell \in \{1, 2\}$ ,  $\|\mathcal{Q}^n \circ \mathcal{P}^\ell\| \leq 3/4$ .

**Proof:** Check that  $\|\mathcal{Q} \circ \mathcal{P}\| = \frac{3}{4}$ ,  $\|\mathcal{Q}^2 \circ \mathcal{P}\| = \frac{5}{8}$ ,  $\|\mathcal{Q} \circ \mathcal{P}^2\| = \frac{5}{8}$ , and  $\|\mathcal{Q}^2 \circ \mathcal{P}^2\| = \frac{11}{6}$ . If  $n \geq 3$ , then assume, inductively, that  $\|\mathcal{Q}^{n-2} \circ \mathcal{P}^\ell\| \leq \frac{3}{4}$  and  $\|\mathcal{Q}^{n-1} \circ \mathcal{P}^\ell\| \leq \frac{3}{4}$ .

Let  $\begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}^2$ , with  $\left\| \begin{bmatrix} x \\ y \end{bmatrix} \right\|_1 = 1$ , and, for all  $n \in \mathbb{N}$ , define  $\begin{bmatrix} x_n \\ y_n \end{bmatrix} := \mathcal{Q}^n \circ \mathcal{P}^\ell \begin{bmatrix} x \\ y \end{bmatrix}$ . Then  $\mathcal{Q}^n \circ \mathcal{P}^\ell \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \frac{x_{n-1} + y_{n-1}}{2} \\ x_{n-1} \end{bmatrix} = \begin{bmatrix} \frac{x_{n-1}}{2} + \frac{y_{n-1}}{2} \\ \frac{x_{n-2}}{2} + \frac{y_{n-2}}{2} \end{bmatrix}$ .

Thus, we have

$$\begin{aligned} \left\| \mathcal{Q}^n \circ \mathcal{P}^\ell \begin{bmatrix} x \\ y \end{bmatrix} \right\|_1 &\leq \left\| \frac{x_{n-1}}{2} + \frac{y_{n-1}}{2} \right\|_1 + \left\| \frac{x_{n-2}}{2} + \frac{y_{n-2}}{2} \right\|_1 \\ &\leq \frac{1}{2} \left\| \begin{bmatrix} x_{n-1} \\ y_{n-1} \end{bmatrix} \right\|_1 + \frac{1}{2} \left\| \begin{bmatrix} x_{n-2} \\ y_{n-2} \end{bmatrix} \right\|_1 \\ &\leq \frac{1}{2} \left\| \mathcal{Q}^{n-1} \circ \mathcal{P}^\ell \begin{bmatrix} x \\ y \end{bmatrix} \right\|_1 + \frac{1}{2} \left\| \mathcal{Q}^{n-2} \circ \mathcal{P}^\ell \begin{bmatrix} x \\ y \end{bmatrix} \right\|_1 \\ &\leq \frac{\|\mathcal{Q}^{n-2} \circ \mathcal{P}^\ell\| + \|\mathcal{Q}^{n-1} \circ \mathcal{P}^\ell\|}{2} \leq \frac{3}{4}. \end{aligned}$$

.....  $\square$  [Claim 2]

---

$\square$

The measure  $\mu$  of Proposition 22 is supported on the subshift of finite type with transition matrix  $\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$ , sometimes called the **Golden Mean**

subshift; this is the set of all sequences in  $\{0, 1\}^{\mathbb{Z}}$  where the symbol “1” never appears twice in a row. Clearly,  $\mu$  does not have full support in  $\{0, 1\}^{\mathbb{Z}}$ , since any cylinder set containing two consecutive “1”s gets zero probability.

There is nothing special about the choice of  $\begin{bmatrix} 1/2 \\ 1/2 \end{bmatrix}$  for the first column of  $\mathbf{Q}$ . The Markov measures induced by matrices of the form  $\begin{bmatrix} (1 - \rho) & 1 \\ \rho & 0 \end{bmatrix}$  are harmonically mixing for values of  $\rho$  ranging at least over  $(0.5, 0.8)$ ; this can be verified computationally by checking that  $\|\mathcal{Q}^n\|$ ,  $\|\mathcal{Q}^n \circ \mathcal{P}^m\|$ , etc. are strictly less than 1.

Unfortunately, the proof method of Proposition 22 breaks down when  $\rho \neq 1/2$ . However, the method can be applied to other subshifts with “equally weighted transitions”. For example, a similar argument demonstrates harmonic mixing for the Markov measure on  $(\mathbb{Z}/3)^{\mathbb{Z}}$  induced by

$$\begin{bmatrix} 1/3 & 0 & 1/3 \\ 1/3 & 0 & 1/3 \\ 1/3 & 1 & 1/3 \end{bmatrix}.$$

However, as yet there is no simple characterization of harmonic mixing for arbitrary Markov measures on subshifts of finite type.

## 7 The Even Shift is Not Harmonically Mixing

Harmonic mixing seems to arise in measures with a high level of “randomness”, such as fully supported Markov random fields. What other “randomness” properties yield harmonic mixing?

A measure  $\mu$  on  $\mathcal{A}^{\mathbb{Z}}$  has the **Kolmogorov** or **K** property if every factor of the measure preserving dynamical system  $(\mathcal{A}^{\mathbb{Z}}, \mu, \sigma)$  has nonzero entropy [8]. Every mixing Markov measure is **K**. The **K** property implies that  $(\mathcal{A}^{\mathbb{Z}}, \mu, \sigma)$  has Lebesgue spectrum and thus is mixing; in a sense, **K** means that  $(\mathcal{A}^{\mathbb{Z}}, \mu, \sigma)$  is “almost” a Bernoulli system. Is the **K** property sufficient for harmonic mixing? We will show that it is not, constructing a **K** measure that is not harmonically mixing.

Let  $\mathbf{X} = (\mathbb{Z}/3)^{\mathbb{Z}}$ , and consider  $\mathbf{X}_A \subset \mathbf{X}$ , the subshift of finite type defined by the transition matrix

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, \text{ where, } \forall i, j \in \mathbb{Z}/3, a_{ij} = \begin{cases} 1 & \text{if } j \rightsquigarrow i \text{ is allowed} \\ 0 & \text{if } j \rightsquigarrow i \text{ is not allowed} \end{cases}$$

Let  $\phi : \mathbf{X}_A \rightarrow (\mathbb{Z}/2)^\mathbb{Z}$  be the factor map of radius 0 which sends 0 into 0 and both 1 and 2 to 1. Then  $\mathbf{Y} := \phi(\mathbf{X}_A)$  is Weiss' *Even Sofic Shift*: if  $\mathbf{y} \in \mathbf{Y}$ , then there are an even number of 1's between any two occurrences of 0 in  $\mathbf{y}$ .

For any  $N \in \mathbb{N}$ , and  $i, j \in \mathbb{Z}/3$ , let  $\mathbf{X}_{ij}^N = \{\mathbf{x} \in \mathbf{X}_A ; x_0 = i, x_N = j\}$ , and let:

$$\mathbf{E}_N := \left\{ \mathbf{y} \in \mathbf{Y} ; \sum_{n=0}^N y_n \text{ is even} \right\}, \text{ and } \mathbf{O}_N := \left\{ \mathbf{y} \in \mathbf{Y} ; \sum_{n=0}^N y_n \text{ is odd} \right\}.$$

**Lemma 23:**  $\forall i, j \in \mathbb{Z}/3$ , either  $\phi(\mathbf{X}_{i,j}^N) \subset \mathbf{E}_N$  or  $\phi(\mathbf{X}_{i,j}^N) \subset \mathbf{O}_N$ . In particular,

$$\begin{aligned} \phi(\mathbf{X}_{0,0}^N \sqcup \mathbf{X}_{1,2}^N \sqcup \mathbf{X}_{2,1}^N \sqcup \mathbf{X}_{0,2}^N \sqcup \mathbf{X}_{1,0}^N) &= \mathbf{E}_N, \\ \text{and } \phi(\mathbf{X}_{1,1}^N \sqcup \mathbf{X}_{0,1}^N \sqcup \mathbf{X}_{2,0}^N \sqcup \mathbf{X}_{2,2}^N) &= \mathbf{O}_N. \end{aligned}$$

**Proof:** Let  $\mathbf{x} \in \mathbf{X}_{ij}^N$ , and  $\mathbf{y} = \phi(\mathbf{x})$ . Note that, if  $k < k^*$  are any

two values so that  $x_k = 0 = x_{k^*}$ , then  $\sum_{n=k}^{k^*} y_n$  is even. In particular, let  $k$  be the first element of  $[0 \dots N]$  where  $x_k = 0$ , and let  $k^*$  be the last element of  $[0 \dots N]$  where  $x_{k^*} = 0$ . Thus,  $\sum_{n=k}^{k^*} y_n \equiv 0 \pmod{2}$ , so that

$$\sum_{n=0}^N y_n \equiv \sum_{n=0}^{k-1} y_n + \sum_{n=k^*+1}^N y_n \pmod{2}.$$

But since  $x_{k-1} \neq 0 \neq x_{k^*+1}$  by construction, the definition of  $\mathbf{X}_A$  forces  $x_{k-1} = 2$  and  $x_{k^*+1} = 1$ . Thus the parity of  $\sum_{n=0}^{k-1} y_n$  depends only on the

value of  $x_0 = i$ . Similarly the parity of  $\sum_{n=k^*+1}^N y_n$  depends only on  $x_N = j$ .

□

Let  $\mu \in \mathcal{M}_{\text{EAS}}[\mathbf{X}_A]$  be a mixing Markov measure on  $\mathbf{X}_A$ , with transition matrix  $\mathbf{Q}$  and Perron measure  $\boldsymbol{\eta} = (\eta_0, \eta_1, \eta_2) \in \mathcal{M}_{\text{EAS}}[\mathbb{Z}/3]$ . Let  $\nu = \phi\mu \in \mathcal{M}_{\text{EAS}}[\mathbf{Y}]$ , so that if  $C \subset \mathbf{Y}$  is measurable, then  $\nu[C] := \mu[\phi^{-1}(C)]$

For all  $N \in \mathbb{N}$ , let  $\chi_N(\mathbf{x}) = \prod_{n=0}^N (-1)^{x_n} \in \widehat{(\mathbb{Z}/2)^{\mathbb{Z}}}$ . Then, by Lemma 23.

$$\begin{aligned} \int \chi_N d\nu &= \nu(\mathbf{E}_N) - \nu(\mathbf{O}_N) \\ &= \mu(\mathbf{X}_{0,0}^N \sqcup \mathbf{X}_{1,2}^N \sqcup \mathbf{X}_{2,1}^N \sqcup \mathbf{X}_{0,2}^N \sqcup \mathbf{X}_{1,0}^N) - \nu(\mathbf{X}_{1,1}^N \sqcup \mathbf{X}_{0,1}^N \sqcup \mathbf{X}_{2,0}^N \sqcup \mathbf{X}_{2,2}^N), \end{aligned}$$

But  $\mu$  is mixing, so  $\lim_{N \rightarrow \infty} \mu(\mathbf{X}_{i,j}^N) = \eta_i \cdot \eta_j$ . Thus,  $\lim_{N \rightarrow \infty} \int \chi_N d\nu = \eta_0^2 + 2\eta_1\eta_2 - \eta_1^2 - \eta_2^2$ . So for example if

$$\mathbf{Q} = \begin{bmatrix} 1/2 & 0 & 1/2 \\ 1/2 & 0 & 1/2 \\ 0 & 1 & 0 \end{bmatrix}$$

with Perron measure  $\boldsymbol{\eta}(\frac{2}{5}, \frac{1}{5}, \frac{2}{5})$  then  $\nu$  is not harmonically mixing.

On the other hand, since  $\mu$  is a Markov measure, it has the **K** property. Every factor of  $(\mathbf{X}_A, \mu; \boldsymbol{\sigma})$  also has the **K** property, including  $(\mathbf{Y}, \nu; \boldsymbol{\sigma})$ . Hence,  $\nu$  is a **K** measure, but is not harmonically mixing.

## 8 Conclusion

We have demonstrated that a broad class of probability measures on  $\mathcal{A}^{\mathbb{M}}$  weak\*-converge to Haar measure in density, when acted on by a wide class of LCA. Many problems remain open, however. For example, in §6, we showed that full support is not necessary for a Markov measure on  $\mathcal{A}^{\mathbb{Z}}$  to be harmonically mixing. Is there a general characterization for harmonic mixing of Markov measures supported on subshifts of finite type? Also, is there any characterization of either diffusion or harmonic mixing when  $\mathbb{M}$  is a nonabelian monoid? Finally, what happens when  $\mathcal{A}$  is a nonabelian group? The natural analogy of LCA for nonabelian  $\mathcal{A}$  are “multiplicative” cellular automata [9], where the local map is computed by (noncommutatively) multiplying the values of neighbouring coordinates. What is the asymptotic behaviour of measures under such automata?

## References

- [1] Alejandro Maass and Servet Martínez. Time averages for some classes of expansive one-dimensional cellular automata. In Eric Goles and Servet Martínez, editors, *Cellular Automata and Complex Systems*, pages 37–54. Kluwer Academic Publishers, Dordrecht, 1999.



- [2] H. Furstenberg. *Ergodic Theory and Combinatorial Number Theory*. Princeton University Press, Princeton, New Jersey, 1981.
- [3] G. Hedlund. Endomorphisms and automorphisms of the shift dynamical systems. *Mathematical System Theory*, 3:320–375, 1969.
- [4] Douglas Lind. Applications of ergodic theory and sofic systems to cellular automata. *Physica D*, 10:36–44, 1984.
- [5] Marcus Pivato and Reem Yassawi. Limit measures for affine cellular automata. To appear in *Ergodic Theory and Dynamical Systems*; preprint available at: <http://arXiv.org>, 2001.
- [6] Servet Martínez Pablo Ferrari, Alejandro Maass. Cesàro mean distribution of group automata starting from Markov measures. (*preprint*), 1998.
- [7] Servet Martínez Pablo Ferrari, Alejandro Maass and Peter Ney. Cesàro mean distribution of group automata starting from measures with summable decay. *Ergodic Theory and Dynamical Systems*, 20(6):1657–1670, 2000.
- [8] Karl Petersen. *Ergodic Theory*. Cambridge University Press, New York, 1989.
- [9] Marcus Pivato. Nonabelian multiplicative cellular automata. Submitted to *Ergodic Theory and Dynamical Systems*; preprint available at: <http://arXiv.org>, August 2001.
- [10] Ross Kindermann and J. Laurie Snell. *Markov Random Fields and their Applications*. American Mathematical Society, Providence, Rhode Island, 1980.
- [11] Laurent Schwartz. *Lectures on disintegration of measures*. Tata Institute of Fundamental Research, Bombay, 1975.