

Open Research Online

The Open University's repository of research publications and other research outputs

The UV surface habitability of Proxima *b*: first experiments revealing probable life survival to stellar flares

Journal Item

How to cite:

Abrevaya, X. C.; Leitzinger, M.; Opezzo, O. J.; Odert, P.; Patel, M. R.; Luna, G. J. M.; Forte-Giacobone, A. F. and Hansmeier, A. (2020). The UV surface habitability of Proxima *b*: first experiments revealing probable life survival to stellar flares. *Monthly Notices of the Royal Astronomical Society: Letters*, 494(1) L69-L74.

For guidance on citations see [FAQs](#).

© 2020 The Author(s)



<https://creativecommons.org/licenses/by/4.0/>

Version: Version of Record

Link(s) to article on publisher's website:

<http://dx.doi.org/doi:10.1093/mnrasl/slaa037>

<https://academic.oup.com/mnrasl/article/494/1/L69/5762693>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

The UV surface habitability of Proxima *b*: first experiments revealing probable life survival to stellar flares

X. C. Abrevaya,^{1,2,3★†} M. Leitzinger,³ O. J. Oppezzo,⁴ P. Odert,³ M. R. Patel,⁵
G. J. M. Luna,^{1,2} A. F. Forte Giacobone^{4,6} and A. Hanslmeier³

¹*Instituto de Astronomía y Física del Espacio (UBA - CONICET), Pabellón IAFE, Ciudad Universitaria, C1428EGA, Ciudad Autónoma de Buenos Aires, Argentina*

²*Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Ciudad Universitaria, C1428EGA, Ciudad Autónoma de Buenos Aires, Argentina*

³*Institute of Physics, IGAM, University of Graz, Universitätsplatz 5, 8010, Graz, Austria*

⁴*Comisión Nacional de Energía Atómica, Centro Atómico Constituyentes, B1650KNA, Buenos Aires, Argentina*

⁵*School of Physical Sciences, The Open University, Milton Keynes MK7 6AA, UK*

⁶*Departamento de Ingeniería, Universidad Nacional de Tres de Febrero, Saenz Peña, B1674AHF, Buenos Aires, Argentina*

Accepted 2020 February 21. Received 2020 February 19; in original form 2019 October 10

ABSTRACT

We use a new interdisciplinary approach to study the UV surface habitability of Proxima *b* under quiescent and flaring stellar conditions. We assumed planetary atmospheric compositions based on CO₂ and N₂ and surface pressures from 100 to 5000 mbar. Our results show that the combination of these atmospheric compositions and pressures provide enough shielding from the most damaging UV wavelengths, expanding the ‘UV-protective’ planetary atmospheric compositions beyond ozone. Additionally, we show that the UV radiation reaching the surface of Proxima *b* during quiescent conditions would be negligible from the biological point of view, even without an atmosphere. Given that high UV fluxes could challenge the existence of life, then, we experimentally tested the effect that flares would have on microorganisms in a ‘worst case scenario’ (no UV-shielding). Our results show the impact that a typical flare and a superflare would have on life: when microorganisms receive very high fluences of UVC, such as those expected to reach the surface of Proxima *b* after a typical flare or a superflare, a fraction of the population is able to survive. Our study suggests that life could cope with highly UV irradiated environments in exoplanets under conditions that cannot be found on Earth.

Key words: astrobiology – planets and satellites: surfaces – stars: activity – stars: flare – ultraviolet: stars.

1 INTRODUCTION

Stellar ultraviolet radiation (UVR) is an important factor that can influence the existence of ‘life as we know it’ on the surface of planetary bodies (for review on the topic see: Abrevaya & Thomas 2018; Hanslmeier 2018). Although several studies have focused on stellar UVR aiming to evaluate the chances of an exoplanet to host life on its surface, they are based on modelling (see e.g. Cockell 1999; Buccino, Lemarchand & Mauas 2006, 2007; Fossati et al. 2012; Rugheimer et al. 2015; O’Malley-James & Kaltenecker 2017, 2019; Estrela & Valio 2018; Howard et al. 2018). Some of these approaches have neglected some important factors that may

lead to: (i) an inaccurate estimation of the UV flux on the surface of the planet, as some studies do not assume the presence of a planetary atmosphere or assume unrealistic atmospheric compositions for the stellar spectral type and activity (e.g. O₂ rich Earth-like atmospheric compositions are unfeasible in planets orbiting M dwarfs, e.g. Johnstone et al. 2019) (ii) an inaccurate estimation of the biological impact of UV by using extrapolations of biologically relevant UV fluences¹ based on empirical data from the literature, such as the UV biological action spectra (UV-BAS; a measure of biological UV damage at a given UV wavelength under a given constant fluence), or the UV ‘lethal dose’ (UV-LD; the estimated amount

* E-mail: abrevaya@iafe.uba.ar

† Visiting scientist at the University of Graz.

of UVR required to kill a determined percentage of the population of the life forms under consideration). Even though UV-BAS or UV-LD can provide information about biological effectiveness of radiation or survival of the population, respectively, there are some drawbacks related to their use. For instance, the values from those extrapolations are taken in general from studies that used low fluence rates,² in contrast to those expected for more realistic astrophysical scenarios. Another factor, usually not considered, is that the survival of the microorganisms will vary according to their growth stage (e.g. growth phase) and the physiological conditions of the population (e.g. starvation). Thus, these kind of extrapolations might not represent accurately the effects of radiation on life considering particular planetary environments and can lead, therefore, to potentially biased results. An example of a preliminary laboratory assay has shown an increase in the biomass of microorganisms exposed to UVC fluences comparable to weak flares (Abrevaya, Cortón & Mauas 2011/12). However, these results are only an approximation and cannot be taken as parameter to predict the survival to flares, as the microorganisms were in a physiological condition unfeasible to find in nature and the experiments included the UV-protective effects of the culture medium, among others. Previously, we proposed how to address the issues derived from the studies above by using an interdisciplinary astrobiological approach that combines astrophysics with microbiology and photobiology in laboratory experiments (Abrevaya et al. 2018/20). In this Letter, we present the first results and conclusions obtained from this approach, taking as a case study Proxima Centauri *b* (Proxima *b*), currently, the closest potentially habitable terrestrial-mass planet outside the Solar system, located in the habitable zone of Proxima Centauri (Proxima Cen), an M6V dwarf (Anglada-Escudé et al. 2016). We applied a radiative transfer modelling to estimate the UVR at the surface of Proxima *b* during quiescent stellar conditions, taking into account different atmospheric compositions and pressures. Those estimations led us to focus on experiments to determine the biological effects of stellar flares on Proxima *b*, considering unattenuated UVC surface fluxes in a ‘worst case scenario’.

This study is part of the EXO-UV programme (Abrevaya et al. 2014), an international interdisciplinary collaboration that seeks to expand the characterization of UVR environments through experimental approaches.

2 METHODS

2.1 Estimation of UV surface fluxes during quiescent stellar conditions

The quiescent stellar spectrum of Proxima Cen was obtained from Meadows et al. (2018) which comprises observational data as well as a model spectrum (PHOENIX 2.0, Husser et al. 2013) for spectral regions where no observational data were available. To investigate how much UVR reaches the planetary surface depending on the assumed atmospheric composition, an extended version of the radiative transfer model by Patel, Zarnecki & Catling (2002) was used, as previously applied for investigations of terrestrial-like atmospheric transmittance on exoplanets orbiting cool white dwarf stars (Fossati et al. 2012). The model calculates the stellar spectrum received at the surface of the planet for a zenith angle of 90°, i.e.

²Defined according to Braslavsky (2007) as: total radiant power, incident from all directions on to a small sphere divided by the cross-sectional area of that sphere. SI unit W m^{-2} .

at the equator, in order to provide the case of maximum irradiance. Very recently, it was predicted that Earth-like atmospheres cannot form in the habitable zones of very active stars (Johnstone et al. 2019), and atmospheres containing CO₂ are likely stable on water-depleted planets orbiting M dwarfs (Gao et al. 2015). Atmospheres dominated by N₂ tend to be unstable under high-XUV fluxes and experience efficient atmospheric escape (Tian et al. 2008; Lammer et al. 2011). However, sufficient levels of CO₂ can protect nitrogen atmospheres from escaping due to infrared cooling (Lichtenegger et al. 2010; Johnstone et al. 2018). Based on these studies, a range of realistic atmospheric compositions were considered: (a) 100 per cent CO₂, (b) 10 per cent CO₂ and 90 per cent N₂, (c) 50 per cent CO₂ and 50 per cent N₂, and (d) 90 per cent CO₂ and 10 per cent N₂, with surface pressures of 100, 200, 500, 1000, 2000, and 5000 mbar.

2.2 Estimation of UV surface fluxes during flares

Two stellar flare scenarios for Proxima Cen were considered, a typical flare and a superflare. For the estimation of a typical flare-related UVC flux enhancement for M dwarfs, we used the study of Welsh et al. (2007), who analysed 52 flares detected by the Galaxy Evolution Explorer in NUV band (175–280 nm), which covers the UVC region. They estimate an average NUV flare energy of 2.5×10^{30} erg for 33 M dwarf flare events, with a range covering 1.2×10^{28} – 1.6×10^{31} erg. Scaling their given flare luminosities to the orbit of Proxima *b* (0.0485 au) yields fluxes in the range of about 5×10^{-4} to 80 W m^{-2} with a mean of $\sim 7 \text{ W m}^{-2}$. Our experiments were then conducted with a value of 8.7 W m^{-2} , which is comparable to this mean value. Since flares follow a power-law distribution according to their energy, there are more flares with low energies and fewer flares with high energies. Therefore, it is clear that most flares from Proxima Cen will be smaller than our assumed ‘typical’ flare value. However, such small flares would provide a negligible addition to the quiescent flux level. Furthermore, we estimated a UVC peak fluence rate of 92 W m^{-2} for the superflare on Proxima based on Howard et al. (2018). The estimated total energy of this superflare is comparable to the ‘great flare’ on AD Leo ($\sim 10^{34}$ erg, Hawley & Pettersen 1991; Segura et al. 2010).

2.3 Microorganisms and irradiation conditions

To evaluate the potential biological damage of flares on the surface of Proxima *b*, considering the UVC range (as it is the most biologically harmful) in a worst case scenario (unattenuated conditions) we performed experiments with microorganisms from different domains: *Haloferax volcanii*, an archeon inhabiting extreme hypersaline environments, and *Pseudomonas aeruginosa*, an ubiquitous bacterium. For the irradiation procedure suspensions of the microorganisms were irradiated with UVC tube lamps at fluence rates of 8.7 W m^{-2} (typical flare) or 92 W m^{-2} (superflare). Irradiation times were from 5 to 4500 s for the flare, considering the typical distribution for the duration of flares described in Gershberg (2005), and from 5 to 420 s for the superflare, considering the peak of the superflare described in Howard et al. (2018). The control groups were non-irradiated suspensions. To test the potential effect of the influence of visible light in the repair capability to DNA damage in the microorganisms, in another experiment, samples of *P. aeruginosa* were exposed for 3 h to visible light after irradiated at a fluence rate of 92 W m^{-2} . The control groups followed the same procedure in the dark. Technical details about the irradiation procedure, growth, and enumeration of microorganisms, and limit

of detection (LOD), can be found in the Appendix. All experiments were performed in triplicate (three independent experiments).

3 RESULTS AND DISCUSSION

3.1 UV radiation on the planetary surface

As presented in Section 2.1, our radiative transfer calculations show the influence of the different atmospheric compositions and pressures on the stellar UVR that reaches the surface of Proxima *b*. Different absorption of the UV (200–380 nm) flux for the different atmospheric scenarios were obtained (see Fig. A), consistent with previous studies by Ranjan, Wordsworth & Sasselov (2017; see comparison in Fig. B).

The UVR is dominated by the UVA range (315–400 nm) providing 92 per cent of the incident radiation. Without atmospheric attenuation the UVA flux is 1.08 W m^{-2} , a value much lower than the UVA flux at the Earth’s surface (e.g. $30\text{--}50 \text{ W m}^{-2}$ for clear sky). This value would be even lower under the attenuation of the considered atmospheres. Thus, the UVA flux on the surface of Proxima *b* would not prevent the existence of ‘life as we know it’. The effectiveness of UVA to induce lethal damage depends also on mechanisms that in the presence of oxygen produce reactive oxygen species (Kielbassa, Roza & Epe 1997; Opezzo, Costa & Pizzarro 2011). According to several studies O-rich atmospheres are not expected to exist in planets orbiting M-type stars (Garcia-Sage et al. 2017; Johnstone et al. 2019), therefore, this would reduce the damage induced by UVA in this context.

For the UVB (280–315 nm) and UVC (200–280 nm) bands, atmospheric absorption would increase with the CO_2 content and with the atmospheric pressure, regardless of the atmospheric compositions (see Figs A1 and A2). The flux in these bands would be reduced by several orders of magnitude for CO_2 contents of 50 per cent or more, and for atmospheric pressures higher than 2000 mbar. This is particularly relevant because UVB and UVC radiations are absorbed by nucleic acids, which can undergo chemical modifications and lose their biological functions upon exposure (Kielbassa et al. 1997). Nevertheless, the stellar UVB fluence rate reaching the surface of Proxima *b* (0.026 W m^{-2}) is lower than that at the Earth’s surface which is around 2 W m^{-2} . Therefore, microorganisms would be expected to survive even without the protective effect of an atmosphere. For the particular case of UVC, since the terrestrial atmosphere blocks this wavelength range, the information obtained by experiments involving solar radiation at the Earth’s surface cannot be used as a reference to evaluate the impact of UVC on life on Proxima *b*. The only data available about biological effects of solar UVC in microorganisms come from space experiments performed in the low Earth orbit. Several microorganisms exposed to unattenuated extraterrestrial solar UVR spectrum in space experiments, for instance, cells of *Chroococcidiopsis* and spores of *Aspergillus sydowii* and *Aspergillus versicolor* survived being exposed during 1.5 yr in the low Earth orbit (Cockell et al. 2011; Novikova et al. 2015). In addition, viable spores of *Bacillus subtilis* were recovered after 6 yr of exposure to solar UVR in space (Horneck, Bucker & Reitz 1994).

We calculated the solar UVC fluence rate at the top of the atmosphere based on solar spectral irradiance data (Meftah et al. 2018) resulting in an UVC fluence rate of 6.27 W m^{-2} , which is around 95 times higher than the UVC flux value estimated at the surface of Proxima *b* (0.066 W m^{-2} , without any atmospheric protection). Therefore, it seems likely that ‘life as we know it’




Figure 1. Survival curve for *H. volcanii* exposed to a fluence rate of 8.7 W m^{-2} (typical flare). N_0 is the viable count of microorganisms at the beginning of the irradiation, N is the viable count after the irradiated time indicated in the abscissa. The LOD value was -6.76 . Error bars indicate three independent experiments. Inset: detail of the curve for the first 15 s of irradiation.

could thrive on the surface of Proxima *b* under the very low UVC irradiation levels expected during quiescent stellar conditions.

In addition, if attenuation of UVC by the atmosphere is taken into account as we predicted in our modelling, the combination of some atmospheric compositions and pressures would provide even more protection against the most harmful UV wavelengths for life (Fig. A1 and A2). In this way, planets with O-rich Earth-like atmospheric compositions are not the only ones offering protection to UVR. This finding expands the results presented in previous studies (see e.g. O’Malley-James & Kaltenecker 2019).

3.2 Biological effects of flares

Given that on the surface of Proxima *b* the UV fluence rates during quiescence are negligible from the biological point of view, we decided to focus on flare activity and we did not conduct experiments for quiescent levels of UVR. Taking into account that UVC is the most deleterious component of UVR, we performed experiments to evaluate the biological damage of UVC fluence rates related to flares on the surface of Proxima *b*, in a worst case scenario (i.e.: no shielding by biological or environmental factors, including no absorption by the atmosphere).

The microorganisms were exposed to fluences comparable to those estimated for the peak of a typical flare (Figs 1 and 2) or a superflare (Fig. 3). The maximum irradiation time for the typical flare (Figs 1 and 2) was 4500 s, which corresponds to an energy of $\sim 2.6 \times 10^{32}$ erg. For the superflare (Fig. 3), the total irradiation time was 420 s, corresponding to a similar energy. The survival curves obtained for the microorganisms exhibit the shape described as typical for the response of a bacterial population to UVC (Coohill & Sagripanti 2008). At the beginning of the irradiation, the survival fraction reduced exponentially, as most of the exposed microorganisms lost their viability.




Figure 2. Survival curve for *P. aeruginosa* exposed to a fluence rate of 8.7 W m^{-2} (typical flare). The meaning of N , N_0 , and error bars is the same as indicated in Fig. 2. The LOD value was -7.32 .




Figure 3. Survival curve for *P. aeruginosa* exposed to a fluence rate of 92 W m^{-2} (superflare). The meaning of N , N_0 , and error bars is the same as indicated in Fig. 2. The LOD value was -7.03 . Inset box: Influence of visible light on the survival of irradiated microorganisms. White bars indicate photoreactivating conditions, grey bars correspond to the control group (incubation in dark conditions).

Subsequently, a remarkable reduction in the slopes of the survival curves was observed when irradiation times were prolonged. This change occurred when the survival fraction reached a value of approximately 10^{-6} for *H. volcanii* (after 25 s, i.e. an energy of $\sim 1.4 \times 10^{30}$ erg), and the viable counts remained above the LOD up to 600 s (6000 J m^{-2} , $\sim 3.5 \times 10^{31}$ erg) (Fig. 1). In the case

of *P. aeruginosa* the slope changed after the same time/energy for the typical flare, and the viable counts remained above the LOD at all considered fluence values (Fig. 2). For the superflare (Fig. 3), the change of slope occurred at a survival fraction between 10^{-4} and 10^{-5} , i.e. after 28 s ($\sim 1.7 \times 10^{31}$ erg), and the viable counts remained above the LOD for all fluence values. This reduction of the slope of the survival curves, usually designated as ‘tail’, indicates the presence of minority subpopulations of cells able to tolerate high fluences. The reason for this phenomenon is not fully understood. It has been ascribed to the formation of clumps (Coohill & Sagripanti 2008), or alternatively to transitory and reversible phenotypic changes (Pennell, Aronson & Blatchey III 2008), similar to those responsible for persistence in bacteria exposed to bactericidal antibiotics (Lewis 2010) and in archaea during starvation conditions or that were treated with various biocidal compounds (Megaw & Gilmore 2017).

Considering the shape of the survival curves shown in Figs 1–3, it seems possible that a small part of a microbial population irradiated during a flare survives. Eventually, survivors could proliferate, reestablishing the microbial population, and therefore the occurrence of a flare would not imply the extinction of the exposed microorganisms. Interestingly, subpopulations able to tolerate high-UVC fluences were found even in cultures of a microorganism that can be considered ubiquitous on the Earth surface, like *P. aeruginosa*, exposed to a superflare in a ‘worst case scenario’ (Fig. 3). Since survival curves to UVC are frequently biphasic, the effect produced by high fluences, like those expected during flares or superflares, cannot be appraised from the initial slope and from the total fluence imparted, such as previous studies assumed by using the UV-LD. For instance, some of them predicted no survival to superflares taking as parameter for lethality fluences greater than 553 J m^{-2} (Estrela & Valio 2018; Howard et al. 2018), which is, according to our experiments, roughly 70 times smaller than the fluence to produce a significant decrease in the viability of the microorganisms on Proxima *b* during a superflare. Moreover, previous theoretical studies using data from curves of UV-BAS for microorganisms (Rugheimer et al. 2015; O’Malley-James & Kaltenecker 2017; Estrela & Valio 2018; O’Malley-James & Kaltenecker 2019) or DNA molecules *in vitro* (Buccino et al. 2006, 2007; Rugheimer et al. 2015) are missing important information, as they work with UV-BAS usually obtained at low fluences in contrast to those related to flares or superflares. Moreover, when using UV-BAS obtained from DNA molecules there is an additional problem. The results can be unrealistic, because the overall biological effect of UVR will depend not only on the damage to DNA, but also to other cellular structures, as well as on the intrinsic mechanisms of the cell, such as photoprotection and DNA repair. Among these mechanisms, microorganisms could use the energy of the visible light to repair UV-induced DNA damage (i.e.: photoreactivation mechanisms, 2.3), improving their survival (Sancar 2000). We tested this mechanism in our experiments, for the case of *P. aeruginosa* exposed to the superflare, and no significant increase in the survival rate was found when the irradiated bacteria were exposed to visible light in comparison to those incubated in the dark (Fig. 3, inset box).

4 CONCLUSIONS

We studied the UV fluxes on the surface of Proxima *b* to evaluate if this kind of radiation could be a factor to limit the existence of ‘life as we know it’. We demonstrate that other planetary atmospheric compositions than O-rich ones (namely N_2 and CO_2)

can be optically thick to UV, in particular for the most biologically damaging wavelengths (UVC and UVB). Moreover, UV shielding increases as a function of atmospheric pressure. This expands the possible atmospheres to be considered as ‘UV-protective’ in habitability studies, beyond O₂ – ozone rich atmospheric compositions. Nevertheless, our results show that the UV fluence rates on the surface, considering the star in quiescence, would probably have negligible effects on life even without an atmosphere. Given that, we focused our study on a ‘worst case scenario’, considering therefore flare activity in the absence of any sources of UV-shielding, such as an atmosphere. Our experiments demonstrate, for the first time, the potential impact that the UVR from a flare would have on life: when microorganisms receive very high fluences of UVC, such as the amount of radiation expected to reach the surface of Proxima *b* after a typical flare or a superflare, a fraction of the microbial population is able to survive. How the flare frequency (e.g. Davenport et al. 2016; Vida et al. 2019) would affect the survival of the microbial population remains an open question and will be addressed in a future study. Our present results based on a single flare event show that microorganisms have higher chances to cope with UVR on the surface of Proxima *b* than estimated in previous studies based on modelling. Beyond the conditions considered in our study, given the potential atmospheric composition and the inherent mechanisms of defence of potential microorganisms against UVR, the probability of life to persist in highly UV irradiated environments could be increased by other biological factors such as clumping and biofilm formation, as well as other environmental factors apart from the atmosphere such as turbidity or the presence of UV-absorbing compounds in a liquid medium or a water column like in an ocean. Our study demonstrates that life can cope with very unfavourable conditions that cannot be found naturally on our planet. These results can be extended to the UVR surface habitability of other terrestrial planets orbiting M dwarfs. Additionally, they can also be relevant to the study of potential surface biosignatures on other planetary bodies.

ACKNOWLEDGEMENTS

We thank the reviewer, Paul Rimmer, for his valuable comments on this paper. We also thank Dr. H. Lammer for helpful discussions at the beginning of this programme and Dr. S. Ranjan for providing data for model comparisons. XCA acknowledges the Austrian Academy of Sciences (ÖAW) for the Joint Excellence in Science & Humanities fellowship grant, and partial funding by Proyecto de Unidad Ejecutora IAFE (CONICET) (2016–2021). AFFG acknowledges funding by Universidad Nacional de Tres de Febrero. ML acknowledges the Austrian Science Fund (FWF): P30949-N36. We also acknowledge funding from UKRI/STFC and data presented here can be made available on request.

REFERENCES

Abrevaya X. C., Thomas B. C., 2018, in Gordon R., Sharov A. A., eds, *Habitability of the Universe Before Earth* [in series: *Astrobiology: Exploring Life on Earth and Beyond*], Elsevier B.V., Amsterdam

Abrevaya X. C., Cortón E., Mauas P. J. D., 2011/2, *Proc. IAU Symp.* 286: Comparative Magnetic Minima: Characterizing Quiet Times in the Sun and Stars. Cambridge Univ. Press, Cambridge, p. 405

Abrevaya X. C., Leitzinger M., Nuñez Pölcher P. N., Odert P., Lammer H., Hanslmeier A., 2014, Workshop ‘Star-Planet Interactions and the Habitable Zone’ – CEA – Irfu. France. Available online: <http://irfu.cea.fr/habitability/>

Abrevaya X. C., Leitzinger M., Oppezzo O. J., Odert P., Luna G. J. M., Patel M., Forte Giacobone A. F., Hanslmeier A., 2018/20, in Elmegreen B. G., Tóth L. V., Güdel M., eds, *Proc. IAU Symp.* 345, *Origins: From the Protosun to the First Steps of Life*. Kluwer, Dordrecht, p. 222

Anglada-Escudé G. et al., 2016, *Nature*, 536, 437

Braslavsky S. E., 2007, *Pure Appl. Chem.*, 79, 293

Buccino A. P., Lemarchand G. A., Mauas P. J. D., 2006, *Icarus*, 183, 491

Buccino A. P., Lemarchand G. A., Mauas P. J. D., 2007, *Icarus*, 192, 582

Cockell C. S., 1999, *Icarus*, 141, 399

Cockell C. S., Rettberg P., Rabbow E., Olsson-Francis K., 2011, *ISME J.*, 5, 1671

Coohill T. P., Sagripanti J. L., 2008, *Photochem. Photobiol.*, 84, 1084

Davenport J. R. A., Kipping D. M., Sasselov D., Matthews J. M., Cameron C., 2016, *ApJ*, 829, L31

Estrela R., Valio A., 2018, *Astrobiology*, 18, 1414

Fossati L., Bagnulo S., Haswell C. A., Patel M. R., Busuttill R., Kowalski P. M., Shulyak D. V., Sterzik M. F., 2012, *ApJ*, 757, L15

Gao P., Hu R., Robinson T. D., Li C., Yung Y. L., 2015, *ApJ*, 806, 249

Garcia-Sage K., Gloer A., Drake J. J., Gronoff G., Cohen O., 2017, *ApJ*, 844, L13

Gershberg R. E., 2005, *Solar-Type Activity in Main-Sequence Stars*, Springer, Berlin Heidelberg

Hanslmeier A., 2018, *Planetary Habitability and Stellar Activity. Advances in Planetary Science: Vol. 3*. World Scientific Publishing Company, Singapore, New York, London

Hawley S. L., Pettersen B. R., 1991, *ApJ*, 378, 725

Horneck G., Bucker H., Reitz G., 1994, *Adv. Space Res.*, 14, 41

Howard W. S. et al., 2018, *ApJ*, 860, L30

Husser T.-O., Wende-von Berg S., Dreizler S., Homeier D., Reiners A., Barman T., Hauschildt P. H., 2013, *A&A*, 553, A6

Johnstone C. P., Güdel M., Lammer H., Kislyakova K. G., 2018, *A&A*, 617, A107

Johnstone C. P., Khodachenko M. L., Lüftinger T., Kislyakova K. G., Lammer H., Güdel M., 2019, *A&A*, 624, L10

Kielbassa C., Roza L., Epe B., 1997, *Carcinogenesis*, 18, 811

Lammer H., Lichtenegger H. I. M., Khodachenko M. L., Kulikov Y. N., Griessmeier J., 2011, in Beaulieu J. P., Dieters S., Tinetti G., eds, *ASP Conf. Ser. Vol. 450, Molecules in the Atmospheres of Extrasolar Planets*. Astron. Soc. Pac., San Francisco, p. 139

Lewis K., 2010, *Annu. Rev. Microbiol.*, 64, 357

Lichtenegger H. I. M., Lammer H., Griessmeier J., Kulikov Y. N., von Paris P., Hausleitner W., Krauss S., Rauer H., 2010, *Icarus*, 210, 1

Meadows V. S. et al., 2018, *Astrobiology*, 18, 133

Meftah M. et al., 2018, *A&A*, 611, A1

Megaw J., Gilmore B. F., 2017, *Front. Microbiol.*, 8, 1589

Novikova N., Deshevaya E., Levinskikh M., Polikarpov N., Poddubko S., Gusev O., Sychev V., 2015, *Int. J. Astrobiol.*, 14, 137

O’Malley-James J. T., Kaltenecker L., 2017, *MNRAS*, 469, L26

O’Malley-James J. T., Kaltenecker L., 2019, *MNRAS*, 485, 5598

Oppezzo O., Costa C. S., Pizzarro R. A., 2011, *Trends Photochem. Photobiol.*, 13, 37

Patel M. R., Zarnacki J. C., Catling D. C., 2002, *Planet. Space Sci.*, 50, 915

Pennell K. G., Aronson A. I., Blatchey E. R., III, 2008, *J. Appl. Microbiol.*, 84, 1084

Ranjan S., Wordsworth R., Sasselov D. D., 2017, *ApJ*, 843, 110

Rugheimer S., Segura A., Kaltenecker L., Sasselov D., 2015, *ApJ*, 806, 137

Sancar G. B., 2000, *Mutat. Res.*, 451, 25

Segura A., Walkowicz L. M., Meadows V., Kasting J., Hawley S., 2010, *Astrobiology*, 10, 751

Tian F., Kasting J. F., Liu H., Roble R. G., 2008, *J. Geophys. Res.*, 113, E05008

Vida K., Oláh K., Kóvári Z., van Driel-Gesztelyi L., Moór A., Pál A., 2019, *ApJ*, 884, 160

Welsh B. Y. et al., 2007, *ApJS*, 173, 673

SUPPORTING INFORMATION

Supplementary data are available at [MNRASL](#) online.

Appendix A. Additional Information.

Please note: Oxford University Press is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

APPENDIX: ADDITIONAL INFORMATION

Figs A1, A2, and B, and details about methods are available online.

This paper has been typeset from a \TeX/L\AA\TeX file prepared by the author.