Open Research Online

The Open University's repository of research publications and other research outputs

Wiman–Valiron Discs and the Dimension of Julia Sets

Journal Item

How to cite:

Waterman, James (2021). Wiman–Valiron Discs and the Dimension of Julia Sets. International Mathematics Research Notices, 2021(12), article no. rnaa029.

For guidance on citations see [FAQs](#).

© 2020 The Authors

<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Version: Accepted Manuscript

Link(s) to article on publisher's website:
<http://dx.doi.org/doi:10.1093/imrn/rnaa029>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

WIMAN–VALIRON DISCS AND THE DIMENSION OF JULIA SETS

JAMES WATERMAN

ABSTRACT. We show that the Hausdorff dimension of the set of points of bounded orbit in the Julia set of a meromorphic map with a simply connected direct tract and a certain restriction on the singular values is strictly greater than one. This result is obtained by proving new results related to Wiman–Valiron theory.

1. INTRODUCTION

Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be a transcendental meromorphic function and denote the n th iterate of f by f^n for $n = 1, 2, 3, \dots$. The *Fatou set* $F(f)$, is defined to be the subset of $\mathbb{C}$ where the iterates f^n of f form a normal family. The *Julia set* is defined to be its complement. Finally, the *escaping set* $I(f)$ is the set of $z \in \mathbb{C}$ for which $f^n(z) \rightarrow \infty$ as $n \rightarrow \infty$ and we denote by $K(f)$ those points in $\mathbb{C}$ with bounded orbit. An introduction to the Fatou set, Julia set, and escaping set and their properties can be found in the survey article of Bergweiler, see [4]. More information on $K(f)$ can be found in [6] and [21].

The Hausdorff dimension of the Julia set, $\dim J(f)$, of transcendental entire and meromorphic functions has been widely studied. Misiurewicz [19] proved that the Julia set of the exponential function is the entire plane, and hence $\dim J(f) = 2$ in this case. Major work was done by McMullen [18], who gave examples for which $\dim J(f) \cap I(f) = 2$ but the Julia set is not the entire plane. It was shown by Baker [1] that the Julia set of every transcendental entire function contains a continuum and hence $\dim J(f) \geq 1$. It was further shown by Stallard [27] that for each $d \in (1, 2)$ there exists a transcendental entire function for which $\dim J(f) = d$.

More is known if one restricts to a specific class of entire functions. Many results have been proven for functions in the Eremenko–Lyubich class $\mathcal{B}$. For example, if f is a function of finite order in the class $\mathcal{B}$, then $I(f) (\subset J(f))$ has Hausdorff dimension 2; see [2] and [25]. Stallard [26] proved that for all entire functions in the Eremenko–Lyubich class $\mathcal{B}$, $\dim J(f) > 1$. This was then improved by Barański, Karpińska, and Zdunik [3] who showed that, for a meromorphic function with a logarithmic tract, the Hausdorff dimension of the set of points in the Julia set with bounded orbit is strictly greater than 1. While Stallard’s result made use of escaping points, Barański, Karpińska, and Zdunik used completely different methods of proof based on points in $J(f) \cap K(f)$.

Until recently it was an open question as to whether there existed an entire function f for which the Hausdorff dimension of the Julia set of f was equal to 1. However, Bishop [10] has constructed an entire function f with $\dim J(f) = 1$. This function has multiply connected wandering domains, and so its direct tract, defined in Section 3, does not have an unbounded boundary component. This suggests the question of whether Barański, Karpińska, and Zdunik’s result can be generalized to show that, for a transcendental meromorphic function f with a direct tract with an unbounded boundary component, we have that $\dim J(f) > 1$. We prove this in several situations, and specifically in the case where the direct tract is simply connected and there is an additional restriction on the singular values associated with the direct tract.

Theorem 1.1. *Let f be a transcendental meromorphic function with a simply connected direct tract D . Suppose that there exists $\lambda > 1$ such that for arbitrarily large r there exists an annulus $A(r/\lambda, \lambda r)$ containing no singular values of the restriction of f to D . Then $\dim J(f) \cap K(f) > 1$.*

Note that there are relatively few results concerning the set $K(f)$ for transcendental entire functions. Following the proof of [3], Bergweiler showed that $\dim J(f) \cap K(f)$ is not greater than one in general. Indeed, he proved in [6] that there exist transcendental entire functions for which $\dim K(f)$ can be arbitrarily small. Note that Bergweiler’s examples all have direct tracts with no unbounded boundary component.

Our proof of Theorem 1.1 partially follows the approach in Barański, Karpińska, and Zdunik [3], with some significant differences. The hypotheses in [3] specify a logarithmic tract, for which one can make use of the logarithmic transform and the resulting expansion estimate in the tract. However, in our case, we no longer have this expansion estimate throughout the direct tract. Instead, our proof uses Wiman–Valiron theory, which gives the existence of a disc in the direct tract, around certain points where the function takes its maximum modulus, within which the function behaves like a polynomial. We use tools developed by Bergweiler, Rippon, and Stallard [9] for general direct tracts, and apply them to direct tracts with certain properties. These results trace back to Wiman–Valiron theory [14] and Macintyre’s theory of flat regions [17]. We show that in our case a larger disc than that given by Wiman–Valiron theory lies in the direct tract, and on this disc an estimate of the function holds which is somewhat weaker than the Wiman–Valiron estimate. While this estimate on the new disc is weaker than the Wiman–Valiron estimate, it enables us to cover a much larger annulus than does the Wiman–Valiron estimate many times over. These results are of independent interest.

The paper is organized as follows. Section 2 introduces the notion of a direct tract and the classification of singularities. In Section 3, we give our results on the size of enlarged

Wiman–Valiron discs. We then use these results in Section 4 in order to prove a theorem on the Hausdorff dimension of certain Julia sets, of which Theorem 1.1 is a special case. Finally, in Section 5 we give two nontrivial examples of transcendental entire functions to which we can apply our results.

2. DIRECT TRACTS AND THE LOGARITHMIC TRANSFORM

First, we recall the classification of the singularities of the inverse function due to Iversen [15], as well as the definition of a tract, following terminology found in [9]. Let f be a meromorphic function and consider $a \in \widehat{\mathbb{C}}$. For $R > 0$, let U_R be a component of $f^{-1}(D(a, R))$ (where $D(a, R)$ is the open disc centered at a with radius R with respect to the spherical metric) chosen so that $R_1 < R_2$ implies that $U_{R_1} \subset U_{R_2}$. Then either $\bigcap_R U_R = \{z\}$ for some unique $z \in \mathbb{C}$ or $\bigcap_R U_R = \emptyset$.

In the first case, $a = f(z)$ and a is an *ordinary point* if $f'(z) \neq 0$, or a is a *critical value* if $f'(z) = 0$. If $f'(z) = 0$, we call z a *critical point*. In the second case, f has a *transcendental singularity* over a . The transcendental singularity is called *direct* if $f(z) \neq a$ for all $z \in U_R$, for some $R > 0$. Otherwise it is *indirect*. Further, a direct singularity is called *logarithmic* if $f : U_R \rightarrow D(a, R) \setminus \{a\}$ is a universal covering. These components U_R are called *tracts* for f . Note that the structure of U_R depends on R . We will mainly restrict f to a specific tract, which motivates the following definition; see for example [9].

Definition 2.1. Let D be an unbounded domain in $\mathbb{C}$ whose boundary consists of piecewise smooth curves and suppose that the complement of D is unbounded. Further, let f be a complex valued function whose domain of definition contains the closure $\overline{D}$ of D . Then D is called a *direct tract* of f if f is holomorphic in D , continuous in $\overline{D}$, and if there exists $R > 0$ such that $|f(z)| = R$ for $z \in \partial D$ while $|f(z)| > R$ for $z \in D$. If, in addition, the restriction $f : D \rightarrow \{z \in \mathbb{C} : |z| > R\}$ is a universal covering, then D is called a *logarithmic tract*. Further, we call R the *boundary value* of the direct tract.

Of note is that meromorphic functions with a direct singularity have a direct tract. Further, every transcendental entire function has a direct tract. Although, for a meromorphic function, the boundaries of these tracts are piecewise analytic, the geometry of these tracts can be incredibly varied and sometimes quite wild. Direct tracts need not be simply connected or even have an unbounded boundary component. However, logarithmic tracts are always simply connected and direct tracts in the class $\mathcal{B}$ are logarithmic for a sufficiently large R in the above.

Logarithmic tracts have been of great use in the iteration of entire and meromorphic functions with many major results and constructions such as [24] and [9] making use of their properties and the additional tools that they give. A technique to identify a logarithmic tract

was given in [31]. The major tool used is the logarithmic transform which was first studied in the context of entire functions by Eremenko and Lyubich [12]. We now introduce the logarithmic transform, F , in the setting of a direct tract with an unbounded boundary component. While the logarithmic transform has been studied in many papers on transcendental entire dynamics, it has almost always been applied to a logarithmic tract.

Let f be a meromorphic function with a direct tract D with an unbounded complementary component; any simply connected direct tract must have such a complementary component. By performing a translation if necessary, assume further that 0 is contained in an unbounded complementary component of D . Then, denote by F the logarithmic transform of f ; that is, $\exp \circ F = f \circ \exp$. Note that F is periodic with period $2\pi i$ and maps each component of $\log D$ into a right half-plane. Unlike for a logarithmic tract, F is not necessarily univalent on each component of $\log D$. However, as there exists an unbounded complementary component, one may still lift f by the branches of the logarithm.

By applying the Koebe 1/4-theorem, we are able to obtain an estimate on the entire function in the direct tract on a domain on which the function is univalent. This result is a modification of an expansion estimate due to Eremenko and Lyubich [12] and is similar to the approach of Rippon and Stallard in [23, Lemma 2.5].

Lemma 2.1. *Let f be a transcendental meromorphic function with a direct tract D , and let $\lambda > 1$. Suppose, for some $r > 0$, that the restriction of f to D has no singular values in $A(r/\lambda, \lambda r)$ and $\gamma \subset D$ is a simple unbounded curve on which $|f(z)| = r$. Then,*

- (a) *the function $\Phi(z) = \log f(z)$, for $z \in D$, is univalent on a simply connected domain Ω such that $\gamma \subset \Omega \subset D$, and*

$$\Phi(\Omega) = S = \{w : \log(r/\lambda) < \operatorname{Re} w < \log \lambda r\};$$

- (b) *the function $F(w) = \Phi(e^w) = \log f(e^w)$ maps each component of $\log \Omega$ univalently onto S ;*

- (c) *any analytic branch H of $F^{-1} : S \rightarrow \log D$ satisfies*

$$(1) \quad |H'(w)| \leq \frac{4\pi}{\operatorname{dist}(w, \partial S)}, \quad \text{for } w \in S,$$

or equivalently

$$(2) \quad \left| \frac{zf'(z)}{f(z)} \right| \geq \begin{cases} \frac{1}{4\pi} \log |\lambda r / f(z)|, & \text{for } |f(z)| \geq r \\ \frac{1}{4\pi} \log |\lambda f(z) / r|, & \text{for } |f(z)| \leq r, \end{cases}$$

for $z \in \Omega$.

Proof. Take a base point $z_0 \in \gamma$ and let $w_0 = \log f(z_0)$. Then, let Ψ denote the inverse branch of $z \mapsto \Phi(z) = \log f(z)$ such that $\Psi(w_0) = z_0$. By our hypothesis on the singular

values of f , the branch Ψ can be analytically continued along every curve in S starting from w_0 , with image values in D . Since S is simply connected, we deduce from the monodromy theorem that Ψ extends to be analytic throughout S , and $\Omega := \Psi(S) \subset D$.

Two cases can then arise [20, p. 283]: either Ψ is univalent in S or Ψ is periodic in S , with period $2\pi im$, where $m \in \mathbb{N}$. In the latter case, however, $\Psi(S)$ must be bounded, which is impossible, since $\gamma \subset \Psi(S) = \Omega$.

Now, we use the fact that $H = \log \Psi$, for some branch of the logarithm, and hence H maps S univalently onto a domain that contains no disc of radius greater than π . Therefore, (1) holds by the Koebe 1/4-theorem, and (2) follows by the change of variables $z = e^w$. $\square$

3. WIMAN–VALIRON THEORY IN TRACTS

First, we introduce some preliminary concepts, discuss the original Wiman–Valiron result of Bergweiler, Rippon, and Stallard, and give some basic lemmas; see [9].

Let $v : \mathbb{C} \rightarrow [0, \infty)$ be a non-constant subharmonic function. Then the function

$$B(r, v) = \max_{|z|=r} v(z)$$

is increasing, convex with respect to $\log r$, and $B(r, v)/\log r \rightarrow \infty$ as $r \rightarrow \infty$. So,

$$a(r, v) = \frac{dB(r, v)}{d \log r} = rB'(r, v)$$

exists except for possibly a countable set of r values and is non-decreasing.

Let f be a meromorphic function with a direct tract D with an unbounded boundary component and boundary value R . Define

$$(3) \quad v(z) = \log \frac{|f(z)|}{R},$$

for $z \in D$ and $v(z) = 0$ elsewhere; then $B(r, v) = \max_{|z|=r} \log |f(z)|/R$.

It was proved in [9] that any direct tract of f contains a disc about points where the maximum modulus is attained, on which f behaves asymptotically like a polynomial. As we will refer back to this result several times, we state it here.

Theorem 3.1 ([9], Theorem 2.2). *Let D be a direct tract of f and let $\tau > \frac{1}{2}$. Let v be defined as in (3) and let $z_r \in D$ be a point satisfying $|z_r| = r$ and $v(z_r) = B(r, v)$. Then there exists a set $E \subset [1, \infty)$ of finite logarithmic measure such that if $r \in [1, \infty) \setminus E$, then $D(z_r, r/a(r, v)^\tau) \subset D$. Moreover,*

$$f(z) \sim \left(\frac{z}{z_r} \right)^{a(r, v)} f(z_r), \quad \text{for } z \in D(z_r, r/a(r, v)^\tau),$$

as $r \rightarrow \infty$, $r \notin E$.

We now show that a larger disc than given by Theorem 3.1 is possible inside a simply connected direct tract. In a general direct tract this is not possible, as shown by an example due to Bergweiler [5]. Note that, Bergweiler's example has a direct tract with no unbounded boundary component.

First, we recall a standard estimate on $a(r, v)$; see [9, Lemma 6.10].

Lemma 3.1. *Let $v : \mathbb{C} \rightarrow [-\infty, \infty)$ be subharmonic and let $\epsilon > 0$. Then there exists a set $E \subset [1, \infty)$ of finite logarithmic measure such that*

$$a(r, v) \leq B(r, v)^{1+\epsilon}$$

for $r \geq 1$, $r \notin E$.

Next, we state a more general version of [9, Lemma 11.2], giving a slightly different estimate of $B(r, v)$ for a general subharmonic function which is suited for our purpose. This statement is extracted from the inequalities given in the proof of [9, Lemma 11.2] (in which the function $\Phi(x) = B(r, v)$ and $\Phi'(x) = a(r, v)$, with $r = e^x$).

Lemma 3.2. *Let $v \rightarrow [-\infty, \infty)$ be subharmonic and let $\beta > \alpha > 0$. Then there exists a set $E \subset [1, \infty)$ of finite logarithmic measure such that*

$$B(s, v) \leq B(r, v) + a(r, v) \log \frac{s}{r} + a(r, v)^{1-\alpha-\beta}, \quad \text{for } \left| \log \frac{s}{r} \right| \leq \frac{1}{a(r, v)^\beta}, r \notin E,$$

uniformly as $r \rightarrow \infty$.

We shall use Lemma 3.2 to prove a generalization of Theorem 3.1, which provides a much larger disc in a direct tract that satisfies certain extra conditions, at the expense of less control on the function as we move further out from the original Wiman–Valiron disc. First, if there exists a direct tract surrounding a sparse collection of zeros, we show that a larger disc exists inside the direct tract than given by Theorem 3.1. Note that this hypothesis is satisfied by a simply connected direct tract.

Theorem 3.2. *Let f be a meromorphic function with a direct tract D with an unbounded boundary component. Let v be defined as in (3), and let $z_r \in D$ be a point for which $|z_r| = r$ and $v(z_r) = B(r, v)$. Suppose that there exists $\lambda > 1$ and a set $E \subset [1, \infty)$ of finite logarithmic measure such that, for $r \in [1, \infty) \setminus E$, we have that $z_r \in L_r$, a simply connected component of $A(r/\lambda, \lambda r) \cap D$. Also, let $\frac{1}{2} > \tau > 0$. Then, for $r \in [1, \infty) \setminus E'$, where E' has finite logarithmic measure, there exists $D(z_r, r/a(r, v)^\tau) \subset D$.*

Proof. Choose α and β such that $0 < \alpha < \beta < \tau < 1/2$ and $1 - \alpha - \beta = \sqrt{1 - 2\tau} = \xi(\tau)$, say. Let E' be the union of E and the exceptional sets in Lemma 3.1 and Lemma 3.2 for these values of α and β , where $\epsilon = \beta$. Further, set $\rho = 2ra(r, v)^{-\tau}$, where r is large enough

that $a(r, v) \neq 0$. Consider the subharmonic function

$$(4) \quad u(z) = v(z) - B(r, v) - a(r, v) \log \frac{|z|}{r} \leq a(r, v)^{\xi(\tau)},$$

for $r \notin E'$ and $z \in \overline{D}(z_r, 512\rho)$ by Lemma 3.2, and the fact that for $z \in \overline{D}(z_r, 512\rho)$,

$$\left| \frac{z - z_r}{z_r} \right| \leq \frac{512\rho}{r} = \frac{1024}{a(r, v)^\tau} = o(1) \text{ as } r \rightarrow \infty,$$

since $\lim_{r \rightarrow \infty} a(r, v) = \infty$, and so, for large r ,

$$(5) \quad \left| \log \frac{|z|}{r} \right| = \left| \log \left| 1 + \frac{z - z_r}{z_r} \right| \right| \leq 2 \left| \frac{z - z_r}{z_r} \right| \leq \frac{2048}{a(r, v)^\tau} \leq \frac{1}{a(r, v)^\beta}.$$

Now, we show that $D(z_r, r/a(r, v)^\tau) \subset D$ for $r \notin E'$ sufficiently large. Following [9, p. 395–396], suppose not. First, by assumption, $D(z_r, 256\rho)$ does not contain any bounded complementary components, since the exceptional set of r has finite logarithmic length and L_r is assumed simply connected. Then, since D has an unbounded boundary component, there exists an unbounded component of the complement of D that intersects $\partial D(z_r, t)$ for $\rho \leq t \leq 256\rho$. Let V be the component of $D \cap D(z_r, 512\rho)$ that contains z_r and let $\Gamma = \partial V \cap \partial D(z_r, 512\rho)$. Note that V is simply connected, as L_r is simply connected. Further, in this case, let $t\theta^*(z_r, t)$ be the linear measure of the intersection of $\partial D(z_r, t)$ with D . Then, $\theta^*(z_r, t) \leq 2\pi$ for $\rho \leq t \leq 256\rho$, since $\partial D(z_r, t)$ is not wholly contained in D . Hence, by a result of Tsuji; see [28, p. 112],

$$\omega(z_r, \Gamma, V) \leq 3\sqrt{2} \exp \left(-\pi \int_\rho^{256\rho} \frac{dt}{t\theta^*(z_r, t)} \right) \leq 3\sqrt{2} \exp \left(-\frac{1}{2} \int_\rho^{256\rho} \frac{dt}{t} \right) = \frac{3\sqrt{2}}{2^4} < \frac{1}{2},$$

where $\omega(z_r, \Gamma, V)$ denotes the harmonic measure of Γ at z_r in V . Thus, for $\Sigma = \partial V \setminus \Gamma$ we have

$$\omega(z_r, \Sigma, V) = 1 - \omega(z_r, \Gamma, V) > \frac{1}{2}.$$

Next, for $z \in \Gamma$, we have $v(z) = 0$, since z is on the boundary of D , so from (4), (5), and Lemma 3.1 with $\epsilon = \beta$, if $r \notin E'$ is sufficiently large, then

$$\begin{aligned} u(z) &= -B(r, v) - a(r, v) \log \frac{|z|}{r} \\ &\leq -B(r, v) + a(r, v)^{1-\beta} \\ &\leq -B(r, v) + B(r, v)^{(1-\beta)(1+\epsilon)} \\ &\leq -\frac{1}{2}B(r, v). \end{aligned}$$

Hence, by (4) and the above, we can apply the two constant theorem (see for example [22, Theorem 4.3.7]) for large $r \notin E'$ and obtain

$$\begin{aligned}
u(z_r) &\leq -\frac{1}{2}\omega(z_r, \Sigma, V)B(r, v) + a(r, v)^{\xi(\tau)}(1 - \omega(z_r, \Sigma, V)) \\
&\leq -\frac{1}{4}B(r, v) + a(r, v)^{\xi(\tau)} \\
&\leq -\frac{1}{4}B(r, v) + B(r, v)^{\xi(\tau)(1+\epsilon)} \\
&\leq -\frac{1}{8}B(r, v),
\end{aligned}$$

which gives a contradiction, since $u(z_r) = 0$, while $B(r, v) \rightarrow \infty$ as $r \rightarrow \infty$. Hence, $D(z_r, r/a(r, v)^\tau) \subset D$ for $r \notin E'$ sufficiently large. $\square$

We now show that, given a disc of the form $D(z_r, r/a(r, v)^\tau)$, for $0 < \tau < 1/2$, inside our tract, we can obtain an estimate for the function f inside that disc. This estimate includes the case of the original Wiman–Valiron estimate when $\tau > 1/2$. A larger disc than that given by Wiman–Valiron theory introduces a worse error to the estimate the larger the new disc is. Note that for the following theorem, we do not insist on any additional hypotheses on the tract, such as being simply connected; we just require the existence of a suitably large disc inside the direct tract. Further, note that the function $\xi(\tau) = \sqrt{1 - 2\tau}$ is chosen so that $\xi(\tau) \rightarrow 1$ as $\tau \rightarrow 0$ and $\xi(\tau) \rightarrow 0$ as $\tau \rightarrow 1/2$, while also $\xi(\tau) < 1 - \tau$. These properties will be important in this and subsequent proofs.

Theorem 3.3. *Let f be a meromorphic function with a direct tract D and $\tau > 0$. Further, let v be defined as in (3), and let $z_r \in D$ be a point for which $|z_r| = r$ and $v(z_r) = B(r, v)$. Then there exists a set $E \subset [1, \infty)$ of finite logarithmic measure, such that, if there exists a disc $D(z_r, r/a(r, v)^\tau) \subset D$ for $r \notin E$ sufficiently large, then there exists an analytic function g in $D(z_r, r/a(r, v)^\tau)$ such that*

$$\log f(z) = \log f(z_r) + a(r, v) \log \frac{z}{z_r} + g(z), \quad \text{for } z \in D(z_r, r/a(r, v)^\tau),$$

where

$$g(z) = \begin{cases} O(a(r, v)^{\xi(\tau)}) & \text{for } z \in D(z_r, r/a(r, v)^\tau) \text{ and } \tau < 1/2, \\ o(1) & \text{for } z \in D(z_r, r/a(r, v)^\tau) \text{ and } \tau > 1/2, \end{cases}$$

and $\xi(\tau) = \sqrt{1 - 2\tau}$ as $r \rightarrow \infty$, $r \notin E$.

Proof. Set

$$(6) \quad g(z) = \log \frac{f(z)}{f(z_r)} - a(r, v) \log \frac{z}{z_r},$$

where the branches of the logarithms are chosen so that $g(z_r) = 0$. By the Borel–Carathéodory inequality [30, p. 20],

$$(7) \quad \max_{|z-z_r| \leq t} |g(z)| \leq 4 \max_{|z-z_r| \leq 2t} \operatorname{Re} g(z) \leq 4a(r, v)^{\xi(\tau)},$$

for $0 < t < \rho/2$ and $r \notin E$, by Lemma 3.2 with α and β chosen so that $\beta > \alpha > 0$ and $1 - \alpha - \beta = \xi(\tau)$.

Thus, by (6), for $z \in D(z_r, ra(r, v)^{-\tau})$ and $r \notin E$,

$$\log f(z) = \log f(z_r) + a(r, v) \log \frac{z}{z_r} + g(z), \quad \text{for } z \in D(z_r, r/a(r, v)^\tau),$$

where $g(z) = O(a(r, v)^{\xi(\tau)})$ for $\tau < 1/2$ by (7) and $g(z) = o(1)$ inside $D(z_r, r/a(r, v)^\tau)$ for $\tau > 1/2$ by Theorem 3.1. $\square$

Note that in the above result, the case $0 < \tau < 1/2$ is new. Further, for $\tau = 1/2$ it is natural to ask what the error, g , in the Wiman–Valiron estimate is and whether it is possibly $O(1)$. However, this proof does not show it.

We now use Theorem 3.3 in order to obtain an estimate on the size of the image of this new disc. First, we label three sets where we choose the principal branch of the logarithm; that is the branch corresponding to the principal value of the argument. Let

$$(8) \quad S_r = \left\{ w : |\operatorname{Re} w - \log r| \leq \frac{1/2}{a(r, v)^\tau}, |\operatorname{Im} w - \arg z_r| \leq \frac{1/2}{a(r, v)^\tau} \right\}.$$

Further, let

$$(9) \quad Q = \left\{ w : |\operatorname{Re} w - \log |f(z_r)|| < \frac{1}{8}a(r, v)^{1-\tau}, |\operatorname{Im} w - \arg f(z_r)| < \frac{1}{8}a(r, v)^{1-\tau} \right\},$$

and

$$(10) \quad \hat{Q} = \left\{ w : |\operatorname{Re} w - \log |f(z_r)|| < \frac{1}{4}a(r, v)^{1-\tau}, |\operatorname{Im} w - \arg f(z_r)| < \frac{1}{4}a(r, v)^{1-\tau} \right\}.$$

Theorem 3.4. *Let f be a meromorphic function with a direct tract D with an unbounded boundary component and τ , v , z_r , and E be as in Theorem 3.3. Consider the logarithmic transform F of f . If, for $r \in [1, \infty) \setminus E$, there exists $D(z_r, r/a(r, v)^\tau) \subset D$, then $S_r \subset \log D(z_r, r/a(r, v)^\tau)$ is mapped univalently by F and $F(S_r) \supset \hat{Q} \supset Q$.*

Proof. Let $w_r = \log z_r + i \arg z_r$, where we choose the principal branch of the logarithm and recall that F is the logarithmic transform of f . From Theorem 3.3,

$$(11) \quad F(w) = F(w_r) + a(r, v)(w - w_r) + O(a(r, v)^{\xi(\tau)}),$$

for $w \in \log D(z_r, r/a(r, v)^\tau)$ and as $r \rightarrow \infty$.

FIGURE 1. A sketch of the argument in the proof of Theorem 3.4.

The domain

$$S_r = \left\{ w : |\operatorname{Re} w - \log r| \leq \frac{1/2}{a(r, v)^\tau}, |\operatorname{Im} w - \arg z_r| \leq \frac{1/2}{a(r, v)^\tau} \right\}$$

$$\subset \log D(z_r, r/a(r, v)^\tau)$$

is a square of side length $1/a(r, v)^\tau$ centered at w_r and mapped by

$$w \mapsto F(w_r) + a(r, v)(w - w_r)$$

to a square, $\tilde{Q}$, of side $a(r, v)^{1-\tau}$ centered at $F(w_r)$. Since

$$F(w) = F(w_r) + a(r, v)(w - w_r) + O(a(r, v)^{\xi(\tau)}),$$

the image of ∂S_r under F is a closed curve lying entirely outside of a square of side

$$a(r, v)^{1-\tau} - O(a(r, v)^{\xi(\tau)})$$

centered at $F(w_r)$ for large enough r . This closed curve winds exactly once around $F(w_r)$, since $1 - \tau > \sqrt{1 - 2\tau} = \xi(\tau)$. Hence, by the argument principle, F is univalent on S_r and $F(S_r)$ contains a square of side length greater than $\frac{1}{2}a(r, v)^{1-\tau}$. $\square$

4. HAUSDORFF DIMENSION

In this section, we show that the set of points of bounded orbit in the Julia set of a meromorphic map with a direct tract with conditions on the singular values, zeros, and the boundary of the tract has Hausdorff dimension strictly greater than one. Theorem 1.1 follows from Theorem 4.1, below, together with Theorem 3.2, replacing λ by $\lambda/2$ in Theorem 4.1 and using the fact that the exceptional set has finite logarithmic length. Recall that v is defined as in (3), and z_r is a point for which $|z_r| = r$ and $v(z_r) = B(r, v)$.

FIGURE 2. Construction of J_B in Theorem 4.1.

Theorem 4.1. *Let f be a meromorphic map with a direct tract D with an unbounded boundary component. Fix $\lambda > 1$ and $\frac{1}{2} > \tau > 0$, and let E be the set of finite logarithmic measure in Theorem 3.3. Suppose that for arbitrarily large $r \in [1, \infty) \setminus E$ there exists an annulus $A(r/\lambda, \lambda r)$ such that both*

- (i) $D(z_r, r/a(r, v)^\tau) \subset A(r/\lambda, \lambda r) \cap D$, and
- (ii) $A(r/\lambda, \lambda r)$ contains no singular values of the restriction of f to D .

Then $\dim J(f) \cap K(f) > 1$.

Proof. First, let E be the set of finite logarithmic measure in Theorem 3.3. Fix $\frac{1}{2} > \tau > 0$ and $\lambda > 1$. Then, take $r \in [1, \infty) \setminus E$ so large that (i) and (ii) are satisfied. We use the sets S_r , Q , and $\hat{Q}$ introduced in (8), (9), and (10) before the statement of Theorem 3.4. By Theorem 3.4, the image of $S_r \subset D(z_r, r/a(r, v)^\tau)$ under $\log f$ covers the squares Q and $\hat{Q}$.

Following [3] we will construct an iterated function system based on taking inverse branches of F^2 that map Q to Q , where F is the $2\pi i$ periodic logarithmic transform of f which satisfies $\exp \circ F = f \circ \exp$; see Figure 2. We show that the dimension of the attractor of this system is strictly greater than 1. The method for constructing this iterated function system is, however, somewhat different from that of Barański, Karpińska, and Zdunik, and we make use of the tools from Wiman–Valiron theory developed in Section 3.

Step 1. We use our results from Section 3 to construct branches of F^{-1} from Q to $\log D(z_r, r/a(r, v)^\tau)$. We let $w_r = \log z_r + i \arg z_r$, where $\arg$ is the principal argument. We now let $w_{r,s} = w_{r,0} + 2\pi i s$, for $s \in \mathbb{Z}$, where $w_{r,0} = w_r$, and let $S_{r,s} = S_r + 2\pi i s$, for $s \in \mathbb{Z}$. Further, let G_s be the branch of F^{-1} satisfying $G_s(F(w_{r,0})) = w_{r,s}$. Then, $G_s : \hat{Q} \rightarrow S_{r,s}$ is univalent, since F is univalent on $S_{r,s}$ and $F(S_{r,s}) \supset \hat{Q} \supset Q$, by Theorem 3.4. By (11) and

by Cauchy's estimate

$$F'(w) = a(r, v) + O(a(r, v)^{\xi(\tau)+\tau}) = a(r, v)(1 + o(1)), \text{ for } w \in S_{r,s}, F(w) \in Q, \text{ and } s \in \mathbb{Z},$$

as $r \rightarrow \infty$. Hence there exists an absolute constant $K > 0$ such that

$$(12) \quad \max_{w \in Q} |(G_s)'(w)| < \frac{K}{a(r, v)}, \text{ for } s \in \mathbb{Z}.$$

Note that $F'(w_{r,s}) = a(r, v)$, and so

$$(13) \quad |(G_s)'(F(w_{r,s}))| = \frac{1}{a(r, v)}, \text{ for } s \in \mathbb{Z}.$$

Step 2. Next, using similar methods to [3], we construct another family of branches of F^{-1} . First, choose r so large that $\max_{w \in l} \operatorname{Re} F(w) > \log r$, where $l = \{w : \operatorname{Re} w = \log r\}$. This is possible since

$$\frac{\max_{w \in l} \operatorname{Re} F(w)}{\log r} = \frac{B(r, v)}{\log r} \rightarrow \infty$$

as $r \rightarrow \infty$. Since D has an unbounded boundary component with boundary value R , any domain of the form $\{z \in D : |f(z)| > R'\}$, for $R' > R$, must also have an unbounded boundary component. Thus we can choose an unbounded curve l_0 , say, in $\log D$ on which $\operatorname{Re} F(w) = \log r$ and $l_0 \cap \{w : \operatorname{Re} w < \log r\} \neq \emptyset$. Then, by Lemma 2.1 part (b), $F(l_0) = l$. Let $\zeta_0 \in l_0$ with $F(\zeta_0) \in l$ and let $\zeta_u = \zeta_0 + 2\pi i u$, for $u \in \mathbb{Z}$. Finally, let H_u be the branch of F^{-1} that maps $F(\zeta_0)$ to ζ_u , as given by Lemma 2.1 part (b).

Since there exist no singular values of f restricted to D that lie in $A(r/\lambda, \lambda r)$, we can obtain the following estimate on the derivative of H_u by applying Lemma 2.1 part (c) for $w \in \{w : \log(r/\sqrt{\lambda}) < \operatorname{Re} w < \log \sqrt{\lambda} r\}$:

$$(14) \quad \max_{w \in S_{r,s}} |(H_u)'(w)| < \frac{4\pi}{\log(\lambda r) - \log(\sqrt{\lambda} r)} = \frac{4\pi}{\log \sqrt{\lambda}}.$$

Step 3. Now, we estimate the diameter of $Q_{u,s} = H_u \circ G_s(Q)$, a second preimage of Q . Recall that $G_s(Q) \subset S_{r,s}$, so by (12) and (14),

$$\max_{w \in Q} |(H_u \circ G_s)'(w)| < \frac{4\pi K}{a(r, v) \log \sqrt{\lambda}}.$$

So, there exists $d \in (0, 1)$ such that

$$(15) \quad \operatorname{diam} Q_{u,s} < \max_{w \in Q} |(H_u \circ G_s)'(w)| \operatorname{diam} Q < \frac{4\pi\sqrt{2}Ka(r, v)^{1-\tau}}{4a(r, v) \log \sqrt{\lambda}} < d,$$

for r sufficiently large.

Let

$$Q' = \left\{ w : |\operatorname{Re} w - \log |f(z_r)|| < \frac{1}{8}a(r, v)^{1-\tau} - d, |\operatorname{Im} w - \arg f(z_r)| < \frac{1}{8}a(r, v)^{1-\tau} - d \right\}$$

and

$$Q'' = \left\{ w : |\operatorname{Re} w - \log |f(z_r)|| < \frac{1}{16}a(r, v)^{1-\tau}, |\operatorname{Im} w - \arg f(z_r)| < \frac{1}{16}a(r, v)^{1-\tau} \right\},$$

where d satisfies (15).

Step 4. We now consider level curves of F that meet Q'' . Let l_u be the image under H_u of $l = \{w : \operatorname{Re} w = \log r\}$, that is a translate of the curve l_0 introduced in Step 2. Denote by $l_{u, Q'}$ the intersection of l_u and Q' . From Step 2, we know that l_u is unbounded, and so if $l_{u, Q'} \cap Q'' \neq \emptyset$, then

$$(16) \quad \text{length}(l_{u, Q'}) \geq \frac{1}{16}a(r, v)^{1-\tau} - d.$$

Note that since $l_0 \cap l \neq \emptyset$ (from Step 2) there are at least $a(r, v)^{1-\tau}/16\pi$ values of u such that $l_u \cap Q'' \neq \emptyset$.

Step 5. Next, we obtain an estimate on $\sum_u \sum_s |(H_u \circ G_s)'(w)|$ for $w \in Q$, and u and s such that $H_u \circ G_s(Q) = Q_{u, s} \subset Q$. Now, $l_{u, Q'} \subset H_u(l)$. Since H_u is univalent on $\{w : \log(r/\lambda) < \operatorname{Re} w < \log(\lambda r)\}$, it follows from the Koebe distortion theorem (applied to a covering of l by disks of a uniform size) that there exists a constant $C_2 > 0$ such that

$$(17) \quad \text{length}(l_{u, Q'}) < C_2 \sum_s |(H_u)'(w_{r, s})|,$$

for any u such that $l_{u, Q'} \cap Q'' \neq \emptyset$, summing over all s such that $H_u(w_{r, s}) \in l_{u, Q'}$. Note that if $H_u(w_{r, s}) \in l_{u, Q'}$, then $Q_{u, s} \subset Q$ by (15).

The distortion of $H_u \circ G_s$ is uniformly bounded on Q , since $G_s : \hat{Q} \rightarrow S_{r, s}$ is univalent on $\hat{Q} \supset Q$, and H_u is univalent on $\{w : \log(r/\lambda) < \operatorname{Re} w < \log(\lambda r)\}$. Therefore, there exists a constant $C > 0$ such that

$$|(H_u \circ G_s)'(w)| > \frac{1}{C} |(H_u \circ G_s)'(F(w_{r, s}))|, \quad \text{for } w \in Q.$$

So, by (13), (16), and (17), for $w \in Q$, and u and s such that $Q_{u, s} \subset Q$,

$$\begin{aligned} \sum_s |(H_u \circ G_s)'(w)| &> \frac{1}{C} \sum_s |(H_u \circ G_s)'(F(w_{r, s}))| \\ &= \frac{1}{Ca(r, v)} \sum_s |(H_u)'(w_{r, s})| \\ &> \frac{\frac{1}{16}a(r, v)^{1-\tau} - d}{CC_2a(r, v)} \\ &> \frac{a(r, v)^{(1-\tau)}}{C_3a(r, v)}, \end{aligned}$$

for r sufficiently large, where $C_3 > 0$ is some constant. As we noted at the end of Step 4, there exist at least $a(r, v)^{1-\tau}/16\pi$ curves l_u that meet Q'' . Hence, for $w \in Q$, and u and s

such that $Q_{u,s} \subset Q$,

$$(18) \quad \sum_u \sum_s |(H_u \circ G_s)'(w)| > \frac{a(r, v)^{2(1-\tau)}}{C_3 16\pi a(r, v)}.$$

Step 6. Following [3, p. 622 - 623], we now obtain a conformal iterated function system from the family of maps $H_u \circ G_s : Q \rightarrow Q$. The sets $H_u \circ G_s(Q) = Q_{u,s}$ are pairwise disjoint and the system gives a maximal compact invariant set, J_B , say. Note that by construction $F^2(J_B) = J_B$ and $F(J_B) \subset \cup_{s \in \mathbb{Z}} D_s$. Further, $\lim_{n \rightarrow \infty} |(F^n)'(w)| = \infty$ for every $w \in J_B$ by (12) and (14).

Recall that the Hausdorff dimension of J_B is the unique zero of the pressure function

$$P(t) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \sum_{g^n} \|(g^n)'\|^t,$$

where $g^n = g_{i_1} \circ \dots \circ g_{i_n}$, with $g_{i_j} = H_u \circ G_s$ for u and s such that $Q_{u,s} \subset Q$. Further recall that the pressure function is strictly decreasing, so in order to prove that $\dim J_B > 1$ it is sufficient to show that $P(1) > 0$. For an introduction to the pressure function see [11] and [16].

Now, estimating the pressure using (18),

$$\begin{aligned} P(1) &= \lim_{n \rightarrow \infty} \frac{1}{n} \log \sum_{g^n} \|(g^n)'\| \\ &\geq \lim_{n \rightarrow \infty} \frac{1}{n} \log \left(\inf_{w \in Q} \sum_{(u,s)} |(H_u \circ G_s)'(w)| \right)^n \\ &\geq \log \left(\frac{a(r, v)^{2(1-\tau)}}{C_3 16\pi a(r, v)} \right) \\ &> 0, \end{aligned}$$

for r sufficiently large, since, by hypothesis, $\tau < 1/2$. This implies that the Hausdorff dimension of the invariant set J_B that arises from this system is greater than 1.

Let $X = \exp(J_B \cup F(J_B))$. As $\exp \circ F = f \circ \exp$ and $J_B \cup F(J_B)$ is F -invariant, X is f -invariant. Further, the exponential function is a smooth covering map, so $\dim X = \dim J_B \cup F(J_B) > 1$. Finally, we have that $\lim_{n \rightarrow \infty} |(F^n)'(w)| = \infty$ for $w \in J_B$ by (12) and (14). Therefore, $\lim_{n \rightarrow \infty} |(f^n)'(z)| = \infty$ for $z \in X$. However, $f^n(X)$ is bounded, so by normality, $X \subset J(f)$. Therefore, $\dim J(f) \cap K(f) > 1$. $\square$

5. EXAMPLES

In this section, we give two examples in order to illustrate the application of our results. First, we consider a function f with one direct tract, which is not simply connected. However,

FIGURE 3. The direct tract of $\cos(z)\exp(z)$ in white with boundary value 1 and zeros at $\pi/2 + k\pi$ for $k \in \mathbb{Z}$. Its boundary includes the point 0.

this function satisfies the hypotheses of Theorem 4.1 and so the Hausdorff dimension of $J(f) \cap K(f)$ is strictly greater than one.

Example 5.1. Let

$$f(z) = \cos(z)\exp(z).$$

Then $\dim J(f) \cap K(f) > 1$.

Consider the direct tract D of f with boundary value 1; see Figure 3. Then D contains $\{iy : y \neq 0\}$. Note that the only zeros of f are real and the only finite asymptotic value of f is 0, by the Denjoy-Carleman-Ahlfors theorem [20, XI.4], since f has order 1 and is symmetric in the real axis. We show that the points where f attains its maximum in the upper half-plane lie asymptotic to the line $y = x$ and that D contains a disc satisfying Theorem 4.1.

We use the following notation and results of Tyler [29] on maximum modulus curves; that is curves on which $|f(z)| = M(|z|)$. First, let

$$A(z) = \frac{zf'(z)}{f(z)} \text{ and } B(z) = zA'(z).$$

Further, let

$$b(r) = \frac{d^2 \log M(r)}{d(\log r)^2}.$$

Then, by [29, p. 2562] we have the following two properties:

- (a) $|f(z)| = M(r) \implies A(z)$ is real and positive, for $|z| = r$,
- (b) $|f(z)| = M(r) \implies B(x) = b(r) \geq 0$, for $x = r$.

For $f(z) = \cos(z) \exp(z)$, we have that $A(z) = z(1 - \tan z)$. Now, $A(z)$ is real and positive either in various intervals on the real line or asymptotically close to the line $y = x$, since $\tan(z) \rightarrow i$ as $\operatorname{Im} z \rightarrow \infty$.

By considering $B(x)$, we show that there are no large maximum modulus points on the positive real axis. First,

$$B(x) = x(1 - \tan x - x \sec^2 x) = x(1 - \tan x - x(1 + \tan^2 x)).$$

Hence, for $x > 0$,

$$B(x) \geq 0 \iff x \leq \frac{1 - \tan x}{1 + \tan^2 x} \iff x \leq 0.43 \dots,$$

approximately. Therefore, there are no maximum modulus points on the positive real axis greater than 0.44. So, the points satisfying both (a) and (b) in the upper half-plane must lie on an unbounded curve in the upper half-plane asymptotically close to the line $y = x$.

Now, all the zeros of f lie on the real axis. Since,

$$|f(x + iy)| = e^x (\cos^2 x + \sinh^2 y)^{\frac{1}{2}} \geq e^x \sinh y > 1, \quad \text{for } x > 0 \text{ and } |y| \geq 1,$$

the direct tract D contains $\{x + iy : x > 0, |y| \geq 1\}$.

Hence for $\tau > 0$ there exists a disc of radius $r^{1-\tau}$ with center on the line $y = x$ inside a tract of f . Further, the critical values of f grow like the exponential function; that is, the modulus of the critical values of f are $\exp(\pi k + \pi/4)/\sqrt{2}$ for $k \in \mathbb{Z}$. So, there exists λ and an annulus $A(r/\lambda, \lambda r)$ which contains no singular values of the restriction of f to D for arbitrarily large r . Hence the conditions of Theorem 4.1 are satisfied, so we deduce from Theorem 4.1 that $\dim J(f) \cap K(f) > 1$.

Next, we give an example with a direct tract with no logarithmic singularities. This example is the reciprocal of the entire function studied in [7]; for an illustration of the tracts of this function, see [7, Figure 1].

Example 5.2. Consider the entire function

$$f(z) = \exp(-g(z)), \quad \text{where } g(z) = \sum_{k=1}^{\infty} \left(\frac{z}{2^k}\right)^{2^k}.$$

Then $\dim J(f) \cap K(f) > 1$.

We will use several key results about f from [7, p. 254-258]. The first is the existence of an infinite tree on which f is very large. Introducing some notation, fix $0 < \varepsilon \leq 1/8$ and set $r_n = (1 + \varepsilon)2^{n+1}$ and $r'_n = (1 - 2\varepsilon)2^{n+2}$ for $n \in \mathbb{N}$. Then, for $j \in \{0, 1, \dots, 2^n - 1\}$, let

$$B_{j,n} = \left\{ r \exp\left(\frac{\pi i}{2^n} + \frac{2\pi i j}{2^n}\right) : r_n \leq r \leq r'_n \right\}$$

and

$$C_{j,n}^{\pm} = \left\{ r \exp \left(\frac{\pi i}{2^n} + \frac{2\pi i j}{2^n} \pm \frac{r - r'_n}{r_{n+1} - r'_n} \frac{\pi i}{2^{n+1}} \right) : r'_n \leq r \leq r_{n+1} \right\}.$$

We then obtain an infinite binary tree

$$T = [-ir_1, ir_1] \cup \bigcup_{n=1}^{\infty} \bigcup_{j=0}^{2^n-1} (B_{j,n} \cup C_{j,n}^{\pm}).$$

If n is large enough, then $\operatorname{Re} g(z) < -2^{2^n}$, for $z \in B_{j,n} \cup C_{j,n}^{\pm}$, $j = 0, 1, \dots, 2^n - 1$.

The second key result in [7] is that if $r_n \leq |z| \leq r'_n$, then

$$g(z) = (1 + \eta(z)) \left(\frac{z}{2^n} \right)^{2^n},$$

where $\eta(z) \rightarrow 0$ as $n \rightarrow \infty$.

Let $\rho_n = (1 + 2\varepsilon)2^{n+1}$ and $\rho'_n = (1 - 3\varepsilon)2^{n+2}$. The authors then show a third key result that

$$|zg'(z) - 2^n g(z)| < 1/2 |g(z)|,$$

for $\rho_n \leq |z| \leq \rho'_n$.

The final key result we need is that for $\rho_n \leq r \leq \rho'_n$, we have that $\arg g(re^{i\theta})$ is an increasing function of θ and increases by $2^n 2\pi$ as θ increases by 2π . The authors conclude that for $\rho_n \leq r \leq \rho'_n$ the circle $\{z : |z| = r\}$ contains exactly 2^n arcs where $\operatorname{Re} g(re^{i\theta}) < \log \rho$ for r and n sufficiently large that $\{z : |z| = r\} \cap D \neq \emptyset$ and $\log \rho > -2^{2^n}$.

Let D be a direct tract of f with boundary value $R > 0$. As f has no zeros, D is simply connected. Further, from [31, Example 3.3], f has a bounded set of asymptotic values, and we can assume they all lie in $D(0, R)$. Hence, it remains to check that the critical values of f are suitably well separated in order to satisfy condition (ii) in Theorem 4.1. We deduce this by using the key results above discussed in [7], as well as an application of Rouché's theorem and properties of level curves.

First, from the third key result above $g'(z) \neq 0$ in $\overline{A(\rho_n, \rho'_n)}$ for any n and so any critical points of f must lie in $\cup_{n \in \mathbb{N}} A(\rho'_{n-1}, \rho_n)$. Now, since

$$|zg'(z) - 2^n g(z)| < 1/2 |g(z)|,$$

for $\rho_n \leq |z| \leq \rho'_n$, by Rouché's theorem $g(z)$ and $zg'(z)$ have the same number of zeros in $D(0, \rho_n)$. Further, $g(z)$ has 2^n zeros in $D(0, \rho_n)$ by the second key result and another application of Rouché's theorem. Therefore, f has exactly $2^n - 1$ critical points in $D(0, \rho_n)$ and hence 2^{n-1} critical points in $A(\rho'_{n-1}, \rho_n)$, for $n \geq 1$. All such critical points of f , apart from 0, must lie in components of $A(\rho'_{n-1}, \rho_n) \cap D$, for some $n \geq 1$. From each such critical point there must originate at least 4 unbounded level curves which go to ∞ through $A(\rho_n, \rho'_n) \cap D$. By the fourth key result above, exactly 2 of these curves can pass through

each component of $A(\rho_n, \rho'_n) \cap D$, and the unions of these level curves must meet the tree T . Hence, the modulus of the level curves must be at least the minimum modulus of f on the tree T in $A(\rho'_{n-1}, \rho_n) \cap D$. Therefore, the modulus of each of the critical values of the critical points in $A(\rho'_{n-1}, \rho_n) \cap D$ is at least

$$(19) \quad \min\{|f(z)| : z \in T, |z| = \rho'_{n-1}\} \geq \exp(2^{2^{n-1}}),$$

for $n \geq 1$, by the first key result and since $r_{n-1} < \rho'_{n-1} < r'_{n-1}$.

We claim that for any $k > 1$ there exist arbitrarily large r such that the annulus $A(r, kr)$ contains no critical values of f . To prove this, first note that, for all $n \geq 1$, the function f has $2^n - 1$ critical points in $D(0, \rho_n)$. Hence, by (19) (with $n - 1$ replaced by n), there are at most $2^n - 1$ critical values of f in $A(\exp(2^{2^{n-1}}), \exp(2^{2^n}))$. Now, the number of annuli of the form $A(\exp(2^{2^{n-1}})k^m, \exp(2^{2^n})k^{m+1})$ that lie in $A(\exp(2^{2^{n-1}}), \exp(2^{2^n}))$ is at least $2^{2^{n-1}} / \log k$, because

$$\exp(2^{2^n}) \geq \exp(2^{2^{n-1}})k^{2^{2^{n-1}} / \log k}.$$

This proves the claim, since at most $2^n - 1$ of these annuli can contain a critical value of f . Therefore, we may apply Theorem 4.1 and obtain that $\dim J(f) \cap K(f) > 1$.

Remark. Bergweiler and Karpińska [8, Theorem 1.1] show that if f is a transcendental entire function that satisfies a certain regularity condition, then $\dim J(f) \cap I(f) = 2$. This result applies to our Example 5.1, but not to Example 5.2 since the regularity condition in [8] implies that f has finite order [8, p. 533], which is not the case for Example 5.2 by a theorem of Pólya [13, Theorem 2.9].

Acknowledgements. The author would like to thank his supervisors Gwyneth Stallard and Phil Rippon for their guidance and help in the preparation of this paper. The author also thanks Walter Bergweiler for a helpful conversation about direct tracts.

REFERENCES

- [1] I. N. Baker. The domains of normality of an entire function. *Ann. Acad. Sci. Fenn. Ser. A I Math.*, 1(2):277–283, 1975.
- [2] K. Barański. Hausdorff dimension of hairs and ends for entire maps of finite order. *Math. Proc. Cambridge Philos. Soc.*, 145(3):719–737, 2008.
- [3] K. Barański, B. Karpińska, and A. Zdunik. Hyperbolic dimension of Julia sets of meromorphic maps with logarithmic tracts. *Int. Math. Res. Not. IMRN*, (4):615–624, 2009.
- [4] W. Bergweiler. Iteration of meromorphic functions. *Bull. Amer. Math. Soc. (N.S.)*, 29(2):151–188, 1993.
- [5] W. Bergweiler. The size of Wiman-Valiron discs. *Complex Var. Elliptic Equ.*, 56(1-4):13–33, 2011.
- [6] W. Bergweiler. On the set where the iterates of an entire function are bounded. *Proc. Amer. Math. Soc.*, 140(3):847–853, 2012.
- [7] W. Bergweiler and A. Eremenko. Direct singularities and completely invariant domains of entire functions. *Illinois J. Math.*, 52(1):243–259, 2008.

- [8] W. Bergweiler and B. Karpińska. On the Hausdorff dimension of the Julia set of a regularly growing entire function. *Math. Proc. Cambridge Philos. Soc.*, 148(3):531–551, 2010.
- [9] W. Bergweiler, P. J. Rippon, and G. M. Stallard. Dynamics of meromorphic functions with direct or logarithmic singularities. *Proc. Lond. Math. Soc. (3)*, 97(2):368–400, 2008.
- [10] C. J. Bishop. A transcendental Julia set of dimension 1. *Invent. Math.*, 212(2):407–460, 2018.
- [11] R. Bowen. Hausdorff dimension of quasicircles. *Inst. Hautes Études Sci. Publ. Math.*, (50):11–25, 1979.
- [12] A. E. Eremenko and M. Y. Lyubich. Dynamical properties of some classes of entire functions. *Ann. Inst. Fourier (Grenoble)*, 42(4):989–1020, 1992.
- [13] W. K. Hayman. *Meromorphic functions*. Oxford Mathematical Monographs. Clarendon Press, Oxford, 1964.
- [14] W. K. Hayman. The local growth of power series: a survey of the Wiman-Valiron method. *Canad. Math. Bull.*, 17(3):317–358, 1974.
- [15] F. Iversen. *Recherches sur les fonctions inverses des fonctions méromorphes*. PhD thesis, Helsingfors, 1914.
- [16] J. Kotus and M. Urbański. Fractal measures and ergodic theory of transcendental meromorphic functions. In *Transcendental dynamics and complex analysis*, volume 348 of *London Math. Soc. Lecture Note Ser.*, pages 251–316. Cambridge Univ. Press, Cambridge, 2008.
- [17] A. J. Macintyre. Wiman’s method and the ‘flat regions’ of integral functions. *Quart. J. Math., Oxford Ser. 9*, pages 81–88, 1938.
- [18] C. McMullen. Area and Hausdorff dimension of Julia sets of entire functions. *Trans. Amer. Math. Soc.*, 300(1):329–342, 1987.
- [19] M. Misiurewicz. On iterates of e^z . *Ergodic Theory Dynam. Systems*, 1(1):103–106, 1981.
- [20] R. Nevanlinna. *Analytic functions*. Translated from the second German edition by Phillip Emig. Die Grundlehren der mathematischen Wissenschaften, Band 162. Springer-Verlag, New York-Berlin, 1970.
- [21] J. Osborne. Connectedness properties of the set where the iterates of an entire function are bounded. *Math. Proc. Cambridge Philos. Soc.*, 155(3):391–410, 2013.
- [22] T. Ransford. *Potential theory in the complex plane*, volume 28 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 1995.
- [23] P. J. Rippon and G. M. Stallard. Singularities of meromorphic functions with Baker domains. *Math. Proc. Cambridge Philos. Soc.*, 141(2):371–382, 2006.
- [24] G. Rottenfusser, J. Rückert, L. Rempe, and D. Schleicher. Dynamic rays of bounded-type entire functions. *Ann. of Math. (2)*, 173(1):77–125, 2011.
- [25] H. Schubert. *Über die Hausdorff-Dimension der Juliamenge von Funktionen endlicher Ordnung*. PhD thesis, 2007.
- [26] G. M. Stallard. The Hausdorff dimension of Julia sets of entire functions. II. *Math. Proc. Cambridge Philos. Soc.*, 119(3):513–536, 1996.
- [27] G. M. Stallard. The Hausdorff dimension of Julia sets of entire functions. IV. *J. London Math. Soc. (2)*, 61(2):471–488, 2000.
- [28] M. Tsuji. *Potential theory in modern function theory*. Chelsea Publishing Co., New York, 1975. Reprinting of the 1959 original.
- [29] T. F. Tyler. Maximum curves and isolated points of entire functions. *Proc. Amer. Math. Soc.*, 128(9):2561–2568, 2000.
- [30] G. Valiron. *Lectures on the general theory of integral functions*. Édouard Privat, Toulouse, 1923.

- [31] J. Waterman. Identifying logarithmic tracts. To appear in *Ann. Acad. Sci. Fenn.* Preprint arXiv:1902.04330.

SCHOOL OF MATHEMATICS AND STATISTICS, THE OPEN UNIVERSITY, WALTON HALL, MILTON KEYNES
MK7 6AA, UK

Email address: james.waterman@open.ac.uk