Open Research Online

The Open University's repository of research publications and other research outputs

The Piancaldoli meteorite: A forgotten primitive LL 3.1 ordinary chondrite

Journal Item

How to cite:

Marocchi, Yves; Bonal, Lydie; Gattacceca, Jerome; Piani, Laurette; Beck, Pierre; Greenwood, Richard; Basque, Anne; Nuccio, Mario and Foresta Martin, Franco (2020). The Piancaldoli meteorite: A forgotten primitive LL 3.1 ordinary chondrite. *Meteoritics and Planetary Science*, 55(8)

For guidance on citations see [FAQs](#).

© 2020 The Meteoritical Society

<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Version: Accepted Manuscript

Link(s) to article on publisher's website:
<http://dx.doi.org/doi:10.1111/maps.13552>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

---

oro.open.ac.uk

The Piancaldoli meteorite: A forgotten primitive LL3.10 ordinary chondrite

Yves Marrocchi^{1,*}, Lydie Bonal², Jérôme Gattacceca³, Laurette Piani¹, Pierre Beck²,
Richard Greenwood⁴, Jolantha Eschrig², Anne Basque¹, Pasquale Mario Nuccio⁵,
Franco Foresta Martin^{6,7}

¹CRPG, CNRS, Université de Lorraine, UMR 7358, Vandoeuvre-lès-Nancy, 54501, France

²Institut de Planétologie et d'Astrophysique de Grenoble, Grenoble, France

³Aix-Marseille Univ, CNRS, IRD, Coll France, INRAE, CEREGE, Aix-en-Provence, France

⁴PSS, Open University, Walton Hall, Milton Keynes MK7 6AA, UK

⁵ Full professor of Geochemistry and Volcanology, Università di Palermo, Italy

⁶Instituto Nazionale di Geofisica e Vulcanologia, Sezione di Palermo, 90146 Palermo, Italy

⁷ Laboratorio Museo di Scienze della Terra, Ustica, Palermo, Italy

*Corresponding author: yvesm@crpg.cnrs-nancy.fr

Abstract

The Piancaldoli ordinary chondrite fell in northern Italy on 10 August 1968. Preliminary studies led to its classification as a LL3.4 unequilibrated ordinary chondrite. However, recent developments in classification procedures have prompted us to re-examine its mineralogical, petrographic, spectroscopic, chemical, and isotopic features in a multi-technique study. Raman spectra and magnetic properties indicate that Piancaldoli experienced minimal thermal metamorphism, consistent with its high bulk hydrogen content and the Cr contents of ferroan olivines in its type-II chondrules. In combination with findings of previous studies, our data thus confirm the variability of Cr contents in ferroan olivines in type-II chondrules as a proxy of thermal metamorphism. Furthermore, our results reveal that Piancaldoli is less altered than previously reported and should be reclassified as an LL3.10 unequilibrated ordinary chondrite. Our results also imply that the bulk deuterium enrichment, as observed in Piancaldoli (LL3.10), Bishunpur (LL3.15), and Semarkona (LL3.00), is a specific signature of the most primitive unequilibrated ordinary chondrites. Based on our results, we propose that, to date, Piancaldoli is the second least-altered unequilibrated ordinary chondrite fall after Semarkona. This work reiterates the importance of meteorite collections worldwide as fundamental resources for studying the formation conditions and evolution our solar system.

Keywords: Ordinary chondrites, meteorite classification, thermal metamorphism, Piancaldoli

1. Introduction

51

52   On 26 April 1803, thousands of stones fell upon the small city of L'Aigle, France (~150
53 km west of Paris). At that time, the extraterrestrial nature of meteorites was not yet recognized
54 and they were still considered to be ‘thunderstones’ resulting from atmospheric processes. The
55 precise description of the L'Aigle fall by dozens of people and the unique nature of the
56 recovered rocks led physicist Jean-Baptiste Biot to conclude their origin to be unambiguously
57 extraterrestrial. The extraterrestrial origin of meteorites had already been proposed on the basis
58 of meteor ballistic trajectories and chemical analyses performed by Ernst Florens Friedrich
59 Chladni (see Marvin, 2010 for a review) and Edward Howard (Howard et al., 1802). Since these
60 works, meteorites have been recognized as objects of scientific importance, and natural history
61 museums worldwide have established invaluable meteorite collections. Despite the recent
62 boom in space exploration, most of our current understanding of the conditions and chronology
63 of the formation of the solar system has been established thanks to the meticulous work of
64 meteorite collection and curation (Heck et al. 2019).

65   Over the past two decades, the number of meteorites reported to the Meteoritical Society
66 has increased exponentially thanks to field trips organized to cold and hot deserts expressly to
67 search for meteorites (e.g., Harvey, 2003; Gattacceca et al. 2011; Evatt et al. 2020). These
68 harvests have enabled the characterization of some of the most primitive meteorites known so
69 far, which are now available to the scientific community. However, meteorite finds are exposed
70 to terrestrial weathering and contamination, which largely affect their primitive characteristics
71 (Alexander 2017; Stephant et al. 2018; Vacher et al. 2020). For instance, Antarctic and Saharan
72 samples are commonly affected by Na loss and/or metal alteration, and Antarctic samples by
73 oxygen isotopic exchange with Antarctic water (Alexander et al. 2018). Thus, as meteorite falls
74 are collected before any significant terrestrial alteration can occur, they are of primary
75 importance for deciphering the formation of the solar system.

In addition to terrestrial weathering, most chondrites experienced secondary alteration on their asteroidal parent bodies, resulting in changes to their primary texture, mineralogy, and chemical and isotopic compositions (Brearley 2006; Marrocchi et al. 2018). These petrographic and chemical modifications are the basis of petrologic type assignments (Van Schmus and Wood 1967) on a scale from 1 to 6, reflecting the progressive roles of low-temperature aqueous alteration (types 3 to 1) and thermal metamorphism (types 3 to 6). In theory, type-3 chondrites thus correspond to primitive materials that underwent minimal modifications after their agglomeration in the protoplanetary disk 4.56 Ga (Amelin et al., 2010). However, new classification procedures based on thermoluminescence (TL; Sears et al. 1980), chemical characteristics (e.g., Cr content in ferroan olivines in type-II chondrules; Grossman and Brearley 2005) and Raman spectroscopy (Bonal et al. 2006, 2007, 2016) permit the quantification of discrete variations among type-3 chondrites. Accordingly, type-3 chondrites are now subdivided into categories ranging from 3.00 to 3.9, with 3.00 being the most *primitive* extraterrestrial materials, virtually unchanged since their agglomeration in the disk.

Because these advances in the classification of minimally altered chondrites are relatively recent (Grossman and Brearley 2005; Bonal et al. 2006, 2016), many previously analyzed chondrites have not been characterized according to the new classification criteria nor studied in further detail after their initial classification. This is fundamental, however, as the identification of primitive chondrites with minimal parent-body alteration improves our understanding of the formation and evolution of solids in the protoplanetary disk. In particular, Al-Mg chronology requires the characterization of refractory inclusions and chondrules in chondrites of petrologic subtypes ≤ 3.1 due to the fast self-diffusion of Mg in minerals and glassy mesostases (Kita and Ushikubo 2012; Van Orman et al. 2014; Marrocchi et al. 2019).

While vacationing in Sicily in 2018, the first author (YM) fortuitously met Franco Foresta Martin, a geologist and a journalist at the Italian national daily newspaper *Corriere*

della Sera. He brought to my knowledge the history of the Piancaldoli meteorite that fell near Florence in northern Italy on 10 August 1968. A fireball was observed over Yugoslavia and central and northern Italy, followed by explosions accompanying the break-up of the meteorite. Three meteorite fragments totaling 13.1 g were found on the roof of a house in Piancaldoli. The Piancaldoli meteorite was first classified as a L3 chondrite (Carapezza and Nuccio 1971) before being reclassified as LL3 (Carapezza et al. 1975, 1976). Although no Raman or TL data were available for Piancaldoli, Rubin et al. (1982) proposed its reclassification as petrologic type 3.4 $\pm$ 0.2 based on silicate compositions. Consequently, Piancaldoli is listed as a LL3.4 ordinary chondrite in the Meteoritical Bulletin Database. Here we report a comprehensive description of Piancaldoli and show that its secondary thermal history was overestimated. Our results indicate that Piancaldoli is the second least-altered unequilibrated ordinary chondrite fall known to date.

2. Material and methods

2.1 Mineralogical and petrographic observations

We surveyed two sections of Piancaldoli: thin section USNM 5649 from the National Museum of Natural History, Smithsonian Institution (Washington D.C., USA) and a thick section prepared at the Centre de Recherches Pétrographiques et Géochimiques (CRPG-CNRS, Nancy, France) from a parent sample provided by the Dipartimento scienze della Terra e del Mare (Palermo, Italy). We also characterized a thin section of Semarkona (LL3.00; Grossman and Brearley, 2005; Muséum national d'Histoire naturelle, Paris, France) and a thick section of Bishunpur (LL3.15; Grossman and Brearley, 2005; Muséum national d'Histoire naturelle, Paris, France) for comparison. The sections were imaged by scanning electron microscopy on a JEOL JSM-6510 equipped with a Genesis energy dispersive x-ray (EDX) detector at the CRPG using a 3 nA electron beam accelerated at 15 kV. The chemical compositions of olivine grains were quantified using a Cameca SXFive electron microprobe at the Université Pierre et Marie Curie

(UPMC, Camparis, Paris, France) using a 150 nA focused beam accelerated at 15 kV. We analyzed Na, Mg, Si, Al, K, Ca, Fe, Ti, Cr, and Mn in olivine grains. The high beam current allowed detection limits for silicates to be 100 ppm for Al, Ca, and Ti, 150 ppm for Mn and Si, and 200 ppm for Na, K, Cr, Fe, and Mg. The PAP software was used for matrix corrections.

2.2 Raman spectroscopy

Raman spectroscopy was performed at the Ecole Normale Supérieure de Lyon (Laboratoire de Géologie de Lyon – Terre, Planètes, Environnement, France) using a LabRam Raman spectrometer (Horiba Jobin-Yvon) equipped with a 600 gr/mm grating and a Spectra Physics Ar⁺ laser ($\lambda = 514$ nm). Because carbonaceous matter is sensitive to laser-induced heating and the Raman bands of polyaromatic carbonaceous matter are dispersive, we followed the analytical procedures of Bonal et al. (2016, summarized here) to avoid any laser alteration of the carbonaceous matter and to facilitate meaningful comparisons with reference meteorites from the literature. The laser was focused through a 100× objective to obtain a ~ 2 μ m spot size. The power on the sample was 500 μ W. Each acquisition comprised two 30-s integrations that were averaged to make the final spectrum. Spectra were acquired under atmospheric conditions over the wavenumbers 500–2,200 cm^{-1} , covering the first-order carbon bands. Raman spectra of carbonaceous matter in the matrix of Piancaldoli were obtained both on isolated matrix fragments (50 spectra) and *in situ* in the thick section prepared at the CRPG (32 spectra). Around 30 matrix fragments (typical apparent diameter around 30 μ m) were manually selected from a gently crushed raw piece of Piancaldoli (initial sample of 50 mg) according to their color and texture under a binocular microscope. The selected matrix fragments were pressed between two glass slides that served as the substrate for the Raman analyses. The G and D bands were fitted with Breit-Wigner-Fano and Lorentzian profiles, respectively, to retrieve spectral parameters. We compared the width (full width at half maximum) of the D band (FWHM_D , cm^{-1}

¹) and its intensity relative to that of the G band (I_D/I_G) to those of reference samples as these spectral parameters are the most sensitive to the maturity range of type-3 chondrites (e.g., Bonal et al., 2016).

2.3 Infrared spectroscopy

Infrared (IR) spectra were obtained with a Bruker HYPERION 3000 infrared microscope at the Institut de Planétologie et d'Astrophysique de Grenoble (IPAG, Grenoble, France). The IR beam was focused through a 15× objective to a typical spot size on the sample of $40 \times 40 \mu\text{m}^2$. Spectra were acquired at 4 cm^{-1} spectral resolution with a MCT detector cooled with liquid nitrogen. Particular care was devoted to sample preparation, which is critical in IR microspectroscopy. Samples must be sufficiently thin ($<100 \mu\text{m}$) and their surfaces flat and parallel to avoid absorption-band saturation and scattering artifacts, respectively (Raynal et al. 2000). Small matrix fragments (30–50 μm) were selected under a binocular microscope and crushed between two diamond windows for analyses in the $4000\text{--}650 \text{ cm}^{-1}$ spectral range. The diamond windows were loaded into an environmental cell, designed and built at IPAG, capable of achieving temperatures up to 300°C under primary or secondary dynamic vacuum (10^{-4} to 10^{-7} mbar). Optical access is via both sides of the cell through KBr windows, enabling measurements in transmission mode. Samples were progressively heated and analyzed at 20, 100, and 300°C for 1 hour at each temperature.

The transmission spectra were automatically converted to absorbance ($A = -\log (T/T_0)$, where T_0 and T are the background and sample transmittance, respectively). To remove interferences and scattering effects, a spline baseline was calculated and subtracted from the raw data. The resulting spectra were normalized to the intensity of the silicate Si-O stretching band (1000 cm^{-1}) at $10 \mu\text{m}$ thickness, which does not evolve over the temperature range of the measurements. Reflectance spectra were obtained on powdered fragments of Piancaldoli using

the SHADOWS instrument (Potin et al., 2018). Spectra were normalized to spectralon^{TM} and
infragold^{TM} and measured under standard mode.

2.4 Magnetic properties

Hysteresis measurements were performed on a 122-mg bulk (matrix + chondrules)
fragment of Piancaldoli with a Princeton Micromag vibrating sample magnetometer with a
maximum applied field of 1 T and a sensitivity of $\sim 5 \times 10^{-9}$ A m². We analyzed hysteresis loops
to obtain the ratio of saturation remanent magnetization (M_{RS}) to saturation magnetization (M_S)
and the coercive force (B_C). M_S was determined by correcting the hysteresis loop for the high-
field susceptibility determined by a linear fit of the hysteresis loops for applied fields >0.9 T.
Remanent coercive force (B_{CR}) was determined by back-field experiments performed with the
magnetometer. The coercivity spectrum was estimated from the derivative of the isothermal
remanent magnetization by stepwise measurements using the magnetometer. The evolution of
low field magnetic susceptibility at low temperature was studied using an Agico MFK1
apparatus equipped with a CSL cryostat. All magnetic measurements were performed at the
Centre Européen de Recherche et d'Enseignement des Géosciences de l'Environnement (Aix-
en-Provence, France).

2.5 Hydrogen isotopes

Hydrogen concentrations and isotopic compositions were measured using the Thermo
Scientific EA IsoLink - DeltaV IRMS System at the CRPG according to the procedure detailed
in Lupker et al. (2012), which we summarize here. A small piece of the meteorite was crushed
into powder and two aliquots (5.97 and 7.82 mg, respectively) were loaded into tin capsules
and degassed at 120 °C under vacuum for 48 h to minimize the contribution of adsorbed

atmospheric water (Lupker et al. 2012; Vacher et al. 2016, 2020). Hydrogen isotopic compositions are expressed relative to that of Standard Mean Ocean Water (SMOW, $D/H_{SMOW} = 155.76 \times 10^{-6}$) as $\delta D [‰] = [(D/H_{sample} / D/H_{SMOW}) - 1] \times 1,000$. Reproducibilities estimated from reference materials are better than 10% (2σ) for H concentration and 0.5×10^{-6} for D/H (or 5‰ for δD).

2.6 Oxygen isotopes

High-precision oxygen isotopic measurements were performed at the Open University (Milton Keynes, UK) using an infrared laser-assisted fluorination system (Miller et al., 1999; Greenwood et al., 2017). Two individual analyses of whole rock chips of Piancaldoli were undertaken, with each replicate having a mass of about 2 mg. After fluorination, the O_2 released was purified by passing through two cryogenic nitrogen traps and over a bed of heated KBr. O_2 analyses were performed using a MAT 253 dual inlet mass spectrometer. Analytical precision (2σ), based on replicate analyses of an internal obsidian standard, is $\pm 0.053 ‰$ for $\delta^{17}O$, $\pm 0.095 ‰$ for $\delta^{18}O$ and $\pm 0.018 ‰$ for $\Delta^{17}O$ (Starkey et al., 2016). Oxygen isotopic analyses are reported in standard δ notation, where $\delta^{18}O$ has been calculated as: $\delta^{18}O = [(^{18}O/^{16}O_{sample}/^{18}O/^{16}O_{ref.}) - 1] \times 100$ and similarly for $\delta^{17}O$ using the $^{17}O/^{16}O$ ratio. $\Delta^{17}O$, which represents the deviation from the terrestrial fractionation line, has been calculated as: $\Delta^{17}O = \delta^{17}O - 0.525 \times \delta^{18}O$ in order to compare our results with those obtained by Clayton et al (1991).

3. Results

3.1 Petrographic overview

The Piancaldoli sections contain abundant chondrules (Figs. 1, 2) and chondrule fragments (Fig. 3) in an optically dark matrix. Type-I chondrules (characterized by abundant Fe-Ni metal beads, FeO-poor silicates and volatile element depletion) are predominantly

olivine-rich and relatively homogeneous (Fig. 2A), whereas type-II chondrule (characterized by FeO-rich silicates and more chondritic abundances of volatile elements) are compositionally variable with normally zoned and relict, Mg-rich, olivine grains (Villeneuve et al., 2020; Fig. 2B). Some ferroan olivine grains show igneous zoning profiles, with rims enriched in Cr (Fig. 4), and chromite exsolutions are commonly observed in type-II chondrule olivines (Fig. 5). Although some type-II chondrules show devitrified mesostases, both types are characterized by well-preserved glassy mesostases (Figs. 2–4) with no specific evidence of aqueous alteration. Dusty olivine grains were observed in about 5% of 350 observed chondrules. Opaque mineral assemblages are rounded and occur predominantly along chondrule exteriors, but also rarely within the interiors of some chondrules (Fig. 1). Ferroan olivines in type-II chondrules of Piancaldoli contain 0.2–0.5 wt.% Cr₂O₃, averaging 0.36 ± 0.20 wt.% Cr₂O₃ (1 σ , Fig. 6, Table S1); those of Semarkona and Bishunpur contain 0.45 ± 0.11 and 0.23 ± 0.13 wt.% Cr₂O₃, respectively (Fig. 6).

3.2 Raman and Infrared spectral characteristics

Every Raman spectrum obtained of the Piancaldoli matrix exhibits the D and G bands, indicating the presence of polyaromatic carbonaceous matter. We obtained mean values of FWHM_D (cm⁻¹) = 173.4 ± 20.2 and $I_D/I_G = 0.912 \pm 0.089$ for *in-situ* thin-section analyses ($n = 32$) and FWHM_D (cm⁻¹) = 159.2 ± 37.8 and $I_D/I_G = 0.996 \pm 0.145$ for matrix fragments ($n = 50$; the errors reflecting the variable structural order of the polyaromatic carbonaceous matter). These spectral parameters are thus variable throughout Piancaldoli: the structural order of the polyaromatic carbonaceous matter in the CRPG section of Piancaldoli reflects a metamorphic grade similar to that of Bishunpur (LL3.15), whereas the spectral parameters obtained for the matrix fragments are more dispersed and tend to reflect a slightly higher metamorphic grade

(Fig. 7). This discrepancy might be explained by the brecciated nature of Piancaldoli (discussed in section 4).

The presence of aliphatic bands in the 2800–3000 cm^{-1} range of IR spectra (Fig. 8) confirms our manual selection of matrix fragments because the matrix is the only petrographic component containing organics in chondrites. These IR spectra mostly exhibit bands related to Si-O stretching around 1000 cm^{-1} , indicating olivine to be the dominant silicate. The stretching mode of molecular water is poorly visible at $\sim 3400 \text{ cm}^{-1}$, and the OH^- band at $\sim 3670 \text{ cm}^{-1}$ is absent (Fig. 8).

The reflectance spectrum obtained for Piancaldoli presents the two typical absorption bands at 1 and 2- μm diagnostic of Fe-bearing silicates (olivine + pyroxene). The shape bands, in particular the reflectance maxima around 1.6 μm , is typical of LL chondrites (Fig 9). However, the depth of the two bands is lower than typical ordinary chondrites, which is likely related to the unequilibrated nature of the sample. The visible reflectance at 550 nm is lower than type 4-6 ordinary chondrites, and in low-side of the range of value measured for type 3 ordinary chondrites.

3.3 Magnetic properties

The hysteresis properties of the bulk Piancaldoli fragment are $M_S = 6.79 \text{ A m}^2/\text{kg}$, $M_{RS} = 0.216 \text{ A m}^2/\text{kg}$, $B_C = 9.02 \text{ mT}$, and $B_{CR} = 62.8 \text{ mT}$, in agreement with observed values for LL3 chondrite falls (Gattacceca et al. 2014). The M_S value indicates a metal content similar to other LL3 chondrites. The coercivity spectrum of Piancaldoli (Fig. 10) is difficult to fit with a typical combination of log-normally distributed components, and is notably shifted toward low coercivities compared to other LL chondrite falls, including LL3 chondrites. In particular, the

absence of a peak in the high coercivity range (>500 mT) precludes the presence of ordered tetrataenite in cloudy zone microstructures of zoned taenite (Gattacceca et al. 2014). Such microstructures form only by the cooling of Fe-Ni metal from above 320°C at cooling rates slower than $\sim 1000^{\circ}\text{C/Myr}$ (Gattacceca et al. 2014; Maurel et al. 2019) and can be disordered by impact-induced thermal events at or above shock stage S3/S4 (Gattacceca et al. 2014). In the case of Piancaldoli (shock stage S1), the absence of ordered tetrataenite constrains the peak metamorphism temperature to $<320^{\circ}\text{C}$. LL chondrites with broadly similar coercivity spectra include Bishunpur, Krymka, and Chainpur, although Piancaldoli has the lowest coercivities. Conversely, Vicencia (LL3) and all other equilibrated LL chondrites have high-coercivity components attributable to ordered tetrataenite in the cloudy zone structure (Fig. 10). It is therefore likely that the classification of Vicencia as LL3.2 (Keil et al. 2015) should be revised to a higher petrologic subtype, consistent with the observed Raman spectral parameters (see their Fig. 8, and Fig. 7 herein). Indeed, Vicencia appears to be more metamorphosed than Chainpur (LL3.4) and less than Tieschitz (H/L3.6).

The low temperature magnetic measurements performed on Piancaldoli do not reveal a Verwey transition at 120K that would indicate the presence of magnetite. Magnetite has been detected through low temperature magnetic measurements in a few primitive LL3 falls, such as Semarkona and Krymka (Keil et al., 2015), and through petrographic observation in, e.g., Semarkona (Hutchison et al., 1987). The absence of the Verwey transition in Piancaladoli confirms the absence of significant aqueous alteration.

3.4 H and O isotopes

The two analyzed aliquots of Piancaldoli contain 0.124 and 0.127 wt.% H, equivalent to 1.12 and 1.14 wt.% H_2O , respectively, and have respective D/H values of 193.2×10^{-6} ($\delta\text{D} = -57.7\text{‰}$) and 187.8×10^{-6} ($\delta\text{D} = -57.6\text{‰}$). These values are broadly typical of mildly

metamorphosed LL3 chondrites (Robert et al. 1979; McNaughton et al. 1982; Yang and Epstein 1983; Alexander et al. 2012; Vacher et al. 2020). Although the hydrogen concentrations are on the higher end of those measured for LL3 chondrites similar to Semarkona ($\sim 0.04\text{--}0.12$ wt.% H), the D/H ratios are intermediate among the range observed for LL3 chondrites (158×10^{-6} to 505×10^{-6} ; Fig. 11).

The results of oxygen isotope measurement of Piancaldoli are plotted in Fig. 12. The two individual analyses show consistent results with $\delta^{18}\text{O} = 5.16 \pm 0.21$ ‰ (2σ) and $\delta^{17}\text{O} = 3.63 \pm 0.18$ ‰ (2σ). Compared to other ordinary chondrites, Piancaldoli plots at the edge of the LL chondrite field with a relatively low $\Delta^{17}\text{O}$ value of 0.95 ± 0.08 ‰ (2σ ; Fig. 12).

4. Discussion

The Piancaldoli chondrite contains sharply defined type-I and type-II chondrules (Fig. 2) with a mean apparent diameter of 901 ± 445 μm , and a median apparent diameter of 804 μm ($n=352$). Rubin et al. (1982) described a Piancaldoli clast with ~ 100 radial-pyroxene micro-chondrules 0.2–64 μm in apparent diameter. Chondrule olivine and pyroxene crystals show heterogeneous compositions with (i) olivine varying between 66 and 98 mol% of forsterite and (ii) pyroxene ranging from 2 to 24 mol% of ferrosilite (Carapezza et al., 1976; Rubin et al., 1982). The chemical composition of Piancaldoli is typical of LL ordinary chondrites with $\text{FeO}/\text{SiO}_2 = 0.48$, $\text{SiO}_2/\text{MgO} = 1.62$, and $\text{Fe}^0/\text{Fe}_{\text{tot}} = 0.12$ (Carapezza et al. 1976; Wasson and Kallemeyn, 1988). In addition, the chemical compositions of the fine-grained opaque matrix and the whole rock were used to propose that Piancaldoli is an unequilibrated LL ordinary chondrite of petrologic type 3.4 ± 0.2 (Carapezza et al. 1976; Rubin et al. 1982). However, all data collected herein indicate that Piancaldoli is more primitive than this estimation. Thus, in this section, we (i) assess the extent of thermal alteration experienced by Piancaldoli and (ii) propose a new classification.

The distribution of Cr contents in ferroan olivines in type-II chondrules is considered a proxy of incipient thermal metamorphism. We obtained a mean value of 0.36 ± 0.19 wt.% Cr₂O₃ (Fig. 6; Table S1) for type-II chondrule olivines in Piancaldoli, and our measurements of those in Semarkona and Bishunpur during the same analytical session are in good agreement with previous reports (Grossman and Brearley 2005; Fig. 6). This suggests that Piancaldoli experienced minimal thermal metamorphism corresponding to petrologic type 3.10 (Fig. 6; Grossman and Brearley 2005), supported by the occurrence of igneous Cr zoning patterns (i.e., rims enriched in Cr) in some type-II chondrule olivines (Fig. 4). Such primary textures are extremely sensitive to thermal metamorphism as they are replaced by complex, subparallel, needle-like exsolutions of Cr-rich phases (likely chromite; Fig. 5) upon heating (Grossman and Brearley 2005). As the separation of Cr-rich phases from FeO-rich olivine grains is nearly complete by petrologic type 3.2 (Grossman and Brearley 2005), the observed Cr zonings in Piancaldoli olivine are a strong argument in favor of Piancaldoli's primitive nature.

Tetrataenite dominates the magnetic properties of all but the most primitive LL chondrites (and those that experienced shock stage S4 or greater). The absence of tetrataenite revealed by the magnetic properties of Piancaldoli sets it among the most primitive LL chondrites and constrains the peak metamorphic temperature to <320 °C. This low degree of thermal metamorphism is supported by the structural order of the polyaromatic carbonaceous matter in Piancaldoli. Indeed, the Raman spectral parameters of polyaromatic carbon bands of Piancaldoli are comparable with those of Bishunpur (LL3.15; Fig. 7). Interestingly, spectra obtained on matrix fragments selected from a piece of bulk Piancaldoli exhibit a wider spectral variability, and, on average, a higher maturity (higher I_D/I_G and lower FWHM_D, Fig. 7). Therefore, some parts of Piancaldoli appear to have experienced a higher peak metamorphic temperature consistent with a petrologic type closer to 3.4. This variable metamorphic grade is consistent with previous petrographic observations and could be related to the brecciated nature

of Piancaldoli (as attested by the presence of light- and dark-colored matrix domains in section USNM 5640; Rubin et al. 1982). Indeed, the fine exsolutions of Cr-rich phases commonly observed in Piancaldoli (Fig. 5) suggest that it is more altered than Semarkona, in which such textures are absent (Grossman and Brearley 2005).

Infrared spectra of Piancaldoli's matrix indicate, if not the absence, the really low abundance of secondary hydrated minerals, and thus that Piancaldoli did not experience significant aqueous alteration. This is in contrast to the matrix of Semarkona (LL3.00), which clearly contains hydrated material, most likely smectite (Alexander et al. 1989; Quirico et al. 2003).

The hydrogen content and isotopic composition of Piancaldoli show important similarities with the two least-altered ordinary chondrite falls, Semarkona and Bishunpur (Fig. 11). The hydrogen content of Piancaldoli (~ 0.125 wt.%) is among the highest observed in ordinary chondrites (despite the removal of adsorbed atmospheric water before measurement in our methodology; Vacher et al. 2016, 2020) and is similar to the highest values reported for Semarkona (0.12 wt.% H; Alexander et al. 2012), further indicating that Piancaldoli experienced minimal thermal metamorphism, which would have resulted in hydrogen loss (Fig. 11). As neither phyllosilicates nor magnetite were observed in Piancaldoli, the H budget is likely controlled by: (i) insoluble organic matter and (ii) chondrules (silicates and glassy mesostases). However, a detailed TEM study should be performed to clearly rule out the presence of phyllosilicates (Hutchison et al., 1987). Although enriched in deuterium compared to more metamorphosed ordinary chondrites, the rather low D/H signature of Piancaldoli relative to Semarkona and Bishunpur may be a consequence of the D/H heterogeneities observed among unequilibrated LL chondrites. Indeed, D/H values reported for Semarkona and Bishunpur are highly heterogeneous, ranging from 393 to 609×10^{-6} and from 253 to 504×10^{-6} , respectively (McNaughton et al. 1982; Yang and Epstein 1983; Alexander et al. 2012; Piani et al. 2015).

Therefore, it seems that heterogeneous H isotopic compositions can be taken as an argument for the primitive nature of unequilibrated LL ordinary chondrites.

5. Concluding remarks

We studied the Piancaldoli chondrite via scanning electron microscopy, electron microprobe analyses, hydrogen and oxygen isotopic analyses, magnetic measurements, and Raman and IR spectroscopy. Characteristics of the Raman spectra and magnetic properties of Piancaldoli are consistent with its primitive nature as inferred from the variability of Cr contents in ferroan olivines in type-II chondrules. Furthermore, the high bulk H content of Piancaldoli indicates that it experienced only minimal thermal metamorphism. Our petrographic and chemical results therefore suggest that Piancaldoli should be reclassified as a LL3.10 unequilibrated ordinary chondrite, rather than a LL3.4 as previously proposed. We further suggest that the deuterium observed in Piancaldoli, Bishunpur, and Semarkona are specific signatures of the most primitive unequilibrated LL ordinary chondrites. Our results thus demonstrate that (i) Piancaldoli is the second least-altered unequilibrated ordinary chondrite fall after Semarkona, and (ii) primitive meteorites are most likely sitting, unstudied, in meteorite collections worldwide.

401
402
403
404
405
406
407
408
409
410
411
412
413
414
415

Acknowledgments

We thank Colette Guilbaud, Michel Fialin, Nicolas Rividi, Maxime Piralla, Lionel G. Vacher, and Thomas Rigaudier for their assistance during sample preparation and analysis. We thank Ashley King and Tim Gregory for helpful reviews and Timothy Jull for editorial handling. This is CRPG-CNRS contribution #2732.

421
422
423
424
425

426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450

Figure captions

Fig. 1: Composite back-scattered electron images of Piancaldoli: (A) USNM 5649 and (B) Piancaldoli-CRPG.

Fig. 2: Back-scattered electron images of representative (A) type-I and (B) type-II chondrules with well-preserved mesostases.

Fig. 3: Back-scattered electron image of a fish-shaped chondrule fragment.

Fig. 4: EDX maps of (A) Si + Mg and (B) Cr in euhedral ferroan olivine grains in a type-II chondrule of Piancaldoli. Olivine rims are enriched in Cr, representing an igneous zoning profile.

Fig. 5: Back-scattered electron image of a ferroan olivine grain in a type-II chondrule showing complex, subparallel, needle-like exsolutions of a Cr-rich phase (likely chromite).

Fig. 6: Standard deviation on Cr₂O₃ content *versus* mean Cr₂O₃ content of ferroan olivines in type-II chondrules in Piancaldoli compared to Semarkona and Bishunpur (data from this study and Grossman and Brearley, 2005). Piancaldoli lies between Semarkona and Bishunpur, suggesting it is of petrologic subtype 3.10.

Fig. 7: Spectral parameters (FWHMD vs. ID/IG, averages and one standard deviations) of Raman bands attributed to carbonaceous materials in Piancaldoli compared to other meteorites (see Bonal et al., 2016). Data for Vicencia (LL), Axtell (CV3) and ALHA 77307 (CO3) are also reported (see section 3.3).

Fig. 8: Baseline-corrected transmission IR spectra (normalized to the height of the Si-O band at 1000 cm⁻¹) of matrix fragments of Piancaldoli, acquired at 20 °C under dynamic vacuum. A vertical offset was applied to ease comparison. Notably, the water band is very weak and olivine is the dominant silicate.

Fig. 9: Left: Reflectance spectra of Piancaldoli powder compared to average ordinary chondrites spectra from the RELAB database. Spectra were normalized at 550 nm. Right:

Reflectance value at 550 nm of Piancaldoli compared to ordinary chondrites, as a function of petrographic type.

Fig. 10: Coercivity spectra, i.e., the derivative of the isothermal remanent magnetization, IRM, (normalized to saturation remanence) as a function of the acquisition field, of LL chondrite falls. The petrologic types and masses of each chondrite is indicated.

Fig. 11: Hydrogen content (reported as $1/H_{wt\%}$) and isotopic composition of Piancaldoli compared to H, L, and LL ordinary chondrites. The hydrogen concentration of Piancaldoli plots in the range of poorly metamorphosed LL3 chondrites. Error bars represent 1σ standard deviations for chondrites having multiply reported hydrogen compositions (data from Robert et al. 1979; McNaughton et al. 1982; Yang and Epstein 1983; Alexander et al. 2012; Vacher et al. 2020).

Fig. 12: The oxygen isotopic composition of Piancaldoli compared to H, L, and LL ordinary chondrites (data from Clayton et al., 1991 and Greenwood et al., 2020) . Piancaldoli plots at the edge of the LL chondrite field with a relatively low $\Delta^{17}O$ value.

500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524

References

- Alexander C. M. O'D., Barber D. J., and Hutchison R. 1989. The microstructure of Semarkona and Bishunpur. *Geochimica et Cosmochimica Acta* 53:3045–3057.
- Alexander C. M. O'D., Bowden R., Fogel M. L., Howard K. T., Herd C. D. K., and Nittler L. R. 2012. The Provenances of Asteroids, and Their Contributions to the Volatile Inventories of the Terrestrial Planets. *Science* 337:721–723.
- Alexander C. M. O'D. 2017. The origin of inner Solar System water. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 375:20150384–20.
- Alexander C. M. O'D., McKeegan K. D., and Altwegg K. 2018. Water Reservoirs in Small Planetary Bodies: Meteorites, Asteroids, and Comets. *Space Science Review* 214:36.

525 Amelin Y., Kaltenbach A., Iizuka T., Stirling C. H., Ireland T. R., Petaev M., Jacobsen S. B.
 526 2010. U–Pb chronology of the Solar System’s oldest solids with variable $^{238}\text{U}/^{235}\text{U}$. *Earth*
 527 *and Planetary Science Letters* 300:343–350.

528 Bonal L., Quirico E., Bourot-Denise M., and Montagnac G. 2006. Determination of the
 529 petrologic type of CV3 chondrites by Raman spectroscopy of included organic matter.
 530 *Geochimica et Cosmochimica Acta* 70:1849–1863.

531 Bonal L., Bourot-Denise M., Quirico E., Montagnac G., and Lewin E. 2007. Organic matter
 532 and metamorphic history of CO chondrites. *Geochimica et Cosmochimica Acta* 71:1605–
 533 1623.

534 Bonal L., Quirico E., Flandinet L., and Montagnac G. 2016. Thermal history of type 3
 535 chondrites from the Antarctic meteorite collection determined by Raman spectroscopy of
 536 their polyaromatic carbonaceous matter. *Geochimica et Cosmochimica Acta* 189:312–337.

537 Brearley A. J. 2006. The action of water. In *Meteorites and the early solar system II*, edited by
 538 Lauretta D. S. and McSween H. Y. Tucson, Arizona: The University of Arizona Press. pp.
 539 587–624.

540 Carapezza M., and Nuccio M. 1971. The Piancaldoli meteorite. *Meteoritics* 6, 255.

541 Carapezza M., Nuccio P.M., and Valenza M. 1975. Piancaldoli meteorite: chemistry and
 542 mineralogy. *Meteoritics* 10, 369.

543 Carapezza M., Nuccio P.M., and Valenza M. 1976. Piancaldoli meteorite: chemistry,
 544 mineralogy and petrology. *Meteoritics* 11, 165.

545 Clayton R. N., Mayeda T. K., and Goswami J. N. 1991. Oxygen isotope studies of ordinary
 546 chondrites. *Geochimica et Cosmochimica Acta* 55:2317–2337.

547 Evatt G. W., Smedley A. R. D., Joy K. H., Hunter L., Tey W. H., Abrahams I. D., and Gerrish
 548 L. 2020. The spatial flux of Earth’s meteorite falls found via Antarctic data. *Geology* 48:
 549 G46733.1.

550 Gattacceca J. et al. 2011. The densest meteorite collection area in hot deserts: The San Juan
551 meteorite field (Atacama Desert, Chile). *Meteoritics & Planetary Science* 46:1276–1287.

552 Gattacceca J., Suavet C., Rochette P., Weiss B.P., Winklhofer M., Uehara M., Friedrich J. 2014.
553 Metal phases in ordinary chondrites: magnetic hysteresis properties and implications for
554 thermal history. *Meteoritics and Planetary Science*, 49:652-676.

555 Greenwood R.C., Burbine T.H., and Franchi I.A. 2020. Linking asteroids and meteorites to the
556 primordial planetesimals population. *Geochimica et Cosmochimica Acta* 277:377-406.

557 Grossman J. N., and Brearley A. J. 2005. The onset of metamorphism in ordinary and
558 carbonaceous chondrites. *Meteoritics & Planetary Science* 40:87–122.

559 Harvey R. 2003. The Origin and Significance of Antarctic Meteorites. *Geochemistry* 63:93-147.

560 Heck P. R. et al. 2019. Best practices for the use of meteorite names in publications. *Meteoritics*
561 *& Planetary Science* 54:1397–1400.

562 Howard, E.C., Bournon, J.-L.Comte De, 1802. Experiments and observations on cer- tain stony
563 and metalline substances, which at different times are said to have fallen on the Earth. *Philos.*
564 *Trans. R. Soc. Lond.* 92:168–175.

565 Hutchison R., Alexander C. M. O'D., and Barber D. J. 1987. The Semarkona meteorite: first
566 occurrence of smectite in an ordinary chondrites and its implications. *Geochimica et*
567 *Cosmochimica Acta* 51:1875-1882.

568 Keil K. et al. 2015. The Vicência meteorite fall: A new unshocked (S1) weakly metamorphosed
569 (3.2) LL chondrite. *Meteoritics & Planetary Science* 50: 1089–1111.

570 Kita N. T., and Ushikubo T. 2012. Evolution of protoplanetary disk inferred from ^{26}Al
571 chronology of individual chondrules: Disk evolution and ^{26}Al chronology of chondrules.
572 *Meteoritics & Planetary Science* 47:1108–1119.

573 Lupker M. et al. 2012. Predominant floodplain over mountain weathering of Himalayan
574 sediments (Ganga basin). *Geochimica et Cosmochimica Acta* 84:410–432.

575 Marrocchi Y., Bekaert D. V., and Piani L. 2018. Origin and abundance of water in carbonaceous
576 asteroids. *Earth and Planetary Science Letters* 482:23–32.

577 Marrocchi Y., Villeneuve J., Jacquet E., Piralla M., and Chaussidon M. 2019. Rapid
578 condensation of the first Solar System solids. *Proceedings of the National Academy of*
579 *Sciences of the United States of America* 116:23461–23466.

580 Marvin U.B. 2007. Ernst Florens Friedrich Chladni (1756-1827) and the origins of modern
581 meteorite research. *Meteoritics & Planetary Science* 42:B3-B68.

582 Maurel C., Weiss B. P., Bryson J. F. J. 2019. Meteorite cloudy zone formation as a quantitative
583 indicator of paleomagnetic field intensities and cooling rates on planetesimals. *Earth and*
584 *Planetary Science Letters* 513:166-175.

585 McNaughton N. J., Fallick A. E., and Pillinger C. T. 1982. Deuterium enrichments in type 3
586 ordinary chondrites. *Journal of Geophysical Research* 87:A297.

587 Miller M. F., Franchi I. F., Sexton A. S. and Pillinger C. T. 1999. High precision $\delta^{17}\text{O}$ isotope
588 measurements of oxygen from silicates and other oxides: methods and applications. *Rapid*
589 *Communication in Mass Spectrometry* 13:1211–1217.

590 Piani L., Robert F., and Remusat L. 2015. Micron-scale D/H heterogeneity in chondrite
591 matrices: A signature of the pristine solar system water? *Earth and Planetary Science Letters*
592 415:154–164.

593 Potin S., Brissaud O., Beck P., Schmitt B., Magnard Y., Correia J.-J., Rabou P., and Jocou L.
594 2018. SHADOWS: a spectro-gonio radiometer for bidirectional reflectance studies of dark
595 meteorites and terrestrial analogs: design, calibrations, and performances on challenging
596 surfaces. *Applied Optics* 57:8279.

597 Quirico E., Raynal P.-I., Bourot-Denise M. 2003. Metamorphic grade of organic matter in six
598 unquellibrated ordinary chondrites. *Meteoritics & Planetary Science* 38:795-811.

599 Raynal P.-I., Quirico E., Borg J., Deboffle D., Dumas P., d 'Hendecourt L., Bibring J.-P., and
 600 Langevin Y. 2000. Synchrotron infrared microscopy of micron-sized extraterrestrial grains.
 601 *Planetary and Space Science* 48:1329–1339.

602 Robert F, Merlivat L, and Javoy M. 1979. Deuterium concentration in the early Solar System:
 603 hydrogen and oxygen isotope study. *Nature* 282:785–789.

604 Rubin A. E., Scott E.R.D, and Keil K. 1982. Microchondrule-bearing clast in the Piancaldoli
 605 LL3 meteorite: a new kind of type 3 chondrite and its relevance to the history of chondrules.
 606 *Geochimica et Cosmochimica Acta* 46:1763–1776.

607 Sears D. W., Grossman J. N., Melcher C. L., Ross L. M., and Mills A. A. 1980. Measuring
 608 metamorphic history of unequilibrated ordinary chondrites. *Nature* 287:791–795.

609 Stephant A., Garvie L. A. J., Mane P., Hervig R., and Wadhwa M. 2018. Terrestrial exposure
 610 of a fresh Martian meteorite causes rapid changes in hydrogen isotopes and water
 611 concentrations. *Scientific Reports* 8:12385.

612 Vacher L. G., Marrocchi Y., Verdier-Paoletti M. J., Villeneuve J., and Gounelle M. 2016.
 613 Inward radial mixing of interstellar water ices in the solar protoplanetary disk. *The*
 614 *Astrophysical Journal Letters* 827:1–6.

615 Vacher L.G., Piani L., Rigaudier T., Thomassin D., Florin G., Piralla M., and Marrocchi Y.
 616 (2020). Hydrogen in chondrites: Influence of parent body alteration and atmospheric
 617 contamination on primordial components. *Geochimica et Cosmochimica Acta* 281:53-66.

618 Van Orman J. A., Cherniak D. J., and Kita N. T. 2014. Magnesium diffusion in plagioclase:
 619 Dependence on composition, and implications for thermal resetting of the ^{26}Al – ^{26}Mg early
 620 solar system chronometer. *Earth and Planetary Science Letters* 385:79–88.

621 Van Schmus W.R., and Wood J.A. (1967) A chemical-petrologic classification for the
 622 chondritic meteorites. *Geochimica et Cosmochimica Acta* 31:747-765.

623 Villeneuve J., Marrocchi Y., and Jacquet E. (2020) Silicon isotopic compositions of chondrule
 624 silicates in carbonaceous chondrites and the formation of primordial solids in the accretion
 625 disk. *Earth and Planetary Science Letters* 542:116318.
 626 Wasson J. T., and Kallemeyn G. W. 1988. Compositions of Chondrites. *Philosophical*
 627 *Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*
 628 325:535–544.
 629 Yang J., and Epstein S. 1983. Interstellar organic matter in meteorites. *Geochimica et*
 630 *Cosmochimica Acta* 47:2199–2216.
 631

Fig. 1

634

635

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. 8

Fig. 9

Fig. 10

Fig. 11

Fig. 12