Open Research Online

The Open University's repository of research publications and other research outputs

Augmentation of clozapine with ECT: a retrospective case analysis

Journal Item

How to cite:

Lally, John; Breese, Emily; Osman, Mugtaba; Hua Sim, Cai; Shetty, Hitesh; Krivoy, Amir and MacCabe, James H (2021). Augmentation of clozapine with ECT: a retrospective case analysis. *Acta neuropsychiatrica*, 33(1) pp. 31–36.

For guidance on citations see [FAQs](#).

© 2020 Scandinavian College of Neuropsychopharmacology

<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Version: Accepted Manuscript

Link(s) to article on publisher's website:
<http://dx.doi.org/doi:10.1017/neu.2020.32>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

Augmentation of clozapine with ECT: a retrospective case analysis

Authors

John Lally*⁺¹⁻⁴, Emily Breese*⁵, Mugtaba Osman², Cai Hua Sim¹, Hitesh Shetty⁶, Amir Krivoy^{1,7}, James H MacCabe^{1,8}

*Both are first named authors and should be acknowledged as such.

Affiliations:

¹ Department of Psychosis Studies, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK

² Department of Psychiatry, Royal College of Surgeons in Ireland, Dublin, Ireland

³ Department of Psychiatry, Mater Misericordiae University Hospital, Eccles St., Dublin, Ireland

⁴ Department of Psychiatry, St Vincent's Hospital Fairview, Dublin, Ireland

⁵ Open University (School of Life, Health and Chemical Sciences, The Open University, Walton Hall, Milton Keynes, UK)

⁶ BRC Case Register, South London and Maudsley NHS Foundation Trust, London, UK

⁷ Sackler Faculty of Medicine, Tel-Aviv University, Ramat-Aviv, Israel

⁸ National Psychosis Service, South London and Maudsley NHS Foundation Trust, London, UK

*Dr John Lally MB MSc MRCPsych, PhD

Department of Psychosis Studies, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK; Honorary Clinical Lecturer, Department of Psychiatry, Royal College of Surgeons in Ireland, Dublin, Ireland; Consultant Psychiatrist, Department of Psychiatry, Mater Misericordiae University Hospital, Eccles St., Dublin, Ireland; St Vincent's Hospital Fairview, Dublin, Ireland and

Email: john.lally@kcl.ac.uk

*Ms Emily Breese, MSc,

Open University (School of Life, Health and Chemical Sciences, The Open University, Walton Hall, Milton Keynes, UK)

Email: emily.breese@open.ac.uk

Dr Mugtaba Osman MRCPsych

Department of Psychiatry, Royal College of Surgeons in Ireland, Beaumont Hospital, Dublin, Ireland

Email: mugtabasulman@yahoo.co.uk

Ms Cai Hua Sim MSc,

Department of Psychosis Studies, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK

Email: cai_hua.sim@kcl.ac.uk

Mr Hitesh Shetty, MSc

Clinical Informatician, BRC Case Register, South London and Maudsley NHS Foundation Trust, London, UK

Email: Hitesh.shetty@slam.nhs.uk

Dr Amir Krivoy, MD

Department of Psychosis Studies, Institute of Psychiatry, Psychology & Neuroscience (IoPPN), King's College London and Sackler Faculty of Medicine, Tel-Aviv University, Ramat-Aviv, Israel

Email: amir.krivoy@kcl.ac.uk

Prof James H MacCabe BSc MBBS FRCPsych MSc PhD

Professor of Epidemiology and Therapeutics, of Psychosis, Department of Psychosis Studies, Institute of Psychiatry, Psychology & Neuroscience (IoPPN), King's College London and Honorary Consultant Psychiatrist, National Psychosis Service, South London and Maudsley NHS Foundation Trust, London, UK

Email: james.maccabe@kcl.ac.uk

Corresponding Author:

*Dr John Lally MB, MSc, MRCPsych, PhD

Department of Psychosis Studies, Institute of Psychiatry, Psychology and Neuroscience, King's College London, PO63, De Crespigny Park, London SE5 8AF, UK (e-mail: john.lally@kcl.ac.uk).

Word count:

Abstract 222

Manuscript: 2900

Abstract

Objective

We sought to assess the effectiveness of clozapine augmentation with ECT (C+ECT) in patients with clozapine-resistant schizophrenia.

Methods

We conducted a retrospective review of electronic health records to identify patients treated with C+ECT. We determined the response to C+ECT and the rate of rehospitalisation over the year following treatment with C+ECT.

Results

Forty-two patients were treated with C+ECT over a ten year period. The mean age of the patients at initiation of ECT was 46.3 (SD=8.2) years (range 27-62 years). The mean number of ECTs given was 10.6(SD = 5.3)(Range 3-25) with the majority receiving twice weekly ECT. Seventy six percent of patients (n=32) showed a Clinical Global Impression improvement score (CGI-I) of ≤ 3 (at least minimally improved) following C+ECT. The mean number of ECT treatments was 10.6 (SD = 5.3) (range 3-25) with the majority receiving twice weekly ECT. Sixty four percent of patients experienced no adverse events. Response to C+ECT was not associated with gender, age, duration of illness, or duration of clozapine treatment.

Seventy five percent of responders remained out of hospital over the course of one-year follow up, while 70% of those with no response to C+ECT were not admitted to hospital. Three patients received maintenance ECT, one of whom was rehospitalised.

Conclusion

This study lends support to emerging evidence for the effectiveness of C+ECT in clozapine resistant schizophrenia. These results are consistent with the results of a meta-analysis and the only randomised controlled trial (RCT) of this intervention. Further RCTs are required before this treatment can be confidently recommended.

Keywords:

Schizophrenia; treatment-resistant; electroconvulsive therapy; psychosis;clozapine

Significant outcomes

Highlights

- Our study adds to the small extant literature indicating that clozapine augmentation with ECT (C+ECT) is associated with a high response rate, with 76% showing an acute response
- Our study findings indicate that the response to C+ECT is maintained in a majority of patients (75% remained out of hospital) with continued clozapine use over 1 year follow up (although 70% of non responders were not admitted to hospital)
- This is the largest retrospective study of C+ECT use in TRS and provides further evidence of the effectiveness of C+ECT in those with clozapine resistant schizophrenia

Limitations

Retrospective study, with no blinding, lack of a control arm and concurrent medication use was not controlled

Longer term follow up studies with larger sample sizes receiving M+ECT are required to evaluate the effectiveness of M+ECT in clozapine resistant schizophrenia

It was not possible to evaluate patient perspectives and experiences of ECT

Introduction

Treatment resistant schizophrenia (TRS) is the most disabling of all psychiatric illnesses and affects approximately 30% of those diagnosed with schizophrenia (Meltzer 1997, Lally *et al.*, 2016a). Clozapine remains the treatment of choice for people with TRS, with approximately 50-60% of patients improving with clozapine (Meltzer 1992, Agid *et al.*, 2011). There remain no evidence based pharmacological treatments for the remaining 40-50% of treatment resistant patients who fail to respond to clozapine (Taylor 2017, Lally and MacCabe 2015) (Lally and Gaughran 2019).

Meta-analyses have failed to demonstrate efficacy for clozapine augmentation with psychotropic medication, including antipsychotics and mood stabilisers (Sommer *et al.*, 2012). However, a recent randomised controlled trial (RCT) of clozapine augmentation with ECT (C+ECT) in clozapine-resistant schizophrenia showed encouraging results (Petrides *et al.*, 2015). Combining the results of that RCT with those of all other cases reported in the literature, we calculated the response rate for C+ECT in TRS to be 66% (Lally *et al.*, 2016b). Another meta-analysis demonstrated that ECT combined with non-clozapine antipsychotics was superior to antipsychotic monotherapy in TRS (Zheng *et al.*, 2016), supporting the findings of an earlier Cochrane review on ECT in schizophrenia which concluded that ECT is effective in schizophrenia and may have a place in patients who have failed to respond to other treatments (Tharyan and Adams 2005). The most recent Cochrane review of ECT for clozapine resistant schizophrenia found moderate evidence for improved clinical with C-ECT use compared to standard care (Sinclair *et al.*, 2019). These studies suggest a benefit of ECT in TRS, in the short term, but data on longer-term outcomes are almost completely lacking (Grover *et al.*, 2019). A systematic review of the use of maintenance ECT (M-ECT) in non TRS, identified that while quality evidence is lacking, M-ECT has some evidence for relapse prevention in combination with antipsychotics, with minimal evidence for persistent cognitive deficits (Ward *et al.*, 2018). It is also unclear whether maintenance ECT is needed to maintain improvement following response to C+ECT in TRS.

Study aims

The primary aim of this retrospective study was to describe the short term outcomes in a cohort of patients treated with C+ECT. The secondary aims were: 1) to explore the medium-term outcome of patients who received C+ECT based on the rate of psychiatric rehospitalisation in the year following a successful response to C+ECT, 2) to examine and identify post-ECT treatments which maintain clinical response, 3) to describe the use of maintenance ECT in those initial responders to C+ECT and 4) to evaluate adverse effects associated with C+ECT.

Methods

This was a retrospective observational study performed using data from the Clinical Record Interactive Search (CRIS) database from the South London and Maudsley (SLaM) NHS Foundation Trust.

In this study, we described the response to treatment in patients receiving ECT to augment clozapine (i.e. clozapine is continued during ECT). The primary outcome was the overall improvement in symptoms based on the Clinical Global Impression improvement (CGI-i) scale (**Guy** (1976)).

Data Source

Serving over 1.2 million residents located in four London boroughs (Croydon, Lambeth, Lewisham and Southwark), the SLaM NHS Foundation Trust is the largest provider of mental healthcare in Europe. Clinical records dating back to 1 January 2007 are stored in a combination of structured free text fields. In August 2016 there were over 200,000 individuals records.

The Clinical Record Interactive Search (CRIS) system was developed by the National Institute for Health Research Biomedical Research Centre located within the SLaM NHS trust to enable the automated search and retrieval of anonymised patient records. Data from structured text fields or from free text within the clinical records (e.g. clinical progress notes,

discharge summaries, outpatient correspondence) was extracted. Comprehensive details regarding this data resource can be found in an open access publication.(Stewart *et al.*, 2009)

Ethics and CRIS approval

Oxfordshire Research Ethics Committee C granted ethical approval to CRIS as an anonymised data resource for secondary analysis in 2008. A patient-led oversight committee is responsible for the governance for all CRIS project, who report to the SLaM Caldicott Guardian. This study carried CRIS application number 16-018 and was approved on 17th March 2016.

Inclusion and exclusion criteria

Primary inclusion and exclusion

The primary inclusion criterion was any person who received ECT to augment clozapine response, as part of a treatment plan for schizophrenia, or schizoaffective disorder (ICD-10 codes: F20.0, F25.0, respectively), and whose records contained sufficient information to retrospectively rate their response to treatment and had at least one year's follow up data. The treating clinicians defined an inadequate clozapine response and identified the need for ECT augmentation. The decision to begin and end ECT was made by the Consultant Psychiatrist responsible for the care of the patient, in consultation with the treating team and the patient. There was no age range criterion.

Figure 1 shows the flowchart for inclusion in the study.

Data extraction and Outcome variables

The first ECT treatment registered for each patient admitted during this 10-year period was defined as the index treatment. The following data were obtained for each patient: gender, age, ethnicity, diagnosis, duration of illness, duration of clozapine treatment, duration of

current episode (the episode that warranted ECT treatment), indication for ECT treatment, duration, number and frequency of ECT treatments, clozapine dose concurrent to ECT, other concurrent medications at time of ECT, outcome of ECT treatment, adverse events, subsequent use of maintenance ECT, subsequent prescription of maintenance medications (including subsequent prescription of clozapine, including dose changes and clozapine augmentation), subsequent rehospitalisation post response to C+ECT, and ECT treatment details.

The patient clinical records did not contain any standardised assessments of severity of illness or degree of improvement. CGI scores were retrospectively assigned based on documented clinical symptoms, psychopathology, behaviour, and functioning identified from the clinical records for 1 week prior to the administration of ECT and for one week after the end of the acute ECT course. As symptoms and behaviour can vary over the course of a week; the score was assigned based on the average severity level documented across the seven days. The CGI scores were rated by one of the researchers (EB) and were then checked for reliability by a clinician (JL). Consensus was sought where any differences existed and a final CGI score was recorded. We compared the mean and median CGI-severity (CGI-S) scores at the start and end of C+ECT.

A dichotomous response criterion based on the Clinical Global Impression (CGI) - Improvement scale was recorded for each ECT treatment case, using records from baseline (i.e. prior to C+ECT) and at the end of the ECT course. The CGI-Improvement (CGI-I) scale score was recorded following the end of C+ECT to assess the overall improvement and response (or not) to C+ECT. Any improvement (a rating of 1, 2 or 3) was seen as a successful response and a decline or no change (anything rated 4, 5, 6 or 7) was deemed unsuccessful (4= no change seen).

Statistical analysis

Descriptive statistics were performed using summary statistics (mean, standard deviation, median and range) for quantitative variables and frequencies for binary and nominal variables. The associations between demographic and clinical variables and response to C+ECT were assessed using t tests, or chi-square tests as appropriate, with a two-tailed alpha of 0.05.

Results

Demographic and clinical characteristics

A total of 42 patients (21 males) were treated with C+ ECT. The mean age of the patients at initiation of ECT was 46.3 (SD=8.2) years (range 27-62 years). Out of the total 42 participants, 29 (69%) were of white ethnicity, 12(29%) of black ethnicity and one (2%) was of mixed ethnicity.

Thirty-five (83%) patients had a diagnosis of schizophrenia and 7 (17%) had a diagnosis of schizoaffective disorder.

The mean clozapine dose during C+ECT was 383.7 (SD=211.6) mg. The mean duration of illness and clozapine administration were 5.5 (SD=5.0) years (median =4.3 years; range=.26-20.6 years) and 2.8 (SD=3.2) years (median 2.8 years; range =0.08-14.9 years) respectively. The mean duration of the current psychotic episode was 9.2 (SD=13.3) months (range =11 days-4.6 years). The median duration was 3.3 months.

ECT and clinical outcomes

The clinical indication for ECT augmentation in all cases was an insufficient response to clozapine (n=42/42). All patients received bilateral ECT (the stimulus dose was not consistently recorded). The mean number of ECT treatments was 10.6 (SD = 5.3) (median=10) (Range 3-25 ECT treatments (the majority of patients received twice weekly

ECT (n=17)). The mean duration of ECT treatment was 8.4 (SD=7.0) weeks. Females received a higher mean number of ECT treatments (mean=12.4 (SD=5.6) compared to males (mean =8.7 (SD=4.4)) ($t(35)=2.221, p=0.033$).

Thirty two patients (76%) responded to C+ECT (CGI-I ≤ 3). Eleven (26%) were 'very much' or 'much' improved following C+ECT. The CGI-I stratified by gender and for the total group is shown in table 1.

INSERT TABLE 1 HERE

There was a significant improvement in mean CGI-s scores following C+ECT (pre C+ECT =4.8 (SD=0.8) and post C+ECT 3.8 (SD=0.93) ($t=7.117, df=41, p<0.001$). There was no significant difference in mean clozapine dose during C+ECT in those who responded to C+ECT (mean dose= 379.4 (SD=223.6) mg) compared those who did not respond (mean dose =397.5 (177.3) mg) ($t=0.234, p=0.816$). Associations between clinical characteristics and age, and response to C+ECT are shown in table 2.

INSERT TABLE 2 HERE

Adverse events

Most individuals did not have any adverse effects secondary to ECT (64% (27/42) with no recorded adverse effects). Nine patients reported short term memory loss concurrent to ECT treatment which was transient for all and occurred at the time of the C+ECT.

Medication use concurrent to C+ECT

The response to C+ECT in relation to concurrent medications used at the time of C+ECT is shown in table 3. The use of anticonvulsant medication or benzodiazepines concurrent to C+ECT was not associated with a decreased response to C+ECT.

INSERT TABLE 3 HERE

Rehospitalisation post ECT response

Of the 32 patients who responded favourably to C+ECT, there were (n=5, 16%) who required rehospitalisation during the following year. Seventy five percent (n=24) of responders were not hospitalised over the course of the one year follow up (and 3 remained hospitalised during the follow up year). Thirty percent (n=3) of those with a non-response to C+ECT were rehospitalised over the follow up period.

Time to rehospitalisation (n=29 who were discharged post response to C+ECT)

Five of those who responded to C+ECT were rehospitalised due to illness relapse over the next year. The mean time to rehospitalisation was 106.8 days (range =24-280 days) and the median was 83 days.

In the year following response to C+ECT, 21% (n=6) of responders were treated with clozapine combined with antidepressant medication, 31% (n=9) were treated with clozapine augmented with mood stabilisers, 69% (n=18) had a dose increase in clozapine over the course of the year following ECT

There was no significant association between the use of clozapine combined with mood stabilisers ($\chi^2=2.719$, $p=0.131$), antidepressant ($\chi^2=1.576$, $p=0.283$) or M-ECT use ($\chi^2=0.607$, $p=0.446$) and rehospitalisation.

In total, 21 of the responders to C+ECT had either a dose increase in clozapine (n=17), were treated with M-ECT (n=1), were treated with both a clozapine dose increase and M-ECT (n=1), or were rehospitalised (with no clozapine dose increase or M-ECT use) (n=2).

Discussion

Response to C+ECT

Seventy six percent of patients had a positive response to C+ECT, a similar response rate to that identified in other retrospective case series (66%(Grover *et al.*, 2017)- 69%(Kim *et al.*,

2017)); though slightly lower than the 90% of patients who had an excellent or very good response in the study of Kristensen *et al.*, (Kristensen *et al.*, 2011). Our response rate is slightly higher than the previously identified pooled response rate of 66% to C+ECT in all extant published data (Lally *et al.*, 2016b). Thirty four percent of responders were very much or much improved following C+ECT.

Rehospitalisation post C+ECT

The one year rehospitalisation rate was 17% in those who were discharged post response to C+ECT, compared to 30% in those who did not respond to C+ECT, indicating that the acute response to C+ECT was associated with a low rehospitalisation rate over the following year. A recent Taiwanese mirror-image study using a national healthcare database, investigated the rates of psychiatric hospitalisation, following antipsychotic augmentation with ECT in patients with schizophrenia receiving ECT for the first time. They identified that antipsychotic augmentation with ECT was associated with a reduced rate of psychiatric hospitalization in the year following treatment, a finding that was particularly significant for those treated with C+ECT (n=906) (Lin *et al.*, 2017).

M+ECT and rehospitalisation during follow up

During follow up, 72% of responders had a dose increase in clozapine, were commenced on maintenance ECT (M-ECT) or were rehospitalised over the year following C+ECT. Dose increases in clozapine may have been indicative of re-emerging psychotic symptoms, though only one of these patients was rehospitalised (and this coincided with the commencement of M-ECT). We were unable to identify any significant relationship between concurrent medication use during the follow up period and reduced rates of hospitalisation, though it is of interest that all those patients treated with clozapine and mood stabilisers, or with antidepressants were not rehospitalised in the year following C+ECT. The proportion of responders who were later treated with M-ECT is similar to that identified in the follow up

study of Grover et al., (Grover *et al.*, 2017) though fewer than the 35% identified in a Danish retrospective chart review (Kristensen *et al.*, 2011).

The low rate of rehospitalisation following a response to C+ECT, is equivalent to the maintained response rates of 66% shown in the retrospective review of Grover et al., (though they do not comment on dose changes in clozapine, rather documenting that 66% maintained their improvement while concurrently treated with clozapine) (Grover *et al.*, 2017), or the 65% of those who did not require M-ECT for relapse of symptoms in the Kristensen et al., retrospective chart review (Kristensen *et al.*, 2011). An earlier open label study of 11 patients treated with C+ECT identified a relapse rate of 63% over an average follow up of 14 weeks following response to C+ECT (Kho *et al.*, 2004).

Only 3 of those who responded to C+ECT received M-ECT, with only one of those requiring hospitalisation for the administration of the maintenance treatment. Those treated with M-ECT were treated with 10-20 ECT treatments, two of whom were treated with weekly ECT, and the third patient was treated with weekly M-ECT for seven weeks followed by twice monthly ECT for 7 treatments. Due to the small number of patients who received M-ECT, we are unable to comment on the effectiveness of the adjunctive treatment in our population. Open label studies have provided support for sustained treatment effects with M-ECT. A previous open label study (n= 23) identified that 48% of those with clozapine resistant schizophrenia had a clinical response to acute C+ECT, and subsequently found an improved sustained clinical response with M-ECT compared to those treated with acute ECT and no maintenance ECT alongside continued clozapine (Youn *et al.*, 2019). An open pilot study (n=14) providing M-ECT for those who responded to ECT for clozapine resistant schizophrenia as part of a cross over RCT, showed no deterioration, and the clinical response was sustained for all who received M-ECT for a 6 month period (n=6, completed course of 10 C+ECT treatments in 6 months following acute C+ECT) (Braga *et al.*, 2019).

ECT treatment and associations with response

The mean number of ECT treatments was 10.6, consistent with that identified in the retrospective chart review of C+ECT of Kristensen et al., (=10) (Kristensen *et al.*, 2011), and the mean number of 11.3 identified in a recent meta-analysis of C+ECT (Lally *et al.*, 2016b). The number of ECT treatments was not significantly associated with treatment response or non-response. The duration of illness or the current psychotic episode was not associated with the likelihood of a patient responding, similar to findings of a retrospective chart review of the use of ECT in schizophrenia (Kristensen *et al.*, 2011).

Adverse events

The use of C+ECT was generally well tolerated with most patients reporting no adverse events, although this may be an underestimate due to underreporting of adverse events and under-recording in the clinical records. There was no evidence of prolonged seizures in any of the patients, though transient memory problems were reported in 21%, these were not sustained. A limitation of this study is the lack of formal cognitive testing performed, preventing a quantitative assessment of any cognitive changes over the course of C+ECT. One patient was found to have evidence of an underlying pre-existing neurocognitive disorder, which was identified during C+ECT.

Limitations

The findings from this naturalistic cohort, while demonstrating that C+ECT is effective for over 3/4s of patients, needs to be viewed considering certain limitations. The retrospective nature of the study design and the absence of a comparison group limit the conclusions that can be drawn. Furthermore, it was not possible to blind the ratings, so there is a possibility of observer bias. The retrospective nature of assigning clinical assessment scores, and the lack of prospective measures of clinical state such as PANSS or BPRS is a further limitation and potential source of observer bias. The CGI-S and CGI-I scales are brief assessment tools, that allow for retrospective assessment of clinical states and improvement, though which may not adequately assess the severity of psychotic symptoms. However, the CGI

assess both symptom severity and functioning, which may make it more relevant to clinical practice than the PANSS or BPRS that assess symptom severity only. The small sample size may have limited the power of our study to make statistical inferences and raising the possibility of type II errors, though it remains one of the largest clinical cohorts assessing the use of C+ECT in clinical practice. The lack of a control group limited the ability to conduct statistical analysis of clinical effects. The lack of neurocognitive assessments is a limitation, particularly since 21% of patients reported some transient memory loss concurrent to C+ECT.

There are limited studies investigating the response to C+ECT in clozapine refractory TRS, and even fewer studies with longer term follow up data. (Kristensen *et al.*, 2011, Grover *et al.*, 2017, Petrides *et al.*, 2015, Meltzer 1997, Braga *et al.*, 2019) Our findings indicate that C+ECT is associated with a high response rate, and is well tolerated. The use of clozapine dose increases was a commonly used treatment intervention in the year post C+ECT; though no specific medication intervention was associated with a significantly reduced risk of rehospitalisation. Our study adds to the small, but growing literature indicating that C+ECT may be an effective treatment for clozapine refractory schizophrenia, although our results need to be viewed with caution in the light of the methodological limitations.

Contributors

Authors Lally, Breese and MacCabe conceptualized the study and authors Lally, Breese and MacCabe contributed to the design.

Authors Breese, Hua Sim, Shetty and Lally conducted the data collection.

Authors Lally and Osman undertook the statistical analysis.

Authors Lally and Breese wrote the first draft of the manuscript. All authors (Lally, Breese, Osman, Hua Sim, Shetty, Krivoy, MacCabe) contributed to content revision and have approved the final manuscript.

Acknowledgement

None

Financial support

Dr MacCabe is supported by CRESTAR (CRESTAR project, <http://www.crestar-project.eu/>) EU-FP7 grant number 279227; Dr MacCabe is supported by STRATA (MRC grant no. MR/L011794/) and by the National Institute for Health Research (NIHR) Biomedical Research Centre for Mental Health at the South London and Maudsley NHS Foundation Trust and Institute of Psychiatry, Psychology and Neuroscience, Kings College London.

The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

The funding sources did not partake in the design, data collection or analysis.

Statement of Interest

None

References

- Agid, O., Arenovich, T., Sajeev, G., Zipursky, R. B., Kapur, S., Foussias, G. & Remington, G.** (2011). An algorithm-based approach to first-episode schizophrenia: response rates over 3 prospective antipsychotic trials with a retrospective data analysis. *Journal of Clinical Psychiatry* **72** (11), 1439-44.
- Braga, R. J., John, M., Schooler, N. R., Bailine, S. H., Malur, C., Mendelowitz, A. & Petrides, G.** (2019). Continuation Electroconvulsive Therapy for Patients With Clozapine-Resistant Schizophrenia: A Pilot Study. *The Journal of ECT* **35** (3), 156-60.
- Grover, S., Chakrabarti, S., Hazari, N. & Avasthi, A.** (2017). Effectiveness of electroconvulsive therapy in patients with treatment resistant schizophrenia: A retrospective study. *Psychiatry Research* **249** 349-53.
- Grover, S., Sahoo, S., Rabha, A. & Koirala, R.** (2019). ECT in schizophrenia: a review of the evidence. *Acta Neuropsychiatr* **31** (3), 115-27.
- Guy, W.** (1976). *ECDEU Assessment Manual for Psychopharmacology*, U.S. Department of Health, Education, and Welfare, Public Health Service, Alcohol, Drug Abuse, and Mental Health Administration, National Institute of Mental Health, Psychopharmacology Research Branch, Division of Extramural Research Programs.
- Kho, K. H., Blansjaar, B. A., De Vries, S., Babuskova, D., Zwinderman, A. H. & Linszen, D. H.** (2004). Electroconvulsive therapy for the treatment of clozapinenonresponders suffering from schizophrenia. *European Archives of Psychiatry and Clinical Neuroscience* **254** (6), 372-9.
- Kim, H. S., Kim, S. H., Lee, N. Y., Youn, T., Lee, J. H., Chung, S., Kim, Y. S. & Chung, I. W.** (2017). Effectiveness of Electroconvulsive Therapy Augmentation on Clozapine-Resistant Schizophrenia. *Psychiatry Investigation* **14** (1), 58-62.
- Kristensen, D., Bauer, J., Hageman, I. & Jorgensen, M. B.** (2011). Electroconvulsive therapy for treating schizophrenia: a chart review of patients from two catchment areas. *European Archives of Psychiatry and Clinical Neuroscience* **261** (6), 425-32.
- Lally, J., Ajnakina, O., Di Forti, M., Trotta, A., Demjaha, A., Kolliakou, A., Mondelli, V., Reis Marques, T., Pariente, C., Dazzan, P., Shergil, S. S., Howes, O. D., David, A. S., Maccabe, J. H., Gaughran, F. & Murray, R. M.** (2016a). Two distinct patterns of treatment resistance: clinical predictors of treatment resistance in first-episode schizophrenia spectrum psychoses. *Psychological Medicine* **46** (15), 3231-40.
- Lally, J. & Gaughran, F.** (2019). Treatment resistant schizophrenia - review and a call to action. *Irish Journal of Psychological Medicine* **36** (4), 279-91.
- Lally, J. & Maccabe, J. H.** (2015). Antipsychotic medication in schizophrenia: a review. *British Medical Bulletin*.

- Lally, J., Tully, J., Robertson, D., Stubbs, B., Gaughran, F. & Maccabe, J. H.** (2016b). Augmentation of clozapine with electroconvulsive therapy in treatment resistant schizophrenia: A systematic review and meta-analysis. *Schizophr Res* **171** (1-3), 215-24.
- Lin, H.-T., Liu, S.-K., Hsieh, M. H., Chien, Y.-L., Chen, I. M., Liao, S.-C., Tsai, H.-J. & Wu, C.-S.** (2017). Impacts of Electroconvulsive Therapy on 1-Year Outcomes in Patients With Schizophrenia: A Controlled, Population-Based Mirror-Image Study. *Schizophrenia Bulletin* sbx136-sbx.
- Meltzer, H. Y.** (1992). Treatment of the neuroleptic-nonresponsive schizophrenic patient. *Schizophrenia Bulletin* **18** (3), 515-42.
- Meltzer, H. Y.** (1997). Treatment-resistant schizophrenia--the role of clozapine. *Current Medical Research and Opinion* **14** (1), 1-20.
- Petrides, G., Malur, C., Braga, R. J., Bailine, S. H., Schooler, N. R., Malhotra, A. K., Kane, J. M., Sanghani, S., Goldberg, T. E., John, M. & Mendelowitz, A.** (2015). Electroconvulsive therapy augmentation in clozapine-resistant schizophrenia: a prospective, randomized study. *American Journal of Psychiatry* **172** (1), 52-8.
- Sinclair, D. J. M., Zhao, S., Qi, F., Nyakyoma, K., Kwong, J. S. W. & Adams, C. E.** (2019). Electroconvulsive therapy for treatment-resistant schizophrenia. *Cochrane Database of Systematic Reviews* (3).
- Sommer, I. E., Begemann, M. J., Temmerman, A. & Leucht, S.** (2012). Pharmacological augmentation strategies for schizophrenia patients with insufficient response to clozapine: a quantitative literature review. *Schizophrenia Bulletin* **38** (5), 1003-11.
- Stewart, R., Soremekun, M., Perera, G., Broadbent, M., Callard, F., Denis, M., Hotopf, M., Thornicroft, G. & Lovestone, S.** (2009). The South London and Maudsley NHS Foundation Trust Biomedical Research Centre (SLAM BRC) case register: development and descriptive data. *BMC Psychiatry* **9** (51), 9-51.
- Taylor, D. M.** (2017). Clozapine for Treatment-Resistant Schizophrenia: Still the Gold Standard? *CNS Drugs* **31** (3), 177-80.
- Tharyan, P. & Adams, C. E.** (2005). Electroconvulsive therapy for schizophrenia. *Cochrane Database of Systematic Reviews* (2).
- Ward, H. B., Szabo, S. T. & Rakesh, G.** (2018). Maintenance ECT in Schizophrenia: A Systematic Review. *Psychiatry Research* **10.1016/j.psychres.2018.03.033**.
- Youn, T., Jeong, S. H., Kim, Y. S. & Chung, I. W.** (2019). Long-term clinical efficacy of maintenance electroconvulsive therapy in patients with treatment-resistant schizophrenia on clozapine. *Psychiatry Research* **273** 759-66.
- Zheng, W., Cao, X. L., Ungvari, G. S., Xiang, Y. Q., Guo, T., Liu, Z. R., Wang, Y. Y., Forester, B. P., Seiner, S. J. & Xiang, Y. T.** (2016). Electroconvulsive Therapy Added to Non-Clozapine Antipsychotic Medication for Treatment Resistant Schizophrenia: Meta-Analysis of Randomized Controlled Trials. *PLoS One* **11** (6), e0156510.

Table 1. Outcome of clozapine augmentation with ECT measured by CGI scores

CGI-Improvement Score	1- <i>Very much improved</i>	2- <i>Much improved</i>	3- <i>Minimally improved</i>	4- <i>No change</i>	5- <i>Minimally worse</i>	6- <i>Much worse</i>	7- <i>Very much worse</i>
Men	1	4	9	5	2	0	0
Women	0	6	12	2	1	0	0
Total	1	10	21	7	3	0	0
Proportion	2.4%	23.8%	50%	16.7%	7.1%	0%	0%

Table 2. Clinical characteristics and response to C+ECT (defined as improvement (a rating of 1, 2 or 3) on Clinical Global Impression (CGI) -Improvement scale score)

	Response (N=32)	Non response (N=10)	T-test; p-value
Age (SD)	46.6 (8.5)	45.5 (7.6)	0.363; 0.718
Duration of illness in years(SD)	5.9 (5.0)	4.3 (5.1)	0.849; 0.401
Duration of clozapine treatment in years (SD)	2.6 (3.3)	3.5 (3.2)	0.698; 0.490
Duration of current psychotic episode (months)(SD)	9.5 (13.4)	8.3 (14.5)	0.246; 0.807
Mean number of ECT treatments (SD)	9.9 (4.7)	12.5 (6.6)	1.356; 0.184

Table 3 Medication use and association with response to C+ECT

	Response n= (%)	No Response n= (%)	χ^2 ; p
Anticonvulsant use			
Yes	13 (40.6)	2 (20.0)	1.412; 0.212
No	19 (59.4)	8 (80.0)	
Anticonvulsant and benzodiazepine use			
Yes	16 (51.6)	3 (30.0)	1.420; 0.205
No	15 (48.4)	7 (70.0)	
Antipsychotic augmentation of clozapine			
Yes	19 (59.4)	4 (40.0)	1.155; 0.238
No	13 (40.6)	6 (60.0)	
Antidepressant use			
Yes	6 (18.8)	3 (30.0)	0.573; 0.362
No	26 (81.3)	7 (70.0)	
Lithium carbonate use			
Yes	9 (29.0)	3 (30.0)	0.003; 0.622
No	22 (71.0)	7 (70.0)	