Open Research Online

The Open University's repository of research publications and other research outputs

Recovering lost hay meadows: An overview of floodplain-meadow restoration projects in England and Wales

Journal Item

How to cite:

Rothero, Emma; Tatarenko, Irina and Gowing, David (2020). Recovering lost hay meadows: An overview of floodplain-meadow restoration projects in England and Wales. *Journal for Nature Conservation*, 58, article no. 125925.

For guidance on citations see [FAQs](#).

© 2020 Elsevier GmbH

<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Version: Accepted Manuscript

Link(s) to article on publisher's website:

<http://dx.doi.org/doi:10.1016/j.jnc.2020.125925>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

Recovering lost hay meadows:

An overview of floodplain-meadow restoration projects in England and Wales

Corresponding author Emma Rothero^a, emma.rothero@open.ac.uk

Irina Tatarenko^a, irina.tatarenko@open.ac.uk David Gowing^a david.gowing@open.ac.uk

^aSchool of Environment, Earth and Ecosystem Sciences, Open University, Gass Building, Walton Hall, Milton Keynes MK7 6AA, England

Declaration of Interest: None

Funding sources: The survey activities and data analyses were undertaken by the Floodplain Meadows Partnership team, funded by the John Ellerman Foundation. The John Ellerman Foundation is a philanthropic trust who fund work in part relating to conservation activity in the UK. They had no part in the study design, report writing, data analysis or paper production/submission. They funded staff time to undertake data collection and landowner interviews, resulting in the production of a report to the funder about activity undertaken during the survey period (2016-2018).

Abstract

Restoration of natural habitats plays an important role in nature conservation. After 30 years of efforts to restore species-rich floodplain meadows, most of which had been lost to alternative land uses during the preceding half century, the extent and level of restoration success in the UK remained unknown. A survey of floodplain meadow-restoration projects across England and Wales from 2016-2018 allowed evaluation of restoration progress on 163 sites nationwide.

Restoration success was measured by floristic composition, species richness and balance of functional types in plant communities. To identify factors affecting restoration success, their state prior to restoration, restoration technique, site ownership and quality of ongoing management were analysed. The survey revealed that 25% of restoration sites demonstrated expected success, achieved mainly by private landowners. Restoration failed or showed very poor progress on another 15% of sites, managed predominantly by public or charitable organisations. The remaining sixty percent of the sites showed some signs of improvement. The degree of restoration success showed no significant correlation to the state of the site prior to restoration, or to the restoration method applied. Ownership of the site and site management both influenced restoration success. The degree of success appeared to depend on the consistency and sufficiency of the restoration management.

Keywords

Floodplain meadow, plant community, biodiversity, sustainability, ecological restoration, success criteria, long-term management.

Open access

Subscription/green open access (embargo period 24 months).

Introduction

Many northern European floodplains have been occupied by semi-natural meadows for most of the last millennium. These diverse communities established under long-term consistent management for hay balancing the nutrient input from floods. These meadows developed as sustainable ecosystems, highly valued for animal feed over many centuries (Schaich et al, 2011; McGinlay et al 2016). More than 97% of these habitats were lost from the floodplains of Europe during the period 1930-1990 (e.g. Fuller, 1987; Krause et al, 2011), and to a lesser extent in the following years (Jefferson, 2014). Moist or wet, mesotrophic to eutrophic hay meadows (Eunis habitat type E2.14 and E3.4a, EEA 2019) are classed as Endangered in the European Red List of habitats (Janssen et al,

2016) and the importance of floodplain meadow restoration has been recently acknowledged globally by the European Environment Agency (2016) and United Nations Environment Programme (2019). Floodplain meadows are increasingly recognised for the range of ecosystem services they offer (Lawson et al, 2018), and their loss has prompted the initiation of many projects aiming to re-create and restore them (e.g., Vinther & Hald 2001, Hölzel & Otte 2003).

Globally, habitat restoration processes have become so widespread that ecosystem restoration targets were set by the United Nations Environment Programme (2019). The increasing number of restoration projects creates a need for critical evaluation of the processes affecting them (Jones *et al* 2019). In the UK, some positive outcomes have been reported for particular floodplain meadow sites in England (e.g., Woodcock et al 2011, Hosie et al 2019) and Wales (Shelswell & Squire, 2019). However, estimation of restoration extent and progress on floodplain meadows at a national scale has never been previously attempted. A three-year nationwide survey was initiated in the UK by the Floodplain Meadows Partnership (FMP; www.floodplainmeadows.org) in 2016 (Rothero & Tatarenko, 2018). Over three years, 177 restoration sites were visited in 20 counties across England and Wales, representing 844 hectares of floodplain. Seventeen sites where restoration projects were about to start were also visited. Sixty seven percent of the surveyed sites were entered into an agri-environment scheme. This UK government funded scheme (known as Countryside Stewardship) provides financial incentives for farmers, land owners or managers to maintain and improve the environment (Rural Payments Agency, 2020)()

Restoration methods have varied over the past 30 years of floodplain-meadow restoration in England and Wales. The most common method in the last ten years has been the application of freshly cut herbage (Kiehl et al, 2010), colloquially referred to as “green hay,” taken from an existing species-rich meadow. “Green hay” refers to species-rich herbage from a donor site that is

transferred to the restoration site on the same day it was cut (Edwards et al 2007, Kirkham et al 2012). However, other methods are also used including sowing commercial seed mixtures, reverting permanent pastures to a regime of hay cutting, application of seed mixtures collected from existing species-rich meadows by hand, often with the help of volunteers, along with planting plug plants, or simply a change of management from pasture to hay cut and aftermath grazing. In this survey, botanical data were collected on each site, together with information about historical and current management and restoration techniques, in order to explore the degree of success of restoration projects. Measuring success or even monitoring progress of restoration projects is a challenge, because there is no definitive approach (Kimball et al 2015). Vegetation structure, species diversity, species abundance, presence of target species and functioning of ecological processes are commonly used (Ruiz-Jaen & Aide 2005). In the UK, the success of restoration sites involved in agri-environment schemes is measured by both species' richness (as number of species per 1 m²), and by frequency of particular indicator species selected from targeted plant communities (Natural England, 2012; Natural England, 2016). Assessing the similarity of a restored community to a target community has rarely been undertaken. In the UK, targets for floodplain-meadow restoration have focussed on the rare, species-rich *Sanguisorba officinalis*-*Alopecurus pratensis* (MG4) grassland and the *Cynosurus cristatus*-*Caltha palustris* grassland (MG8) as defined by the British National Vegetation Classification (NVC) (Rodwell, 1992). The match between the vegetation on a restored site and a target community can be measured by calculating a similarity coefficient (Czekanowsky, 1913; Malloch, 1996.) This degree of similarity was applied here as the first criterion of restoration success. Apart from the similarity in floristic composition, a restored community is expected to mirror the functionality of the target community. The C-S-R system of describing plant functional types (Grime, 1974) has been successfully used to compare vegetation samples which differed in management regime (Hunt et al, 2004). The C-S-R signature (Hunt et al 2004), was used as the second criterion in our calculation of restoration success. The combination of these two criteria together with species richness, as a general indicator of biodiversity, were adopted to produce a robust scale of restoration success.

As the success of a project is defined by its results, so assessments of project management are usually based on the achievements of pre-set goals and objectives (Hockings et al, 2006). For example, the management of restoration sites in agri-environment schemes is assessed by the presence of indicator species in the sward (Natural England, 2012). The spectrum of methods that have been used to assess the management of protected areas is broad in scale and the degree to which the criteria are formalised is broad too (Stoll-Kleemann 2010.) However, the role of management in restoration projects has rarely been discussed (e.g. Guerrin, 2015). The socio-economic component of management is even more rarely considered because of the variability of management approaches, especially in the private sector, the lack of records of management activities, and the poor structure and ambiguity of questionnaires collecting management data (McGinlay et al 2016). Restoration projects are expected to be managed in accordance with restoration guidelines and advice (e.g., Natural England 2012). However, the dynamic nature of floodplain environments, e.g. variation in soil nutrients (Klaus et al 2011) and flooding regime (Gerard et al 2008), brings additional challenges for restoration managers. Our approach seeks to explore whether management traits, such as consistency, sufficiency and adaptability help to explain the trajectory of restored vegetation.

Our survey combined information from a diversity of landowners and restoration approaches, with other factors, such as the condition of the site prior to restoration to evaluate variation in the effectiveness of restoration.

Within this paper, we aim to address two questions:

How close are we to restoring lost hay meadows on British floodplains?

Has a particular habitat restoration approach proven successful?

124

125 Our second question gave rise to four separate hypotheses:

- 126 - Restoration success is a function of site ownership
- 127 - Restoration success depends on the restoration technique used
- 128 - Restoration success depends on site condition prior to restoration
- 129 - Restoration success correlates with the consistency, sufficiency or adaptiveness of
- 130 management

131

132 **Materials and methods**

133 *Identifying restoration locations.*

134 The diversity of organisations, landowners and other bodies involved in floodplain-meadow
135 restoration projects in the UK is extensive, but their identification was made feasible by using the
136 network already developed by the Floodplain Meadows Partnership (FMP.) A layered approach was
137 used to identify them:

- 138 • Data supplied under licence from the Government adviser for the natural environment in
139 England, Natural England (Natural England, n.d.), was entered into a Geographic Information
140 System (QGIS association, Switzerland) and used to find sites that were registered in the
141 Countryside Stewardship Scheme. We focussed on land entered under option GS6 [Restoration
142 of species-rich, semi-natural grassland] or GS7 [Creation of species-rich, semi-natural grassland]
143 and located within floodzone 2 of the floodplain (Environment Agency, 2020) The sites identified
144 were then followed up through local Natural England staff, who were able to contact landowners,
145 and in some cases help with access.
- 146 • proactively approaching local contacts (e.g. The Natural England Grassland Network, Local
147 Wildlife Trusts, Local Biodiversity Record Centres) to request information on floodplain-meadow
148 restoration activity in their areas

- advertising the study through the FMP newsletter and social media.

Collection of physical and management information from sites

The following physical data were collected:

- On 127 sites, five 1 m x 1 m quadrats were surveyed per site, listing all plant species present and their percentage cover. Quadrats were placed randomly across each field. Positions of the quadrats were recorded with Mobile Topographer GPS app (Google, 2020) with an accuracy to 0.5 m. Quadrat locations were shown on maps provided to the landowners together with survey results.

Thirty-six further fields were surveyed on a walk-through basis due to time constraints, where species lists were recorded rather than quadrat data collected. All species seen along a walk across the field were recorded with estimates of their abundance and frequency.

- The landowner or manager was questioned about site management and restoration methods (Appendix 1 – Site Assessment Form). In some cases, follow-up emails and telephone calls were required to fill data gaps.

- Of the 177 sites visited, information from fourteen sites (111 ha) remained incomplete, so they were excluded from the data analysis.

Statistical tests

. The Kruskal Wallis H test was applied to address the following hypotheses:

- Restoration success is a function of site ownership
- Restoration success depends on the restoration technique used
- Restoration success depends on site condition prior to restoration

The Kruskal Wallis test was chosen because restoration success was assessed on an ordinal scale, whilst the multiple potential explanatory variables were categorical. Where significant effects were indicated, a Mann-Whitney U-test was used as a *post hoc* test. .

The Spearman rank correlation test was used to address the following hypotheses:

- Restoration success correlates with the consistency of management
- Restoration success correlates with the sufficiency of management
- Restoration success correlates with the adaptiveness of management

The Spearman rank test was chosen because management regimes were scored subjectively on an ordinal scale, the success of restoration and the management regime were paired at site level and visualisation of the data suggested a monotonic relationship.

The number of sites in each success category entered into an agri-environment scheme was analysed using a chi-square contingency table. This approach was taken because both variables were categorical.

Theory and calculations

In order to estimate restoration success, we summed measures of species richness, similarity to target community and functional diversity.

To evaluate characteristics of restoration management (management factors), questionnaire responses were scored against criteria set out below.

Definition and measure of restoration success

In this study, several measures in combination are proposed as a robust approach to assessing restoration success:

197 1. Species richness

198 Species richness is a frequently used measure of habitat quality (e.g. Ruiz-Jaen & Aide 2005).
199 On restoration sites, it has been widely used as a criterion of success (e.g. Natural England,
200 2012). The targeted vegetation types in this context (MG4, MG8 and MG5), tend to be species
201 rich with typically more than 20 plant species per 1 m² (Wallace & Prosser 2016). However, if
202 the number of species is boosted by the presence of ruderal species, as is often the case during
203 the early phases of a restoration scheme, this measure alone can be unreliable.

204 2. Similarity to National Vegetation Classification communities

205 For many of the restoration projects, a specific restoration target was set at the outset;
206 namely to achieve a sward composition as described by the British National Vegetation
207 Classification (Rodwell, 1992). The Modular Analysis of Vegetation Information System
208 (MAVIS; CEH, 2016) was used to measure the similarity coefficient between observed and
209 target communities. Smaller samples were taken than recommended for NVC survey, to be
210 consistent with existing data held for these grassland types (Wallace and Prosser, 2016).
211 MAVIS uses the same form of the Czekanowski coefficient as an earlier software application,
212 MATCH (Malloch, 1996). In this study, a similarity coefficient of over 60% was assumed to
213 represent a good fit to the particular NVC type. Here we followed a methodological study by
214 Semkin (2009,) which demonstrated that exceeding a threshold coefficient of 59% was
215 indicative of high similarity between communities. Scores below 50% were assumed not to
216 conform to the target plant community (Dodd et al, 1994). Scores between 50 and 60%
217 suggest community re-assembly is partially achieved (Tatarenko et al, 2018). Similarity to
218 target communities alone was not considered sufficiently sensitive to evaluate restoration
219 progress fully and therefore additional measures were included.

220 3. C-S-R Functional Types

221 The high diversity of floodplain meadows can be explained by hydrological niche segregation
222 in space and time (Silvertown et al, 1999). This heterogeneity can promote both taxonomic

and functional diversity. The latter has been proposed as a finely tuned characteristic of restoration success, although it is more difficult to measure (England & Wilkes, 2018; Jones et al, 2019). Based on C-S-R functional types, as defined by Grime (1974), a C-S-R signature was suggested as a tool for comparison of herbaceous vegetation (Hunt et al, 2004). Species with competitive (C) or ruderal (R) life-strategies tend to occupy newly cleared areas, such as restored arable fields, much faster than species with a stress-tolerant (S) strategy. The latter group tend to perform poorly in the early stages of vegetation succession (Pywell et al, 2003). Meadow communities restored in tundra, after 15 years, had low S-value in functional evenness compared to C and R-values: e.g. C:S:R 0.38:0.24:0.40 (Novakovsky & Panyukov, 2018). Established species-rich meadow communities are characterised by an even spread of CSR functional types. For example, in alpine hay meadow C:S:R was found as 0.36: 0.30: 0.34 (Onipchenko et al, 2020). Based on the arguments above, we suggest using evenness of C:S:R scores in measuring restoration success. C-S-R signatures as defined by Hunt et al (2004), were calculated in MAVIS for each quadrat, and averaged per site as C:S and S:R ratios.

Therefore, a combination of four measures: species richness, similarity to target community, C:S ratio and S:R ratio, were used to develop a quantitative scale for evaluating restoration success. The scale ranges from 1 (failure), to 5 (success), with 2, 3, and 4 marking different levels of progress in between (Table 1).

The scale was developed in discussion with several grassland ecologists experienced in the classification of plant communities in English and Welsh floodplain meadows.

For fields surveyed by a walk-through, the assessment of restoration progress was in accordance with guidelines in the Countryside Stewardship manual (Natural England, 2016) which includes floodplain meadow plant communities defined within the lowland meadows broad habitat type. The combination of presence and frequency of common and indicator species was used to calculate

restoration success scores, as shown in Table 1. The thresholds were set to create an even spread of sites across the range of scores.,

Table 1. Measures of success for restoration of floodplain meadows.

	Measure of success				
Measure	1 Failure	2	3	4	5 Success
Average scores from five botanical quadrats per field as calculated in MAVIS					
Species richness	<8	8-12	13-15	16-20	>20
NVC similarity score	<50%	50-55%	55-60%	>60%	>60%
C:S ratio (average)	>1.39	1.27-1.39	1.18-1.27	1.10-1.18	<1.10
S:R ratio (average)	<0.79	0.79-0.81	0.81-0.84	0.84-0.89	>0.89
Scores based on walk-through survey using Countryside Stewardship manual					
Number and frequency of species deemed to be -indicators of success	0	1-2 occasional	4-5 occasional	1-4 frequent	>4 frequent
Number of species common to lowland meadows	0-2	3-5	6-7	>7	>7

Role of management in restoration success in dynamic floodplain-meadow ecosystems

How management is delivered differs. We define three characteristics of management, which we use to assess the management regime for each site (Appendix 2). Firstly, consistency of management defined as an annual midsummer hay cut with prompt removal of hay and followed by a period of

aftermath grazing. These practices have been proposed as important for the development and maintenance of a species-rich floodplain meadow (Gerard et al, 2008, Poptcheva et al, 2009) particularly with respect to balancing the soil-nutrient budget (Rothero *et al*, 2016). Secondly, sufficiency of management captures whether the degree of restoration effort is planned and tailored to the specific site to ensure the target can be achieved. On floodplains, lack of management may result in flood waters failing to drain (Leyer, 2004), nutrients accumulating (Timmermann et al, 2015, Fry et al, 2017), propagules being too sparse (Carter & Blair, 2012) or germination being sub-optimal (Abbandonato et al, 2018). All these problems would be noticeable from the early stages of restoration projects as would others such as weed infestation, poaching of the soil and ingress of scrub along boundaries. A third characteristic – adaptiveness of management - reflects the dynamic nature of floodplain habitats (e.g. unpredictable spring and summer floods,) which require adaptive management in response to the condition of the habitat from month to month (Lemke et al 2017).

To identify the management factors involved in the success or otherwise of a restoration project, we determined for each site whether the management was: A – consistent, B – sufficient, and C – adaptive. Information gathered about management during the site visit was compiled and structured along a three-point scale, using the criteria set out in Table 2. A degree of subjectivity was required, but sites for which there was too little information to make an informed judgement about the consistency, sufficiency and adaptiveness of the management were omitted from the analysis.

Table 2. Management factors affecting restoration projects for floodplain meadows

Categories	1	2	3
A – Consistency	Inconsistent: Timing of hay cuts delayed beyond when	Partially consistent:	Consistent: A timely annual hay cut Timely hay removal

	<p>hay was ready. Hay cut missed in some years.</p> <p>Cut material not collected promptly when dry.</p> <p>Lack of grazing in autumn.</p> <p>Overgrazing causing soil compaction</p>	Occasional cases of inconsistency as listed in column 1	Regular grazing with its pressure determined by the availability of grass.
<p>B – Sufficiency</p>	<p>Insufficient</p> <p>Hay making too late or too incomplete to balance nutrient inputs.</p> <p>Drainage infrastructure insufficient to avoid periodic soil anoxia</p> <p>Cover of litter allowed to accumulate.</p> <p>Too few propagules introduced and/or seed introduction methods failed to optimise germination</p>	<p>Partially sufficient</p> <p>Some aspects of the factors mentioned in column 1.</p>	<p>Sufficient:</p> <p>Nutrient input and outputs are in balance over a flood cycle.</p> <p>Flood waters leave site quickly enough to avoid anoxia.</p> <p>Well managed grazing or occasional harrowing to avoid litter accumulation</p> <p>Seed material of good quality, sown at an appropriate density and onto adequately prepared ground.</p> <p>Propagules introduced in more than one year.</p>

C –	Inflexible:	Partially adaptive:	Fully adaptive:
Adaptiveness	Following published protocol (e.g. Countryside Stewardship Manual) to the letter without consideration of local conditions. For example, no additional hay cut in years when nutrient availability peaks after a major flood. No re-application of propagules if initial recruitment is poor.	Making some limited adjustments to the restoration plan to reflect conditions on the ground.	Taking management decisions based upon monitoring data, prevailing site conditions and weather.

276

277 **Results**

278

279 Extent of floodplain-meadow restoration across the UK

280 Data collected from 163 restoration sites in 20 counties across England and Wales (Table 3)

281 representing 733 hectares of floodplain grassland, were included in the data analysis. 70% of

282 restoration projects started between 2006 and 2015, 12% occurred during 1990s, and 6% in the

283 period 2000-2005. Some decline in the number of new restoration projects was observed in recent

284 years. Restoration projects were found in most of the major river floodplains in England, most

285 notably the Thames, Severn, Ouse, Wye, Kent, and Ribble (Fig. 1). A number of sites were found

286 along lakesides or along the coast. The highest density of restoration sites was found in the Thames

287 catchment, in proximity to some of the oldest and best-preserved floodplain meadows, such as

North Meadow and Clattinger Farm (Wiltshire), and Oxford Meadows, Ducklington Mead and Chimney Meadows (Oxfordshire). Restoration sites along the river Thames have started to fill the gaps in between existing species-rich meadows, in some areas starting the forming of a 'flower-rich corridor' along the river valley (Fig. 2). In the North of England, restoration sites were found in connection with existing floodplain-meadow systems in North Yorkshire, such as Clifton Ings and Rawcliffe Meadows along the river Ouse, and the flat landscapes of Lincolnshire, which historically supported vast areas of former fen used for agricultural purposes, including hay production. There are very few remaining species-rich floodplain meadows in the area, but the 38 restoration sites from that one county cover 53 ha, or 8% of the total area, and comprise 23 % of the total number of sites visited. The hilly landscapes of the Lake District with its smaller, flashier river valleys, where remaining traditional hay meadows are again very limited in extent, revealed 16 sites covering 31 ha. Wales has few floodplains considered suitable to support classic floodplain meadows, with only a very small handful of existing sites recorded in the country. A number of future restoration fields are located next to two of the three known MG4 Sites of Special Scientific Interest (SSSI) (Natural Resources Wales, 2020) in Wales – Old Pulford Brook Meadow, and Crabtree Green. Dolydd Hafren meadow restoration project (Montgomeryshire) is located in a very active river floodplain on very thin, young alluvial soils, and C'aer Ddol in Gwynedd is on the margins of Lake Padarn.

Figure 1. Map of restoration sites and existing species-rich meadows in England and Wales

Figure 2. Distribution of restoration sites and existing species-rich meadows along the Thames valley around Oxford, England.

Five percent of restoration sites surveyed in England and Wales were classified as fully ‘successful’ (category 5, see Table 1). Progress was substantial on 20% of sites marked as category 4 (Table 1). These two categories add up to 25% of the restoration sites which can be considered as meeting the expected “restored” status (Table 3 and Appendix 2). The target plant communities of MG4 and MG8 (Rodwell, 1992), as well as MG15, which is closely aligned to MG4 (Wallace & Prosser, 2016), were recorded. These restoration sites tended to be either more than 20-years old, or had received large and repeated applications of propagules.

More than half of all the sites were classified as being in a transitional state, demonstrating some progress towards formation of species-rich meadows (categories 2 and 3 with 27% and 34% of sites

respectively). They showed a tendency to develop grass-dominated communities, such as MG6, MG7C, MG7D, MG9 and MG10 (Rodwell, 1992). The remaining 15% of sites were evaluated as category 1 ('failure').

The number of restoration sites and their success outcome varied between counties in England and Wales (Table 3). The large variability in restoration success observed on the surveyed sites was analysed against a range of factors involved in restoration projects.

Table 3. Number of floodplain-meadow restoration sites surveyed in the counties in England and Wales in 2016-2018. A more complete summary of the data is presented in Appendix 2. Categories of restoration success range from 1 (failure) to 5 (success) as explained in Table 1.

County	Number of restoration sites surveyed in the county	Categories of restoration success				
		1	2	3	4	5
Berkshire	1	0	0	0	0	1
Buckinghamshire	17	1	3	10	2	1
Cambridgeshire	3	1	2	0	0	0
Clywd	2	0	0	1	1	0
Cumbria	16	0	3	6	7	0
Gloucestershire	10	3	2	3	2	0
Gwynedd	3	0	2	1	0	0
Hampshire	2	0	0	1	0	1
Herefordshire	2	0	2	0	0	0
Lancashire	3	1	1	0	1	0
Lincolnshire	26	4	7	6	7	2
Montgomeryshire	1	0	1	0	0	0
Northamptonshire	10	1	1	4	3	1
Oxfordshire	39	5	10	18	5	1
Shropshire	4	1	1	2	0	0
South Yorkshire	2	0	2	0	0	0
Staffordshire	5	2	2	1	0	0
Surrey	2	2	0	0	0	0
Wiltshire	5	0	3	1	0	1
Yorkshire	10	3	1	2	4	0
Total:	163	24	43	56	32	8

Factors affecting restoration success on the sites

a) *Site ownership*

Restoration projects in the UK were categorised by the type of landowner or manager responsible for them. Individual farmers and private landowners of mainly small fields attached to a house were grouped as “private landowners” (PL); they comprised 42% of all owners, managing 255 hectares of restoration sites. Wildlife charities, local councils, and local and national organisations were grouped in the category of “Public & Charitable ownership” (PO) and managed 43% of restoration sites covering an area of 386 ha. Private companies (PC) included represented 14% of owners, managing more than 90 ha. Community groups play a minimal role in the restoration of floodplain meadows at a national scale, representing only 1% of owners.

Success of restoration was found to be associated with ownership category; two thirds of sites within category 5 (success) belonged to private landowners (PL). Differences between PL and PO as the two major categories of land manager, were clear in category 1 (failure,) where the number of public and charitable organisations (PO) and the associated area managed were 3 times higher than that from private landowners (PL) (Fig. 3). Schemes managed by private companies (PC) and community groups (CG) were only represented in categories 2-4 (Fig. 3).

Figure 3. Proportion of site owners in each of the five categories of restoration success as defined in Table 1. PC – private companies, PL – private landowners, PO – public and charitable organisations, CG – community groups.

The success scores were found to differ significantly between three main types of landowners (Kruskal Wallis H test; $n=14, 51, 89$; for PC, PL and PO categories respectively; $H = 6.83$; $d.f = 2$; $p=0.03$). *Post hoc* Mann-Whitney tests comparing median values between PC ($n=14$) and PL ($n=51$) and between PO ($n=89$) and PL ($n=51$) both showed the PL value to be significantly higher.

b) *Site condition prior to restoration and method of restoration used*

The condition of sites prior to restoration showed a wide range of starting points (Fig. 4). Seven types of land use were changed to restore floodplain meadows. The majority of the sites (57%) were restored from some form of species-poor permanent grassland, such as pastures, old degraded meadows and agriculturally improved meadows. More than 200 hectares (27%) were restored from arable land. Land uses such as allotments and amenity grasslands were rare (0.3%) and not included

in the analysis for this reason. The success of restoration was not strongly determined by land-use prior to the restoration attempt (Kruskal Wallis H-test, $n=163$, $H=3.23$, $d.f.=3$, $p=0.36$).

Green hay application was used on 29% of sites, the majority of which were permanent grassland prior to restoration (Fig. 4). Some sites receiving green hay were first scarified, but this level of detail of restoration technique was not considered in the data analysis. Most sites restored by green hay fell into success categories 2 and 3.

Sowing commercial seed mixtures and reverting permanent pastures to a regime of hay cutting were applied on 23% of sites. The application of commercial seed mixtures was successful on half of the arable fields where it was applied, however the number of sites with insignificant progress and failure (categories 1 and 2) was also high (Fig. 5). A simple change of management was applied to pastures, where it resulted in success (category 5) on two fields (Fig. 5). Application of seed mixtures collected from existing species-rich meadows, along with planting plug plants, has been used occasionally, and was recorded on 13% of the sites in our survey. The success of these methods and their combination was variable across the different pre-restoration conditions of the sites (Fig. 5). Most sites that listed commercial seed and seed/plug planting as their restoration method, also had a change of management as part of the process, typically from year-round grazing to an annual hay cut and aftermath grazing. This restoration approach was called 'multiple' because several methods were used on the same site. Five main restoration techniques: ChM – changed management, CSM – commercial seed mixture, GH – green hay, WSM – wild seed mixture, and Multi – multiple methods, have been tested against categories of restoration success as defined in Table 1. The Kruskal Wallis test ($n=163$, $H=3.48$, $d.f.=4$, $p=0.48$) showed no significant difference in the effect of restoration method on the success of a restoration project.

Older methods not apparently used since the 1990's include dry hay application (3 sites) and natural regeneration (4 sites). The latter approach relied on propagules of meadow species arriving naturally

from neighbouring species-rich fields, but their swards remained species-poor 30 years later. Dry hay on its own resulted in failure or weak progress on two sites, however in combination with other restoration methods (plug plants, turf transfer and wild seed sowing) it led to a very good outcome at Copse Meadow along the river Ouse (York). Two sites were successfully restored from pasture by changes in management regime. A few sites successfully practised a mixture of techniques, e.g. green hay, commercial seed and plug planting, or dry hay, wild seeds and plug planting. In those cases, the application of different plant material was spread through several years, while in the majority of sites, plant propagules in one form or another were applied only once. About 6% of sites did not keep records of the restoration methods used (Appendix 2, Table 1).

Figure 4. Restoration success measured by categories 1-5 (as described in Table 1) on different pre-restoration land use types.

Figure 5. Restoration success measured by categories 1-5 (as described in Table 1) with different restoration techniques applied. Restoration techniques: ChM – changed management, CSM – commercial seed mixture, GH – green hay, WSM – wild seed mixture, PP - plug plants, DH – dry hay, Multi – multiple methods, Natural regeneration.

Impact of management on restoration success

Restoration success was measured using a 5-point scale (Table 1) and analysed against three characteristics of management using a 3-point scale (Table 2). Correlation between quality of management and restoration success was positive in all three categories described in Table 2. The strongest correlation (Spearman rank test, $\rho=0.77$; $p<0.001$) was found between sufficiency of management and restoration success. The coefficients from the same test for consistency ($\rho=0.49$; $p<0.001$) and adaptability ($\rho=0.28$; $p=0.002$) were lower, but nevertheless significant. The restoration success rate of the three main types of ownership using the three management characteristics described in Table 2 was tested using the Kruskal-Wallis H-test. It revealed that public

and charitable organisations (PO), private landowners (PL), and private companies (PC), had similar levels of management consistency (n=14, 51, 83; $H=1.36$, d.f.=2, $p=0.50$) and adaptiveness (n=14, 34, 67; $H=3.09$, d.f.=2, $p=0.21$), while sufficiency differed significantly (n=14, 51, 83; $H=8.13$, d.f.=2, $p=0.02$) between the three groups of owners (Fig. 6).

Figure 6. Median values of the management-regime scores of the three main ownership types: private companies (PC,) private landowners (PL) and Public and charitable organisations (PO.) Bars represent the mean and error bars the interquartile range.

Only 18% of all restoration projects were not run under a government-funded Agri- environment scheme. As shown in Fig. 7, sites participating in agri-environment schemes were well distributed across all five categories of restoration success, as were sites not in a scheme. A chi-square contingency table ($n=163$, $d.f.=4$, $p=0.76$) showed no significant effect of scheme membership on the distribution of sites between success categories.

Figure 7. Split of restoration sites within and outside of an agri-environment scheme.

Discussion

Spatial extent of floodplain meadow restoration

Restoration of semi-natural habitats, especially those which declined in the 20th century due to agricultural intensification, was thought to be an effective way of bringing them back (e.g., Wells et al, 1994). There are 2980 ha of species rich floodplain meadows remaining in England and Wales

(Rothero et al, 2016). The 844 hectares represented by restoration sites described here would extend these rare plant communities by 28% if successful, which would be a substantial contribution to their conservation. In some areas, for example the Thames river valley around Oxford, restoration sites have started to fill the gaps between existing species-rich meadows, forming 'wildflower rich corridors' (Fig. 2), which are important for the exchange of propagules and gene flow between fragmented habitats (Krause et al, 2011, Arponen et al, 2013). However, the extent of restoration projects does not determine how effective restoration has truly been in practice (Ramstead et al, 2012). Restoration success varied across the sites in this survey, as well as in most projects carried on wet meadows across Europe (Dicks et al, 2019). The challenges of grassland restoration have been discussed using examples of projects at different scales: from small fields (e.g., Hölzel & Otte, 2003, Smith et al, 2017), up to landscape-scale restoration projects across the world (Gerard et al, 2008, Nakamura et al, 2014, Guerrin, 2015, Lemke et al, 2017). After 20-60 years of restoration, no projects were recognized as fully successful and completed (e.g. Poptcheva et al, 2009, Fagan et al, 2010, Smith et al, 2017). In the UK by 2018, only twenty five percent of surveyed sites achieved restoration success (categories 4 & 5.) On half of the sites visited, the vegetation showed limited signs of progress towards desirable species-rich communities (Tatarenko et al, 2018). On fifteen percent of the sites in this survey, restoration has made little discernible progress and the process probably needs to start again. Below we discuss the limitations which may explain the slow progress with restoration of floodplain meadows in England and Wales.

How long does successful floodplain meadow restoration take?

Restoration projects of semi-natural landscapes and habitats are designed to bring back associations of organisms which used to form communities linked into specific ecosystems. Easily germinating and fast-growing grassland species are considered to be quick and easy to re-assemble into specific plant communities (Stevenson et al, 1995, Jongepierova et al, 2007). The majority of meadow restoration projects and trials have been carried out in dry grasslands (e.g., Hayes & Tallowin, 2007,

Edwards et al, 2007, Jongepierova et al, 2007), where vegetation tends to restore faster than in wet meadows (Galvnek & Lepš, 2009). Agri-environment schemes suggest that meadow restoration can be completed in 5-10 years, while in reality it can take much longer (Smith et al 2007, Poptcheva et al 2009; Woodcock et al, 2011) as the species re-assembly requires time for spatial and temporal niches to develop (Fagan et al, 2008).

The presence of positive indicator species on the site (e.g., Shelswell & Squire, 2019), is only the very first step in a long process of plant re-assembly into a community. After 22 years of good progress on Somerford Mead (Woodcock et al, 2011) and almost 10 years on Clattinger Farm (Hosie et al, 2019), target species are still very patchy. Many meadow species, particularly herbs, demonstrate a low rate of establishment from the existing seed bank, as well as poor natural dispersal (e.g. Bakker & Berendse, 1999; Pywell et al, 2002; Bossuyt, & Honnay, 2008). These considerations explain the slow progress on many restoration fields in this survey. Even though some target species were present in the fields, their distribution was very restricted. As a result, such vegetation gives low similarity scores with reference NVC targets (Tatarenko et al, 2018). Furthermore, the small number of sites in our survey that are undergoing restoration through natural regeneration, show little signs of improvement after 30 years, despite the close vicinity of existing species-rich meadows. Natural recovery of such sites can take many decades (Walker et al, 2004). For example, it was predicted that colonization by invertebrate species that characterize the target habitat type could take over 130 years (Woodcock et al, 2011).

Effectiveness of different restoration techniques

Data from 36 wet meadows in the Netherlands, Germany and the UK showed different success rates of the main restoration methods used on wet meadows (Klimkowska et al, 2007). Our data suggest

that on floodplain meadows, no one restoration technique was more successful than others. Responses to different seeding methods in the restoration of English lowland calcareous grasslands were also shown to be unclear (Fagan et al, 2010).

Topsoil removal was recognised as one of the most promising techniques to begin a restoration project on lowland meadows including those on the floodplains (Hölzel & Otte, 2003, Klimkowska et al 2009, Kiehl et al, 2010). In the UK, this method was used on two fields in York (sites 18 and 19, Appendix 2), which resulted in relatively good restoration success on those sites. Overall in the UK, topsoil removal is rarely used, possibly due to the high costs associated with this exercise.

Amongst the eight restoration methods used on British floodplain meadows, green hay spreading was most common. The green hay method was successful in trials (e.g. Edwards et al, 2007) and recommended for wider use by Natural England (2010).. However, no restoration sites ran germination tests to calculate transfer rates of species, which is required to evaluate the effectiveness of this technique (Kiehl et al, 2010). As a result, the application of green hay without control of propagules transferred within it, may not provide sufficient re-assurance of restoration success. Commercial seed mixtures (CSM) used for re-seeding grasslands, are variable in their origin, (Kiehl et al, 2010) and quality of seeds (Abbandonato et al, 2018). In the UK, CSM often originate from local sources, which is thought to be important for restoration success (Jongepierová et al, 2007, Schmidt et al, 2020). Local origin of CSM can explain the similarity in success rates (category 4) between CSM and wild seed mixtures used on restoration sites (Fig 5). However, CSM vary substantially in the number of species included (Török et al, 2011). Poor mixtures may have led to the high proportion of sites with no or poor restoration success (categories 1 and 2) in our survey (Fig 5).

A positive effect of a high diversity of seeding material and dense seeding has been shown to enhance establishment of species on sites (Carter & Blair, 2012; Manchester et al, 1999). In our survey, the number of sites that were re-seeded over several consecutive years was very small, so this treatment could not be considered independently in the analysis. Several individual sites (e.g. sites 113, 141, 163, Appendix 2) achieved steady restoration success (categories 4 and 5) via repeated application of propagules. Seed limitation is a well-recognised barrier to the recovery of temperate grasslands (e.g. Walker et al, 2004; Johnson et al, 2018). In semi-natural habitats, seed dispersal is an annual, repetitive process under hay cut management, where the date of the hay cut is critically important (Bischoff et al, 2018). Different meadow species have annual fluctuations in their abundance, sizes, and seed production (Pierce et al, 2014), so the amount of propagule available for germination varies from year to year. This variation can affect the quantity and quality of seeding material transferred to a restoration site (Kiehl et al, 2010, Bischoff et al, 2018) and restoration success in general. Multiple applications of propagules should ensure greater species richness in the restored field. This rarely used technique has good potential, which merits more research. However, multiple applications can be expensive and time-consuming, unsuitable for the goal of restoration of large areas at low cost (Liira et al, 2009). The project budget often defines the choice of restoration technique used (Török et al, 2011).

The role of pre-restoration condition of the field

Pre-restoration condition of the fields is a major factor to consider in restoration projects (Walker et al, 2004); Harvolk-Schoning et al, (2020) found that arable fields on floodplains can be restored more successfully than species-poor grassland. In our survey, about half the sites restored from arable use showed no or poor restoration success. The number of restored arable fields in the higher success categories was lower than permanent grasslands including pastures (Fig 4). Statistically, no significant differences in success were found between the four main types of pre-restoration land use (arable, improved pasture, unimproved pasture and species-poor hay meadow). Pre-restoration

soil treatment on the sites (e.g. harrowing) was not analysed in this paper. Soil disturbance has helped to explain differences in restoration success on ex-arable fields with bare soils (Kiehl et al, 2010). Access to the mineral soil surface promotes seed germination (Hellström et al, 2009, Réka et al, 2020), suggesting soil disturbance can be a more powerful factor than pre-restoration condition of the site, though the benefits may be short-lived (Harvolk-Schoning et al, 2020.)

Three qualitative approaches to management of restoration sites

Whilst physical condition of the site and the restoration technique used showed no significant influence on restoration success, the quality of restoration management (its consistency, sufficiency and adaptiveness) did play a significant role. The type of land ownership/management has been found to have an effect on the progress of restoration projects elsewhere (Stoll-Kleemann, 2010). In our survey, private landowners (PL) had a significantly higher rate of success compared to both public and charitable organisations (PO) and private companies (PC). A potential explanation that private landowners had smaller sites, which were easier to restore, was not supported, as the size of field had no significant effect on restoration success within any group of landowners. Another hypothesis, that private landowners have better funding opportunities, availability of resources for hay cut and aftermath grazing, as well as a better focus on their sites, was given some support because their management was deemed sufficient to a greater degree than other managers. The mean degree of consistency and adaptiveness of management showed no significant differences between the three main groups of land managers in our study, but institutional factors have been suggested as playing a role in the failure of restoration projects elsewhere (Guerrin, 2015).

Sufficiency of management showed a significant correlation with restoration success in general. High sufficiency (category 3, Table 2) specifies that all restoration measures were implemented effectively: enough propagules were introduced, nutrient availability was managed in advance and drainage infrastructure was maintained. Insufficiency of post-restoration management can include

reduced livestock grazing (e.g. Timmermann et al, 2015) and poor control of nutrients brought with flood sediments (Lemke et al, 2017). In our survey, consistency of management also showed a significant correlation with restoration success, reflecting the importance of the regular hay cut (Smith et al, 1996) and timely hay removal (Schaffers et al, 1998).

Restoration of floodplain ecosystems and their specific functions can suffer from a lack of knowledge of basic processes and dynamics, such as sedimentation and flood duration (Klaus et al, 2011). On dynamic floodplain landscapes, flexibility in approaching restoration projects appears to be key to success. More than half of the sites surveyed in the first year of our project showed very slow restoration progress because of high nutrient availability in the soil (Rothero & Tatarenko, 2018). Double hay cuts in June and September were found to be beneficial for soil-nutrient management in wet-meadow restoration projects in Germany (Poptcheva et al, 2009). In British meadows, timing of the hay cut is very restrictive for those within an agri-environment scheme, where mowing is often not permitted until after 15th July. An adaptive management (AM) approach (Zedler & Callaway, 2003) should be well suited to the management of such a dynamic ecosystem to allow the variation of environmental conditions across space and time to be considered (Kimball et al, 2015). The flexible, open approach to restoration practice advocated by Higgs et al (2018), is consistent with our findings here.

Agri-environment schemes are designed in part to reinstate past habitat losses, but their effectiveness has been variable (Arponen et al, 2013). The majority of sites in this survey (137 of 163) were entered into such a scheme, which incentivised the landowner to attempt restoration. However, according to our survey, participation in the scheme did not significantly enhance the likelihood of a successful outcome.

Conclusions

From a survey of 163 field sites, neither the restoration method nor the previous land use was found to affect restoration success in floodplain-meadow schemes. However, the category of ownership did influence the outcome, with schemes managed by private landowners being the most successful.

The aspect of a management regime that appeared to have the strongest correlation with success was a set of actions we classified as “sufficiency.” This aspect was a measure of the care and diligence taken by the scheme manager. These results suggest that the pathway by which a meadow is restored is of less importance than the care with which that management is applied. The manager needs to assess the efficacy of their own management and adjust their actions accordingly. Our findings suggest that following a pre-determined set of rules (a recipe) for restoration is not ideal. The manager needs to respond to what they can see on the ground. Where a manager lacks experience, the help of an adviser may be key.

The category of ownership and the sufficiency of management appear to be linked in as much as it was the private landowner category that scored most highly in terms of management sufficiency.

The survey has documented the extent of restoration activity in England and Wales and has demonstrated that the current resource of this valuable habitat is being increased by restoration schemes. However, the data were unable to show any effect of affiliation to an agri-environmental scheme on the outcomes.

Acknowledgements

All farmers and managers of the restoration sites for their kind permission to access their fields and for sharing their knowledge. Richard Jefferson, Andy Cooke and George Hinton (Natural England) for their help in supplying data. Mike Dodd (The Open University) for help with fieldwork and data analysis. Hilary Wallace (Ecological Surveys Bangor) for help with fieldwork and interpretation of

NVC data. Richard Jefferson, Mike Dodd and Hilary Wallace for their inputs into deriving a scale of restoration success. The John Ellerman Foundation for financially supporting this work for 3 years.

References

- Abbandonato, H., Pedrini, S., Pritchard, H.W., De Vitis, M. Bonomi, C. (2018). Native seed trade of herbaceous species for restoration: a European policy perspective with global implications. *Restoration Ecology* Vol. 26, No. 5, pp. 820–826
- Arponen, A., Heikkinen, R. K., Paloniemi, R., Pöyry, J., Similä, J., Kuussaari, M. 2013. Improving conservation planning for semi-natural grasslands: Integrating connectivity into agri-environment schemes. *Biological Conservation*, 160, 234–241.
- Bakker, J.P. & Berendse, F. (1999) Constraints in the restoration of ecological diversity in grassland and heathland communities. *Trends in Ecology & Evolution*, 14, 63–68.
- Bischoff, A., Hoboy, S., Winter, N., Warthemann, G. (2018). Hay and seed transfer to re-establish rare grassland species and communities: How important are date and soil preparation? *Biological Conservation* 221, 182–189.
- Bossuyt, B., & Honnay, O. (2008). Can the seedbank be used for ecological restoration? An overview of seedbank characteristics in European communities. *Journal of Vegetation Science*, 19, 875–884.
- Carter, D. L., & Blair, J. M. 2012. High richness and dense seeding enhance grassland restoration establishment but have little effect on drought response. *Ecological Applications*, 22(4), 1308–1319. <https://doi.org/10.1890/11-1970.1>
- Centre for Ecology and Hydrology, 2016. Modular Analysis of Vegetation Information System (MAVIS). <https://www.ceh.ac.uk/services/modular-analysis-vegetation-information-system-mavis>.

653 Czekanowski J (1913) Zarys Metod Statystycznych. Warsaw: E. Wendego.

654 Dicks, L.V., Ashpole, J.E., Dänhardt, J., James, K., Jönsson, A., Randall, N., Showler, D.A., Smith, R.K.,
655 Turpie, S., Williams D.R. & Sutherland, W.J. (2019) Farmland Conservation Pages 291-330 in: W.J.
656 Sutherland, L.V., Dicks, N., Ockendon, S.O., Petrovan & R.K. Smith (eds) *What Works in Conservation*
657 2019. Open Book Publishers, Cambridge, UK.

658 Dodd, M.E., Silvertown, J., McConway, K., Potts, J., Crawley, M. 1994. Application of the British
659 National Vegetation Classification to the communities of the Park Grass Experiment through time.
660 Folia Geobotanica & Phytotaxonomica, Vol. 29, No. 3, pp. 321-334

661 Edwards A.R., Mortimer S.R., Lawson C.S. Westbury D.B., Harris S.J., Woodcock B.A. & Brown V.K.
662 (2007) Hay strewing, brush harvesting of seed and soil disturbance as tools for the enhancement of
663 botanical diversity in grasslands. *Biological Conservation*, **134**, 372-382.

664 England, J., Wilkes, M.A. 2018. Does river restoration work? Taxonomic and functional trajectories at
665 two restoration schemes. *Science of the Total Environment*. 618. 961–970

666 Environment Agency (2020). Flood Map for Planning (Rivers and Sea) - Flood Zone 2 online. Available
667 at [https://data.gov.uk/dataset/cf494c44-05cd-4060-a029-35937970c9c6/flood-map-for-planning-](https://data.gov.uk/dataset/cf494c44-05cd-4060-a029-35937970c9c6/flood-map-for-planning-rivers-and-sea-flood-zone-2)
668 [rivers-and-sea-flood-zone-2](https://data.gov.uk/dataset/cf494c44-05cd-4060-a029-35937970c9c6/flood-map-for-planning-rivers-and-sea-flood-zone-2). Accessed 2nd September 2020.

669 European Environment Agency 2016. Flood risks and environmental vulnerability. Exploring the
670 synergies between floodplain restoration, water policies and thematic policies. EEA Report, N1,
671 2016, 1-84.

672 European Environment Agency (2019) Habitat factsheets E2.14 and 3.4.
673 <https://eunis.eea.europa.eu/habitats/1635> and <https://eunis.eea.europa.eu/habitats/158>. Accessed
674 8th June 2020.

675 Fagan, K. C., Pywell, R. F., Bullock, J. M., & Marrs, R. H. (2008). Do re-stored calcareous grasslands on
676 former arable fields resemble ancient targets? The effect of time, methods and environment on

677 outcomes. *Journal of Applied Ecology*, 45(4), 1293–1303. <https://doi.org/10.1111/j.1365->
678 2664.2008.01492.x

679 Fagan, K. C., Pywell, R. F., Bullock, J. M., & Marrs, R. H. (2010). The seed banks of English lowland
680 calcareous grasslands along a restoration chronosequence. *Plant Ecology*, 208:199–211.

681 Fry, E. L.; Pilgrim, E. S.; Tallowin, J. R.B.; Smith, R. S.; Mortimer, S. R.; Beaumont, D. A.; Simkin, J.;
682 Harris, S. J.; Sheil, R. S.; Quirk, H.; Harrison, K. A.; Lawson, C. S.; Hobbs, P.J. and Bardgett, R. D.
683 (2017). Plant, soil and microbial controls on grassland diversity restoration: a long-term, multi-site
684 mesocosm experiment. *Journal of Applied Ecology*, 54(5) pp. 1320–1330.

685 Fuller, R.M., 1987. The changing extent and conservation interest of lowland grasslands in England
686 and Wales: a review of grassland surveys 1930-1984. *Biological Conservation*, 40 (4), 281-300.

687 Galváneš, D. Lepš, J. (2009). How do management and restoration needs of mountain grasslands
688 depend on moisture regime? Experimental study from north-western Slovakia (Western
689 Carpathians) *Applied Vegetation Science* 12: 273–282.

690 Gerard, M., El Kahloun, M., Rymen, J., Beauchard, O. and P. Meire. 2008. Importance of mowing and
691 flood frequency in promoting species richness in restored floodplains *Journal of Applied Ecology* 45,
692 1780–1789

693 Google, 2020. Google Play Apps. Mobile Topographer Free online. Available at
694 https://play.google.com/store/apps/details?id=gr.stasta.mobiletopographer&hl=en_GB. Accessed
695 2nd September 2020.

696 Grime, J.P. 1974. Vegetation classification by reference to strategies. *Nature* 250: 26-31.

697 Guerrin, J. (2015). A floodplain restoration project on the River Rhone (France): analysing challenges
698 to its implementation. *Reg. Environ. Change*, 15:559–568.

699 Harvolk-Schoning, S, Michalska-Hejduk, D., Harnisch, M., Otte, A., Donath, T.W. (2020) Floodplain
 700 meadow restoration revisited: Long-term success of large-scale application of diaspore transfer with
 701 plant material in restoration practice. *Biological Conservation*, 241, 108322
 702 Hayes M.J. & Tallowin J.R.B. (2007) Recreating biodiverse grasslands: long-term evaluation of
 703 practical management options for farmers. Pages 135-140. in: Hopkins, J. J., Duncan, A. J.,
 704 Hellström, K. & Huhta, Ari-Pekka & Rautio, Pasi & Juha, Tuomi. (2009). Seed introduction and gap
 705 creation facilitate restoration of meadow species richness. *Journal for Nature Conservation*. 17.
 706 10.1016/j.jnc.2009.04.007.
 707 Higgs, E., Harris, J., Murphy, S., Bowers, K., Hobbs, R., Jenkins, W., Kidwell, J., Lopoukhine, N.,
 708 Sollereder, B., Suding, K., Thompson, A., Whisenant, S. 2018. On principles and standards in
 709 ecological restoration. *Restoration Ecology Vol. 26, No. 3, pp. 399–403*
 710 Hockings, M., Stolton, S. Leverington, F. Dudley, N. Courrau, J. 2006. Assessing Effectiveness—A
 711 Framework for Assessing Management Effectiveness of Protected Areas. Online.
 712 DO - 10.2305/IUCN.CH.2006.PAG.14.en
 713 Hölzel, N., Otte, A. 2003. Restoration of a species-rich flood meadow by topsoil removal and
 714 diaspore transfer with plant material. *Applied Vegetation Science* 6:131–140.
 715 Hosie, C., Jones, E., Rothero, E., & Wallace, H. 2019. Restoration of a floodplain meadow in Wiltshire,
 716 UK through application of green hay and conversion from pasture to meadow management.
 717 *Conservation Evidence* 16: 12–16.
 718 Hunt, R., Hodgson, J.G., Thompson, K., Bungener, P., Dunnett, N.P., Askew, A.P. (2004). A new
 719 practical tool for deriving a functional signature for herbaceous vegetation. *Applied Vegetation*
 720 *Science*, 7, 163-170.
 721 Janssen, J.A.M, Rodwell, J.S. García Criado, M. Gubbay, S., Haynes, T., Nieto, A., Sanders, N.,
 722 Landucci, F., Loidi, J., Ssymank, A., Tahvanainen, T., Valderrabano, M., Acosta, A., Aronsson, M., Arts,
 723 G., Attorre, F., Bergmeier, E., Bijlsma, R.-J., Bioret, F., Biță-Nicolae, C., Biurrun, I., Calix, M., Capelo, J.,
 724 Čarni, A., Chytrý, M., Dengler, J., Dimopoulos, P., Essl, F., Gardfjell, H., Gigante, D., Giusso del Galdo,

725 G., Hájek, M., Jansen, F., Jansen, J., Kapfer, J., Mickolajczak, A., Molina, J.A., Molnár, Z., Paternoster,
 726 D., Piernik, A., Poulin, B., Renaux, B., Schaminée, J.H.J., Šumberová, K., Toivonen, H., Tonteri, T.,
 727 Tsiripidis, I., Tzonev R., and Valachovič M., (2016). European red list of habitats. Part 2. Terrestrial
 728 and freshwater habitats Brussels. European Commissioner for the Environment .
 729 Jefferson, R.G., Smith, S.L.N. and MacKintosh, E.J. (2014). Guidelines for the Selection of Biological
 730 SSSIs. Part 2: Detailed Guidelines for Habitats and Species Groups. Chapter 3: Lowland Grasslands.
 731 Joint Nature Conservation Committee, Peterborough.
 732 Johnson, D.P., Catford, J.A., Driscoll, D.A., Gibbons, P. (2018). Seed addition and biomass removal key
 733 to restoring native forbs in degraded temperate grassland. *Appl Veg Sci.* 21:219–228.
 734 Jones, H.P., Barber, N.A. & Gibson, D.J. 2019. Is phylogenetic and functional trait diversity a driver or
 735 a consequence of grassland community assembly? *Journal of ecology*, 107: 2027-2032.
 736 <https://doi.org/10.1111/13652745.13260>
 737 Jongepierová, I., Mitchley, J. & Tzanopoulos, J. (2007) A field experiment to recreate species rich hay
 738 meadows using regional seed mixtures. *Biological Conservation*, 139, 297-305
 739 Kiehl, K., Kirmer, A., Donath, T.W., Rasran, L., Hölzel, N. (2010). Species introduction in restoration
 740 projects - evaluation of different techniques for the establishment of semi-natural grasslands in
 741 Central and Northwestern Europe. *Basic and Applied Ecology* 11: 285-299
 742 Kimball, S., Lulow, M., Sorenson, Q., Balazs, K., Fang, Y-C., Davis, S.J., O'Connell, M., T. and E.
 743 Huxman 2015. Cost-effective ecological restoration. *Restoration Ecology Vol. 23, No. 6, pp. 800–810*
 744 Kirkham, F. W., Bhogal, A., Chambers. B. J., Dunn, R. M., and Tallowin, J. R. B. (2012). Effects of
 745 spreading species-rich green hay on the botanical composition of an agriculturally improved
 746 hay meadow in northern England. *Grass and Forage Science*, 68, 260–270 doi: 10.1111/j.1365-
 747 2494.2012.00895.x

748 Klaus, V.H., Sintermann, J., Kleinebecker, T., and N. Holzel. 2011. Sedimentation-induced
 749 eutrophication in large river floodplains – An obstacle to restoration? *Biological Conservation* 144
 750 (2011) 451–458.

751 Klimkowska, A., Van Diggelen, R., Bakker, J.P. & Grootjans, A.P. (2007) A review of the success of
 752 commonly used wet meadow restoration methods (rewetting, topsoil removal, and diaspore
 753 addition) in Western Europe. *Biological Conservation*, 140, 318-328.

754 Klimkowska, A., Kotowski, W., van Diggelen, R., Grootjans, A.P., Dzierż, P., Brzezińska, K. (2009).
 755 Vegetation Re-development After Fen Meadow Restoration by Topsoil Removal and Hay Transfer.
 756 *Restoration Ecology* Vol. 18, No. 6, pp. 924–933 doi: 10.1111/j.1526-100X.2009.00554.x

757 Krause, B. Culmsee, H., Wesche, K., Bergmeier, E., Leuschner, C. 2011. Habitat loss of floodplain
 758 meadows in north Germany since the 1950s. [Biodiversity and Conservation](#). 20, 11, 2347–2364

759 Lawson, C., Rothero, E., Gowing, D., Nisbet, T., Barsoum N., Broadmeadow, S., Skinner, A., (2018)
 760 The natural capital of floodplains: management, protection and restoration to deliver greater
 761 benefits. *Valuing Nature Natural Capital Synthesis Report VNP09*.

762 Lemke, M.J., Hagy, H.M., Dungey, K., Casper, A.F., Lemke, A.M., Van Middlesworth, T. D., and Kent,
 763 A. (2017). Echoes of a flood pulse: short-term effects of record flooding of the Illinois River on
 764 floodplain lakes under ecological restoration. *Hydrobiologia* 804:151–175.

765 Leyer, I. 2004. Effects of dykes on plant species composition in a large lowland river floodplain *River*
 766 *Research and Applications*, Vol. 20(7), pp.813-827

767 Liira, J., Issak, M., Jõgar, Ü., Mändoja, M., Zobel, M., 2009. Restoration management of a floodplain
 768 meadow and its cost-effectiveness—the results of a 6-year experiment. *Ann. Bot. Fenn.* 46 (5), 397–
 769 408

770 Malloch, A.J.C (1996). Match Version2.0: A computer program to aid assignment of vegetation data
 771 to the communities and sub-communities of the National Vegetation Classification. Unit of
 772 Vegetation Science, Lancaster University.

773 Manchester, S.J., McNally, S., Treweek, J.R. Sparks, T.H., Mountford, J.O. (1999). The cost and
 774 practicality of techniques for the reversion of arable land to lowland wet grassland—an
 775 experimental study and review. *Journal of Environmental Management*, 55, 91-109,

776 McGinlay, J., Gowing, D. J. G., Budds, J. (2016). Conserving socio-ecological landscapes: An analysis of
 777 traditional and responsive management practices for floodplain meadows in England. *Environmental*
 778 *Science & Policy* 66:234–241.

779 Nakamura, F., Ishiyama, N., Sueyoshi, M., Negishi, J.N., and T. Akasaka. (2014). The Significance of
 780 Meander Restoration for the Hydrogeomorphology and Recovery of Wetland Organisms in the
 781 Kushiro River, a Lowland River in Japan *Restoration Ecology Vol. 22, No. 4, pp. 544–554* JULY 2014

782 Natural England (n.d.). Natural England online. Available at
 783 <https://www.gov.uk/government/organisations/natural-england>. Accessed 2nd September 2020.

784 Natural England (2010). Sward Enhancement: diversifying grassland by spreading species rich green
 785 hay. Natural England Technical Information Note TIN063, edition 2. Available at: [http://publications.](http://publications.naturalengland.org.uk/publication/23025)
 786 [naturalengland.org.uk/publication/23025](http://publications.naturalengland.org.uk/publication/23025) (accessed 20 May 2015).

787 Natural England, (2012). Assessing whether created or restored grassland is a BAP Priority Habitat.
 788 2012. Technical Information Note TIN110.
 789 [https://webarchive.nationalarchives.gov.uk/20150902172513/http://publications.naturalengland.or](https://webarchive.nationalarchives.gov.uk/20150902172513/http://publications.naturalengland.org.uk/publication/1649037)
 790 [g.uk/publication/1649037](https://webarchive.nationalarchives.gov.uk/20150902172513/http://publications.naturalengland.org.uk/publication/1649037)

791 Natural England, (2016). Countryside Stewardship. Baseline Evaluation of Higher Tier
 792 Agreements (BEHTA) Manual Part 2: Technical user guidance on BEHTA feature
 793 identification, condition assessment and data collection in the field.

794 Natural Resources Wales (2020). Sites of Special Scientific Interest, Responsibilities of
 795 Owners and Occupiers online. Available from [https://naturalresources.wales/guidance-and-](https://naturalresources.wales/guidance-and-advice/environmental-topics/wildlife-and-biodiversity/protected-areas-of-land-and-seas/sites-of-special-scientific-interest-responsibilities-of-owners-and-occupiers/?lang=en)
 796 [advice/environmental-topics/wildlife-and-biodiversity/protected-areas-of-land-and-seas/sites-of-](https://naturalresources.wales/guidance-and-advice/environmental-topics/wildlife-and-biodiversity/protected-areas-of-land-and-seas/sites-of-special-scientific-interest-responsibilities-of-owners-and-occupiers/?lang=en)
 797 [special-scientific-interest-responsibilities-of-owners-and-occupiers/?lang=en](https://naturalresources.wales/guidance-and-advice/environmental-topics/wildlife-and-biodiversity/protected-areas-of-land-and-seas/sites-of-special-scientific-interest-responsibilities-of-owners-and-occupiers/?lang=en). Accessed 2nd
 798 September 2020.

799 Novakovskiy, A., & Panyukov, A. (2018). Analysis of Successional Dynamics of a Sown Meadow using
 800 Ramenskii–Grime’s System of Ecological Strategies. Russian Journal of Ecology. 49. 2, 110-118. DOI
 801 10.1134/S106741361802011X.

802 Onipchenko, V.G., Dudova, K.V., Akhmetzhanova, A.A., Khomutovskiy, M.I., Dzhatdueva, T.M.,
 803 Tekeev, T.K., Elumeeva, T.G. (2020). Which plant strategies are related to dominance in alpine
 804 communities? Journal of General Biology, 81, 1, 37-46. DOI: 10.31857/S0044459620010054

805 Pierce, S., Bottinelli, A., Bassani, I., Ceriani, R.M., Cerabolini, B.E.L. (2014). How well do seed
 806 production traits correlate with leaf traits, whole-plant traits and plant ecological strategies? Plant
 807 Ecology, 215:1351–1359. DOI 10.1007/s11258-014-0392-1

808 Poptcheva K., Schwartz P., Vogel A., Kleinebecker T. & Hölzel N. (2009) Changes in wet meadow
 809 vegetation after 20 years of different management in a field experiment (north-west Germany).
 810 Agriculture, Ecosystems and Environment, 134, 108-114

811 Pywell, R.F., Bullock, J.M., Hopkins, A., Walker, K.J., Sparks, T.H., Burke, M.J.W. & Peel, S. (2002)
 812 Restoration of species-rich grassland on arable land: assessing the limiting processes using a multi-
 813 site experiment. Journal of Applied Ecology, 39, 294–309.

814 Pywell, R.F., Bullock, J.M., Roy, D.B., Warman, L., Walker, K.J., Rothery, P. (2003) Plant traits as
 815 predictors of performance in ecological restoration. Journal of Applied Ecology, 40, 65-77.

816 Ramstead, K.M., Allen, J.A. and Springer, A.E. (2012). Have wet meadow restoration projects in the
 817 Southwestern U.S. been effective in restoring geomorphology, hydrology, soils, and plant species

818 composition? Environmental Evidence,1:11.

819 <http://www.environmentalevidencejournal.org/content/1/1/11>

820 Réka, K., Balázs, D., Tóthmérész, B., Migléc, T., Tóth, K., Török, P., Lukács, K., Godó, L., Kormoczi, Z.,

821 Szilvia, R., Kelemen, A., Sonkoly, J., Kirner, A., Tischew, S., Švamberková, E., Valkó, O. (2020).

822 Establishment gaps in species-poor grasslands: artificial biodiversity hotspots to support the

823 colonization of target species. 10.1101/2020.01.23.916155.

824 Rodwell, J.S. ed. (1992). British Plant Communities. 3 *Grasslands and Montane communities*.

825 Cambridge, CUP.

826 Rothero, E., Lake, S. and Gowing, D. (eds) (2016). Floodplain Meadows – Beauty and Utility. A

827 Technical Handbook. Milton Keynes, Floodplain Meadows Partnership.

828 Rothero, E. & Tatarenko, I. (2018). Restoration success of British floodplain meadows *Aspects of*

829 *Applied Biology 139, 2018 Ecosystem and Habitat Management: Research, Policy, Practice* 1-9.

830 Ruiz-Jaen, M. C., and T. M. Aide. (2005). Restoration success: how is it being measured? *Restoration*

831 *Ecology* 13:569–577.

832 Rural Payments Agency (2020). Countryside Stewardship Online. Available at

833 <https://www.gov.uk/government/collections/countryside-stewardship>. Accessed 2nd September

834 2020.

835 Schaffers, A.P., Vasseur, M.C., Sýkora, K.V. (1998). Effects of delayed hay removal on the nutrient

836 balance of roadside plant communities. *Journal of Applied Ecology*, 35, 349-364.

837 Schaich, H., Karier, J., Konold, W. (2011). Rivers, regulation and restoration: land use history of

838 floodplains in a peri-urban landscape in Luxembourg, 1777 - 2000 *Europ. Countrys.* · 4· p. 241-264.

839 Schmidt, A., Kirner, A., Kiehl, K. Tischew, S. (2020). Seed mixture strongly affects species-richness

840 and quality of perennial flower strips on fertile soil. *Basic and Applied Ecology*. 42. 62-72.

841 10.1016/j.baae.2019.11.005.

842 Semkin BI (2009). On the relation between mean values of two measures of inclusion and measures
843 of similarity. *Biulleten' Botanicheskogo Sada-Instituta DVO RAN* 3: 91-101 (in Russian).

844 Shellswell C. H., & Squire, V. (2019). Can species-poor grassland be diversified? A case study of
845 lowland hay meadow restoration at Llanerchaeron, Ceredigion, Wales/*Conservation Evidence* 16, 1-5

846 Silvertown, J., Dodd, M. E., Gowing, D. J. G., & Mountford, J. O. (1999). Hydrologically defined
847 niches reveal a basis for species richness in plant communities. *Nature*, 400, 61–63.

848 Smith B. M., Diaz, A., Winder, L. (2017). Grassland habitat restoration: lessons learnt from long term
849 monitoring of Swanworth Quarry, UK, 1997–2014. *PeerJ* 5:e3942
850 <https://doi.org/10.7717/peerj.3942>

851 Smith, R.S., Buckingham, H., Bullard, M.J., Shiel, R.S. and Younger, A. (1996) The conservation
852 management of mesotrophic (meadow) grassland in northern England. 1. Effects of grazing, cutting
853 date and fertilizer on the vegetation of a traditionally managed sward. *Grass and Forage Science* 51,
854 278–291.

855 Smith, R. S., Shiel, R. S., Bardgett, R. D., Millward, D., Corkhill, P., Evans, P., Quirk, H., Hobbs, P. J. and
856 Kometa, S. T. (2007), Long-term change in vegetation and soil microbial communities during the
857 phased restoration of traditional meadow grassland. *Journal of Applied Ecology*, 45: 670–679, 2007.

858 Stevenson M.J., Bullock J.M. & Ward L.K. (1995). Re-creating Semi-natural Communities: Effect of
859 Sowing Rate on Establishment of Calcareous Grassland. *Restoration Ecology*, 3, 279-289.

860 Stoll-Kleemann, S. (2010). Evaluation of management effectiveness in protected areas:
861 methodologies and results. *Basic Appl. Ecol.* 11 (5), 377–382.

862 Tatarenko, I., Rothero, E., Wallace, H. (2018). Re-assemblage of plant communities: a survey of
863 floodplain meadow restoration projects in the UK. *Proceedings of the II international conference*
864 "Systematic and floristic studies in Northern Eurasia» (on the 90th anniversary of Prof. A. G.

865 Elenevsky), 5 – 8th December 2018, Moscow: Moscow Pedagogical State University. Vol. 3. P. 167-
866 171.

867 Timmermann, A., Damgaard, C., Strandberg, M.T., Svenning, J.-C. (2015). Pervasive early 21st-century
868 vegetation changes across Danish semi-natural ecosystems: more losers than winners and a shift
869 towards competitive, tall-growing species. *Journal of Applied Ecology*, 52, 21–30.

870 Török, P., Vida, E., Deák, B., Lengyel, S., Tóthmérész, B. (2011). Grassland restoration on former
871 croplands in Europe: an assessment of applicability of techniques and costs. *Biodiversity
872 Conservation*, 20:2311–2332.

873 United Nations Environment Programme (2019). New UN Decade on Ecosystem Restoration offers
874 unparalleled opportunity for job creation, food security and addressing climate change. Retrieved
875 from [https:// www.unenvironment.org/news-and-stories/press-release/new-undecade-ecosystem-
876 restoration-offers-unparalleled-opportunity](https://www.unenvironment.org/news-and-stories/press-release/new-undecade-ecosystem-restoration-offers-unparalleled-opportunity). Accessed 2nd September 2020.

877 Vinther, E. & Ha1d A. B. (2001). Restoration of an abandoned species-rich fen meadow in Denmark:
878 changes in species richness and dynamics of plant groups during 12 years. - *Nord. J. Bot.* 20:573-584.

879 Walker, K.J., Stevens, P.A., Stevens, D.P., Mountford, J.O., Manchester, S.J. & Pywell, R.F. (2004). The
880 restoration and re-creation of species rich lowland grassland on land formerly managed for intensive
881 agriculture in the UK. *Biological Conservation*, 119, 1–18.

882 Wallace, H., Prosser, M. (2016). Plant communities on floodplain meadows // *Floodplain Meadows –
883 Beauty and Utility. A Technical Handbook* pp. 45-56. Eds E Rothero, S Lake, D Gowing, Milton Keynes:
884 Floodplain Meadow Partnership.

885 Wells T.C.E., Pywell R.F. & Welch R.C. (1994). Management and restoration of species-rich grassland.
886 Report to the Ministry of Agriculture, Fisheries and Food (BD0306), Institute of Terrestrial Ecology,
887 Monks Wood.

888 Woodcock, B.A., McDonald, A.W. & Pywell, R.F. (2011). Can long-term floodplain meadow recreation
889 replicate species composition and functional characteristics of target grasslands? *Journal of Applied*
890 *Ecology*. 48(5): 1070-1078.

891 Zedler, J. B., Callaway, J. C. (2003). Adaptive restoration: a strategic approach for integrating
892 research into restoration projects. Pages 167–174. In: Rapport, DJ, Lasley WL, Rolston DE, Nielsen
893 NO, Qualset CO, Damania AB (eds) *Managing for healthy ecosystems*. Lewis Publishers, Boca Raton,
894 Florida

895

896

897

898

899

900

901

902

903

904

905

906

907

908

909

910

911

912

913

914 Appendix 1. Standard Site Assessment Form

915 Insert map of site, including location of quadrats and soil samples if collected.

Site Name	Grid Ref	County	
River	Ownership	Designation	Size
Date	Meeting with	Managed by	
Management and History			
Agri environment agreement			
Current management			
Restoration			
Technique used/Dates			
Hydrology			
Flooding regime			
Water management			
Soil-water levels (indicated by auger hole/any other data)			
Historical information			
None known			
Current site interest	Attach excel spreadsheet for botanical data		

Main grasses/herbs	
Phosphorus levels	
Soil auger photo and findings	
Site manager aspirations/objectives	
Management recommendations	

916

917

918