

# Open Research Online

---

The Open University's repository of research publications and other research outputs

## Spatial variations of primordial and recycled noble gases across Iceland

Conference or Workshop Item

How to cite:

Cogliati, Simone; Hartley, Margaret; Holland, Greg; Burgess, Raymond; Sæmundur, Ari Halldórsson; Shorttle, Oliver and Álvarez Valero, Antonio (2021). Spatial variations of primordial and recycled noble gases across Iceland. In: VMSG 2021, 6-8 Jan 2021, Online.

For guidance on citations see [FAQs](#).

© 2021 Simone Cogliati



<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Version: Accepted Manuscript

---

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

---

[oro.open.ac.uk](https://oro.open.ac.uk)

# VMSG 2021 ABSTRACT

## Spatial variations of primordial and recycled noble gases across Iceland

Simone Cogliati<sup>1\*</sup>, Margaret Hartley<sup>1</sup>, Greg Holland<sup>1</sup>, Ray Burgess<sup>1</sup>, Sæmundur A. Haldórsson<sup>2</sup>, Oliver Shorttle<sup>3</sup>, Antonio Álvarez Valero<sup>4</sup>

\* Corresponding Author: [simone.cogliati@manchester.ac.uk](mailto:simone.cogliati@manchester.ac.uk)

<sup>1</sup> SEES, University of Manchester, Oxford Road, Manchester M13 9PL, U.K.

<sup>2</sup> Institute of Earth Sciences, University of Iceland, 101 Reykjavik, Iceland.

<sup>3</sup> Department of Earth Sciences, University of Cambridge, Cambridge, CB2 3EQ, U.K.

<sup>4</sup> Department of Geology, University of Salamanca, 37008 Salamanca, Spain

Noble gas (He, Ne, Ar, Kr, Xe) compositions of mid-ocean ridge basalts (MORB) and ocean island basalts (OIB) have been widely used to investigate the geochemical structure and evolution of Earth's mantle. Many studies provide evidence for the existence of different mantle domains having distinctive chemical and noble gas signatures. Primordial mantle domains have isotopic signatures that have remained largely unmodified since the Earth's formation, while recycled mantle domains have undergone extensive modification following chemical fractionation during melt extraction and magma degassing, mantle convection, and subduction recycling. Iceland represents a perfect natural laboratory to study the inventory of primordial and recycled noble gases within the mantle thanks to its particular location above a mid-ocean ridge and a mantle plume. In this hybrid setting, melts with a deep OIB-like mantle origin and with near-primordial mantle gas signatures interact and coexist with melts formed at shallower levels that exhibit MORB-like recycled mantle chemical characteristics. On Iceland, chemical and lithological mantle heterogeneities exist on both long and short length scales, and primordial and recycled noble gases signatures can both be present even in a single sample set. We investigated the spatial relationships between Iceland's primordial and recycled mantle components by combining new high-precision noble gas (He, Ne, Ar, Kr, Xe) analyses of basaltic glass with a large existing dataset of noble gas data from subglacially erupted basalts collected across the Iceland. Here, we present noble gas data for the Western Volcanic Zone (WVZ), one of the most geologically interesting areas of Iceland. The data indicate a significant and consistent lateral variability in the noble gas signatures in relation to the distance from the plume centre. We discuss possible explanations for these variations, ways to improve our systematic understanding of mantle volatile distribution beneath Iceland, and outline future directions of this research.