

Open Research Online

The Open University's repository of research publications and other research outputs

Remarkably high abundance of presolar grains in interplanetary dust particles collected from the comet Grigg-Skjellerup dust stream

Conference or Workshop Item

How to cite:

Nguyen, A. N.; Busemann, H. and Nittler, L. R. (2007). Remarkably high abundance of presolar grains in interplanetary dust particles collected from the comet Grigg-Skjellerup dust stream. In: 38th Lunar and Planetary Science Conference, 12-16 Mar 2007, League City, Texas, USA.

For guidance on citations see [FAQs](#).

© [not recorded]

Version: [not recorded]

Link(s) to article on publisher's website:
<http://www.lpi.usra.edu/meetings/lpsc2007/pdf/2332.pdf>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

REMARKABLY HIGH ABUNDANCE OF PRESOLAR GRAINS IN INTERPLANETARY DUST PARTICLES COLLECTED FROM THE COMET GRIGG-SKJELLERUP DUST STREAM. A. N. Nguyen¹, H. Busemann^{1,2} and L. R. Nittler¹, ¹Department of Terrestrial Magnetism, Carnegie Institution of Washington, 5241 Broad Branch Road NW, Washington DC 20015, USA (nguyen@dtm.ciw.edu), ²Planetary & Space Science Research Institute, The Open University, Milton Keynes, MK7 6AA, UK.

Introduction: Interplanetary dust particles (IDPs) have been shown to preserve isotopic anomalies, particularly in nitrogen and hydrogen associated with molecular cloud organic material [1-3]. Furthermore, with the heightened capabilities of the NanoSIMS ion microprobe, submicrometer-sized presolar silicate grains have been identified in IDPs [3-6]. Recently, four IDPs collected during the 2003 Earth passage through the comet Grigg-Skjellerup dust stream were analyzed by isotopic and micro-Raman imaging [7]. All of the samples display heterogeneous distributions and anomalous “hotspots” in both D/H and ¹⁵N/¹⁴N ratios (Fig. 1). Raman spectra for these samples indicate disordered C similar to the most primitive interstellar organic material in meteorites [8]. These data support the possibility that the IDPs studied by [7] did indeed originate from comet Grigg-Skjellerup. Of the four samples, L2054 E1, a fragment from a cluster IDP, and individual IDP L2054 G4 have the highest D-enrichments, with δD up to 16,000 ‰. We thus focus on these two samples to conduct further isotopic studies.

Experimental: The IDPs L2054 E1, L2054 F1 (a fragment from the same cluster IDP as E1), L2054 G2, L2054 G3, and L2054 G4, all of which had been pressed into Au foils, were analyzed by raster ion imaging with the Carnegie NanoSIMS 50L ion microprobe. A focused (~100 nm) Cs⁺ primary ion beam was rastered over the IDPs, and negative secondary ions of the three O isotopes, ¹²C, ²⁸Si, and ¹²C¹⁴N were measured simultaneously along with secondary electrons. Integrated ion images were produced and from these, O isotopic ratio images were also calculated. Regions of interest were defined and their O isotopic ratios were normalized to the average ratio of the IDP. Grains having O isotopic compositions distinct from the isotopically normal sub-regions of the sample could thus be identified. The mineral species (i.e., oxide or silicate) was estimated from the ²⁸Si⁻ signal. For L2054 E1 and G4, the three Si isotopes and ²⁴Mg¹⁶O were analyzed in a second set of measurements. In addition, ¹⁶O and ¹⁷O were measured again to ensure that any grains previously identified as having anomalous ¹⁷O/¹⁶O ratios could easily be located and their Si isotopic ratios deduced. The Si isotopic ratios and the ¹⁷O/¹⁶O ratio were normalized to those of the average IDP.

Results: *L2054 E1 and F1:* No anomalous grains




Figure 1. Isotopic ratio images of L2054 E1. The N (NanoSIMS) and D (IMS-6f) measurements were performed previously by [7]. Grains having anomalous compositions appear distinct from the rest of the IDP. Presolar silicate grains are circled in the O ratio images.

were identified in F1 (~15 μm²). However, four grains having anomalous O isotopic compositions were identified in L2054 E1 (~140 μm², Fig. 1). All four grains have ²⁸Si/¹⁶O⁻ ratios similar to the average ratio of the IDP fragment, which is composed mostly of silicate minerals. In addition, all grains display ²⁴Mg¹⁶O⁻, making it likely that these grains are Mg-rich circumstellar silicates. One of these grains is enriched in ²⁹Si and ³⁰Si ($\delta^{29}\text{Si} = 149 \pm 25 \text{ ‰}$; $\delta^{30}\text{Si} = 133 \pm 30 \text{ ‰}$). *L2054 G4:* This ~40 μm² IDP was found to contain 7 grains with anomalous O isotopic compositions. Six of these grains were unambiguously identified during the second set of measurements (though also visible in the first) and thus the ¹⁸O/¹⁶O ratios were not well determined. All of the grains appear to be silicates with some Mg. The Si isotopic compositions for these grains fall in the range observed for mainstream SiC grains and for other presolar silicate grains [9, 10]. In addition, one grain has depletions in ²⁹Si and ³⁰Si ($\delta^{29}\text{Si} = -159 \pm 25 \text{ ‰}$, $\delta^{30}\text{Si} = -74 \pm 39 \text{ ‰}$) similar to those seen in SiC X grains that originate from Type II supernova. The ²⁸Si/¹⁶O⁻ ratio for this grain is greater than the average for this IDP. An attempt to obtain the C isotopic composition for this grain was not successful as the grain had been sputtered away. *L2054 G2 and G3:* No anomalous grains were found in these

two large (total analyzed area $\sim 1800 \mu\text{m}^2$) individual IDPs, which also show less extreme H and N isotopic variability than the other samples.

Discussion: Fig. 2 shows the O isotopic ratios of the presolar silicates from L2054 E1 and G4. The 5 grains for which we have $^{18}\text{O}/^{16}\text{O}$ data fall into the presolar oxide Group 1 believed to originate from low-mass red giant and asymptotic giant branch stars [11]. The Si isotopic ratios for all grains are plotted in Fig. 3 along with the SiC mainstream correlation line. This correlation line mainly reflects the initial composition of the parent stars.

The inferred abundance of presolar silicate grains in the cluster IDP of which E1 and F1 are fragments is 0.35 %. The abundance in IDP L2054 G4 is 1.5 %. In contrast, the estimated average abundance of presolar silicates in isotopically primitive IDPs is ~ 375 ppm [3]. Clearly L2054 E1 and G4 are some of the most primitive extraterrestrial materials studied. This is exemplified not only by the exceptionally high abundances of presolar grains, but also by the large D and ^{15}N anomalies and extremely disordered Raman signatures seen previously [7].

That two of four studied IDPs from the L2054 April 2003 collection, one cluster particle and one individual one, are so strikingly different from almost all previously studied IDPs is statistically unlikely. This




Figure 2. Oxygen isotopic ratios for presolar silicates from IDPs L2054 E1 and G4. The data points on the left represent the silicates from G4 for which $^{18}\text{O}^-$ was not directly measured. Error bars are 1σ .

strengthens the possibility of a link to comet Grigg-Skjellerup. Unusual noble gases reported for some IDPs from the same collection might also support such a link [12]. However, it is interesting to note that the IDPs studied here show much higher abundances of presolar organic matter and presolar grains than has been found in samples from comet Wild-2 [13, 14]. If the IDPs are indeed from Grigg-Skjellerup, then this difference might reflect heterogeneity within or between comets or biases introduced by the different sampling methods.

References: [1] Aléon J. et al. (2001) *Geochim. Cosmochim. Acta*, 67, 3773-3787. [2] Busemann H. et al. (2006) *Science*, 312, 727-730. [3] Floss C. et al. (2006) *Geochim. Cosmochim. Acta*, 70, 2371-2399. [4] Messenger S. et al. (2003) *Science*, 300, 105-108. [5] Floss C. and Stadermann F. J. (2004) *LPS XXXV*, Abstract #1281. [6] Messenger S. et al. (2005) *Science*, 309, 737-741. [7] Nittler L. R. et al. (2006) *LPS XXXVII*, Abstract #2301. [8] Busemann et al. (2007) *Meteoritics & Planet. Sci.*, submitted. [9] Mostefaoui S. and Hoppe P. (2004) *ApJ*, 613, L149-L152. [10] Nguyen A. N. et al. (2007) *ApJ*, submitted. [11] Nittler L. R. et al. (1997) *ApJ*, 483, 475-495. [12] Palma R. L. et al. (2005) *Meteoritics & Planet. Sci. (Supp.)*, 40, Abstract #5012. [13] Sandford S. A. et al. (2006) *Science*, 314, 1720-1724. [14] McKeegan K. D. (2006) *Science*, 314, 1724-1728.




Figure 3. Silicon isotopic ratios for presolar silicates from IDPs L2054 E1 and G4. The triangular data point is a possible SiC X grain from G4. Error bars are 1σ . Also shown is the SiC mainstream correlation line (dash-dotted line).