

Open Research Online

The Open University's repository of research publications and other research outputs

Noble gases from the interstellar medium trapped on the MIR space station and analyzed by in vacuo etching

Conference or Workshop Item

How to cite:

Busemann, H.; Bühler, F.; Agafonov, Y. N.; Baur, H.; Bochsler, P.; Eismont, N. A.; Heber, V. S.; Wieler, R. and Zastenker, G. N. (2004). Noble gases from the interstellar medium trapped on the MIR space station and analyzed by in vacuo etching. In: 67th Annual Meeting of the Meteoritical Society, 4-6 Aug 2004, Rio de Janeiro, Brazil.

For guidance on citations see [FAQs](#).

© [not recorded]

Version: [not recorded]

Link(s) to article on publisher's website:

<http://www.lpi.usra.edu/meetings/metsoc2004/pdf/5170.pdf>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

NOBLE GASES FROM THE INTERSTELLAR MEDIUM TRAPPED ON THE MIR SPACE STATION AND ANALYZED BY IN VACUO ETCHING.

H. Busemann^{1*}, F. Bühler¹, Y.N. Agafonov², H. Baur³, P. Bochsler¹, N.A. Eismont², V.S. Heber^{1,3**}, R. Wieler³ and G.N. Zastenker². ¹Physikalisches Institut, University of Bern, Switzerland. ²IKI, Russian Academy of Sciences, Moscow, Russia. ³Earth Sciences, ETH Zurich, Switzerland. Present addresses: *E-mail: busemann@dtm.ciw.edu. DTM, Carnegie Institution, Washington DC, USA. **Earth Sciences, Open University, Milton Keynes, UK.

Introduction: The composition of the present interstellar medium (ISM) provides an important benchmark in cosmochemistry. It serves as a reference for galactic chemical evolution (GCE) models, solar mixing predictions and provides information for understanding Big Bang nucleosynthesis. The present-day ISM ³He abundance allows, combined with the protosolar ³He, deduced from the Jovian atmosphere or meteorites [1,2], tracing the GCE over the past 4.56 Ga. ${}^3\text{He}/{}^4\text{He} = (2.5 \pm 0.6) \times 10^{-4}$ has been determined for the local ISM [3]. However, the uncertainty is too large to better constrain GCE models and - in combination with the present-day solar wind value - the protosolar D/H [4].

Experiment: The COLLISA experiment [*Collection of Inter-Stellar Atoms*, 5,6] sampled interstellar gas in Cu-Be foils covered with BeO and exposed to the flux of neutrals from the ISM on board the MIR space station. Stepwise heating extraction allowed the detection of interstellar ⁴He [6] and yielded $({}^3\text{He}/{}^4\text{He})_{\text{ISM}} = (1.7 \pm 0.8) \times 10^{-4}$ [7], in agreement with the value for pickup ions observed with SWICS/Ulysses [3].

Further foils are currently analyzed by closed system stepwise etching at ETH Zurich [8]. This technique allows to efficiently separate implanted interstellar He and terrestrial tritogenic ³He, probably residing in the Cu-Be substrate, which had to be taken into account for the determination of interstellar ³He during stepwise heating [7].

Results: Offline tests suggest that HF acid vapor efficiently and uniformly etches BeO. The system blank (in 10^{-14} cm³ STP, ³He ~3, ⁴He ~300, ²⁰Ne ~90) is now sufficiently low to measure the exposed foils. Two unexposed foils (31 and 50 cm²) were etched online and yielded no significantly increased values relative to these blanks, implying that the tritogenic ³He (0.5-1 10^{-14} cm³/cm² foil) indeed resides in deeper foil layers that are not affected by superficial etching. The analysis of a foil artificially irradiated with ³He and ²⁰Ne at energies comparable to those of the ISM neutrals (25 eV/amu) showed that our protocol (10 steps 1-30 min, HF vapor at 20 °C) releases all trapped noble gases. Results of the ongoing etching experiment on foils exposed to the ISM (including a witness foil doped with terrestrial ³He) will be presented. The expected concentrations of interstellar gas [3,6,7] in 50 cm² of exposed foil are (in 10^{-14} cm³ STP) ³He ~25, ⁴He ~200000, ²⁰Ne ~375).

Acknowledgement: This work was supported by the Swiss NSF. Technical support by U. Menet is gratefully acknowledged.

References: [1] Mahaffy P. R. et al. 2000. *JGR* 105:15061-15071. [2] Busemann H. et al. 2001. *Lunar Planet. Sci. Conf. XXXII*:#1598. [3] Gloeckler G. and Geiss J. 2001. *AIP Conf. Proc.* 598:281-289. [4] Geiss J. and Gloeckler G. 2002. *Space Sci. Rev.* 105:3-18. [5] Zastenker G. N. et al. 2002. *Cosmic Research* 40:347-357. [6] Bühler F. et al. 2000. *Astrophys. Space Sci.* 274:19-24. [7] Salerno E. et al. 2003. *Astrophys. J.* 585:840-849. [8] Busemann H. 2000. *Meteorit. Planet. Sci.* 35:949-973.