

Open Research Online

The Open University's repository of research publications and other research outputs

Oxygen isotope evidence for the origin of HEDs and Angrites

Conference or Workshop Item

How to cite:

Greenwood, R.G.; Franchi, I.A. and Jambon, A. (2003). Oxygen isotope evidence for the origin of HEDs and Angrites. In: 66th Annual Meteoritical Society Meeting, 28 Jul - 1 Aug 2003, Munster, Germany.

For guidance on citations see [FAQs](#).

© [\[not recorded\]](#)

Version: [\[not recorded\]](#)

Link(s) to article on publisher's website:

<http://www.lpi.usra.edu/meetings/metsoc2003/pdf/5178.pdf>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's [data policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk


OXYGEN ISOTOPE EVIDENCE FOR THE ORIGIN OF HEDs AND ANGRITES. R C Greenwood¹, I A Franchi¹, A Jambon² ¹PSSRI, Open University, Milton Keynes, MK7 6AA, U.K. ²Laboratoire MAGIE, Université Pierre et Marie Curie, CNRS UMR 7047 case 110, 4 place Jussieu, 75252 Paris Cedex 05 France.

Introduction: HEDs and angrites represent distinct suites of mafic igneous rocks formed early in the history of the Solar System [1]. Despite their mineralogical and chemical differences both groups have similar O-isotope compositions [2]. To examine the origin of HEDs and angrites we have undertaken a high-precision study of their O-isotope systematics.

Experimental Techniques: Oxygen isotope analyses were undertaken by infrared laser-assisted fluorination [3]. Samples were fused prior to fluorination. O₂ was analyzed using a Micromass Prism III dual inlet mass spectrometer. Precision is $\pm 0.04\%$ for $\delta^{17}\text{O}$, $\pm 0.08\%$ for $\delta^{18}\text{O}$ and $\pm 0.025\%$ for $\Delta^{17}\text{O}$ [3]. Meteorites studied: *Diogenites*: Bilanga (4), Johnstown (4), Shalka (2), Tatahouine (2), *Eucrites*: Pasamonte (5), Juvinas (2), Padvarnika (2), Sioux County (2), Stannern (2), Moore County (2), *Howardites*: Kapoeta (2), Pavlovka (2), Molteno (2), *Angrites*: Angra dos Ries (4), LEW 86010* (2), D'Orbigny* (2), NWA 1296* (2). (all falls except as indicated by an asterisk, number of replicate analyses in brackets).

Results: Angrites and HEDs plot as distinct groups (Fig 1) and define single mass fractionation lines ($\Delta^{17}\text{O} = -0.056 \pm 0.007$ for angrites and -0.220 ± 0.012 for HEDs). Previous studies have indicated that Angra dos Reis has an O-isotope composition indistinguishable from HEDs [2]. Using two distinct sub-samples we have found it to have a $\Delta^{17}\text{O}$ value of -0.065 ± 0.009 and plots in the angrite field (Fig. 1). Eucrites and diogenites plot at either end of the HED array (Fig. 1), while howardites occupy a more central position. The large error bars for $\delta^{18}\text{O}$ reflect the inhomogeneous, coarse-grained nature of HEDs, with diogenites having grain sizes in excess of 5cm [1].

Discussion: Experimental work indicates that either an angritic or eucritic melt can be produced from a chondritic parent simply by partial melting under differing conditions of oxygen fugacity [1]. Thus a single asteroid might be capable of producing both the HEDs and angrites. However, O-isotope data indicates that this is highly unlikely and that both suites must come from distinct parent bodies. Furthermore, as both fall on well-defined mass fractionation lines only a single parent body per group is required. Finally, in contrast to recent suggestions [4], O-isotope data presented here clearly demonstrates that Angra dos Ries is an angrite.



References: [1] Mittlefehldt D. W. et al (1998) in *Planetary Materials*, Papike J.J.(ed). [2] Clayton R. N. and Mayeda T.K. (1996) *GCA*, 60, 1999-2017. [3] Miller M.F. et al. (1999) *Rapid Commun. Mass Spectrom.* 13, 1211-1217. [4] Mittlefehldt et al. (2002) *Meteorit. Planet. Sci.*, 37, 345-369.