

Open Research Online

The Open University's repository of research publications and other research outputs

Dar Al Gani 896: A unique picritic achondrite

Conference or Workshop Item

How to cite:

Folco, L.; Bland, P.A.; D'Orazio, M.; Franchi, I.A. and Rocchi, S. (2002). Dar Al Gani 896: A unique picritic achondrite. In: 65th Annual Meteoritical Society Meeting, 21-26 Jul 2002, Los Angeles, California, USA.

For guidance on citations see [FAQs](#).

© [\[not recorded\]](#)

Version: [\[not recorded\]](#)

Link(s) to article on publisher's website:

<http://www.lpi.usra.edu/meetings/metsoc2002/pdf/5176.pdf>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

DAR AL GANI 896: A UNIQUE PICRITIC ACHONDRITE.

L. Folco¹, P. A. Bland², M. D'Orazio³, I. A. Franchi², and S. Rocchi³, ¹Museo Nazionale Antartide, Via Laterina 8, 53100 Siena, Italy, ²Planetary and Space Sciences Research Institute, Open University, Milton Keynes, MK7 6AA, U.K., ³Dipartimento di Scienze della Terra, Università di Pisa, Via S. Maria 53, 56126 Pisa, Italy, (folco@unisi.it)

DaG 896 is the provisional name assigned to a dark, 50% fusion crusted, stony fragment of 22.6 g found in the Dar al Gani region of the Libyan Sahara (~27°08'N – 16°05'E) in November 2000.

Mineralogy: DaG 896 is hypocrystalline with porphyritic texture (porphyritic index ~70). Abundant fine-grained (avg. grain-size 100 µm), skeletal forsterite crystals (avg. Fo₈₂) are set in a ground-mass consisting mainly of rhyolitic glass plus quench microlites of pigeonite (En₅₄ Wo₉). Minor mineral constituents include enstatite (En₈₃ Wo₂), augite (En₃₇ Wo₃₀) and traces of chromite, troilite, Fe-Ni metal and phosphates. Veinlets up to 15 µm thick of secondary carbonates are most likely due to terrestrial weathering. Mineral mode (vol. %) is forsterite 69, glass 17, clinopyroxene (pigeonite >> augite) 8, enstatite 4, carbonates 1, others 1. Strong undulose extinction and planar fracturing in olivine and mm-thick interconnecting veinlets indicate moderate shock S4, after [1]. Fe/Mn a.p.f.u. ratios for olivine and pyroxenes (35 and 18, respectively) are chondritic [2]. No achondrite with the above texture and mineral composition is documented in literature [e.g., 2]. Similar textures and mineral compositions are in turn observed in some igneous inclusions in ordinary chondrites [e.g. 3].

Bulk chemistry: XRF and ICP-MS major and trace element concentrations indicate an high-Mg (mg# 79.2) picritic *IUGS* composition (SiO₂ = 47.62, Na₂O = 0.16, K₂O = 0.24, MgO = 30.3 wt %) with a chondritic [Mn/Mg] = 7.5 *10⁻³ ratio. The overall composition is similar to H-chondrites [4] (e.g., Mg/Si = 0.83, Al/Si = 0.073, Ti/Si = 0.006) except for a strong depletion in siderophile and chalcophile elements (e.g., Fe/Si = 0.50, Fe/Ni = 254 and S/Si = 0.009). A close major element compositional match is found with the harzburgitic igneous inclusion in the Y-794046 H5 chondrite [5], [6] and [7] and the komatiite-like lithic fragment in the Eva H4 chondrite [3]. Anomalously high concentrations of Cs, Rb, Sr and Pb suggest some terrestrial contamination.

Oxygen isotopes: Mean data from two replicates are δ¹⁷O = 2.55, δ¹⁸O = 3.50 and Δ¹⁷O = 0.726 ± 0.007 ‰. The Δ¹⁷O value is typical for equilibrated H-chondrites [8], although the δ¹⁸O ratio is slightly offset to the light side for this grouping.

Conclusions: i) DaG 896 is a unique picritic achondrite; some igneous inclusions in ordinary chondrites are the closest existing analogs. ii) DaG 896 formed through rapid cooling of a high-temperature melt likely representing the silicate fraction of H-chondritic source material, which underwent severe heating, melting and metal plus sulfide loss.

References: [1] Stöffler D. et al. (1991) *GCA*, 55, 3845-3867. [2] Papike J. J., ed. (1998) *Rev. Mineral.*, 36. [3] Ruzicka et al. (1998) *GCA* 62, 1419-1442. [4] Jarosewich E. (1990) *Meteoritics* 25, 323-337. [5] Nakamura N. et al. (1994) *Proc. NIPR Symp. Antarct. Meteorites*, 7, 125-143. [6] Sack R. O. et al. (1994) *JGR*, 99, E12, 26029-26044. [7] Mittlefehldt D. W. et al. (1995) *Proc. NIPR Symp. Antarct. Meteorites*, 8, 251-271. [8] Clayton R. N. et al. (1991) *GCA*, 55, 2317-2337.