

# Open Research Online

---

The Open University's repository of research publications and other research outputs

## Carbon and oxygen isotopes in CO<sub>3</sub> chondrites

### Conference or Workshop Item

How to cite:

Greenwood, R.G.; Franchi, I.A.; Gibson, J.M. and Pillinger, C.T. (2002). Carbon and oxygen isotopes in CO<sub>3</sub> chondrites. In: 65th Annual Meteoritical Society Meeting, 21-26 Jul 2002, Los Angeles, California, USA.

For guidance on citations see [FAQs](#).

© [\[not recorded\]](#)

Version: [\[not recorded\]](#)

Link(s) to article on publisher's website:

<http://www.lpi.usra.edu/meetings/metsoc2002/pdf/5167.pdf>

---

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

---

[oro.open.ac.uk](http://oro.open.ac.uk)

**CARBON AND OXYGEN ISOTOPES IN CO<sub>3</sub> CHONDRITES.**

R. C. Greenwood, I. A. Franchi, J. Gibson and C. T. Pillinger, PSSRI, Open University, Walton Hall, Milton Keynes, MK7 6AA U.K. (R.C.Greenwood@open.ac.uk)

**Introduction:** CO<sub>3</sub> chondrites form a metamorphic series and have been divided into subtypes ranging from 3.0 to 3.7 [1], [2], [3]. To examine the processes and conditions prevailing during metamorphism we have undertaken a detailed investigation of the whole rock oxygen and carbon isotope systematics of CO<sub>3</sub> chondrites.

**Experimental Techniques:** Oxygen isotope analyses were made using an infrared laser fluorination system [4]. All analyses were obtained on powders that were fluorinated using BrF<sub>5</sub> and then heated progressively for periods of up to 50 minutes. O<sub>2</sub> was analysed using a Micromass Prism III dual inlet mass spectrometer. Analytical precision is approximately ±0.04‰ for δ<sup>17</sup>O, ±0.08‰ for δ<sup>18</sup>O and ±0.025‰ Δ<sup>17</sup>O. Carbon isotopes were determined using a Geo 20-20 mass spectrometer with an ANCA elemental analyser preparation system. Analytical precision is ±0.09‰ δ<sup>13</sup>C.

**Results:** The following CO<sub>3</sub> chondrites have been analysed: ALH77307(3.0), Colony(3.0), Kainsaz(3.1\*), Felix(3.2\*), Ornan(3.3\*), ALH82101(3.3), Lance(3.4\*), ALH77003(3.5) Warrenton(3.6\*), Isna(3.7) (figures in brackets are the metamorphic subtypes of [2], asterisks indicate a fall)

*Oxygen isotopes:* With the exception of Colony(3.0) and ALH77307(3.0), samples fall within an extremely restricted area of the oxygen three-isotope diagram, variation being less than that reported by [5]. If finds are excluded, and with the possible exception of Warrenton(3.6), there is a positive correlation between Δ<sup>17</sup>O and metamorphic subtype. Analyses of different sub-samples of Lance(3.4) demonstrate small, but significant, levels of sample heterogeneity (up to approximately 0.2‰ for δ<sup>17</sup>O and 0.5‰ for δ<sup>18</sup>O).

*Carbon isotopes:* A distinct negative correlation is displayed when δ<sup>13</sup>C is plotted against metamorphic grade, the relationship being particularly well developed if finds are excluded. In addition, whole rock carbon abundance declines with increasing grade being 0.8% in ALH77307(3.0) and 0.3% in Isna(3.7).

**Discussion:** The suggestion that there is a correlation between whole rock oxygen isotope compositions and metamorphic subtype [3], [5] is supported by the results of this study, contrary to our initial findings [6]. Our results are consistent with the involvement of an aqueous fluid phase during metamorphism [3]. The presence of phyllosilicates within the matrices of a number of CO<sub>3</sub> chondrites [7] lends further support to this possibility. Whole rock C isotopes show a clear negative correlation with metamorphic grade, as does C abundance. In view of the evidence that alteration took place under relatively oxidising conditions [8] whole rock C isotope systematics are consistent with high partial pressures of CO<sub>2</sub> in the fluid phase during metamorphism. The presence of carbonate in Warrenton(3.6), as detected in step combustion studies [9], provides additional evidence of high CO<sub>2</sub> levels during metamorphic alteration on the CO parent body.

**References:** [1] McSween H.Y (1977) *Geochim. Cosmochim. Acta*, 41, 477-491. [2] Scott E.R.D. and Jones R.H. (1990) *Geochim. Cosmochim. Acta*, 54, 2485-2502. [3] Rubin A.E. (1998) *Meteorit. Planet. Sci.*, 33, 385-391. [4] Miller et al. (1999) *Rapid Commun. Mass Spectrom.* 13, 1211-1217. [5] Clayton R.N. and Mayeda T.K. (1999) *Geochim. Cosmochim. Acta*, 63, 2089-2104. [6] Greenwood et al (2002) *LPS XXXIII Abstract 1609*. [7] Brearley A.J. and Jones R.H. (1998) *Reviews in Mineralogy* 36(3), 1-398. [8] Keller L.P. and Buseck P.R. *Geochim. Cosmochim. Acta*, 54, 1155-1163. [9] J. Newton (1994) *Unpublished Ph.D.Thesis*, Open Uni.