

# Open Research Online

---

The Open University's repository of research publications and other research outputs

## Chemical fractionation and alteration trends between five nakhlites

Conference or Workshop Item

How to cite:

Bridges, J.; Jeffries, T.E. and Grady, M.M. (2003). Chemical fractionation and alteration trends between five nakhlites. In: 66th Annual Meeting of the Meteoritical Society, 28 Jul - 1 Aug 2003, Muster, Germany.

For guidance on citations see [FAQs](#).

© [\[not recorded\]](#)

Version: [\[not recorded\]](#)

Link(s) to article on publisher's website:

<http://www.lpi.usra.edu/meetings/metsoc2003/pdf/5228.pdf>

---

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

---

[oro.open.ac.uk](http://oro.open.ac.uk)

**CHEMICAL FRACTIONATION AND ALTERATION TRENDS BETWEEN FIVE NAKHLITES.** J. C. Bridges, T. E. Jeffries and M. M. Grady, Dept. of Mineralogy, Natural History Museum, London SW7 5BD, UK, j.bridges@nhm.ac.uk.

**Introduction:** We combine new petrographic and geochemical (solution ICPMS) data on 5 nakhlites in order to show igneous fractionation and alteration trends between the samples.

**Trace Elements:** Nakhla (BM1913, 26) and Y00593 have nearly identical relative abundances of the REE e.g. La/Lu = 3.5, 4.0 but Nakhla has 4.5 – 1.3 x CI REE with Y00593 having higher abundances 6.9 – 1.7 x CI. The abundances are consistent with Rayleigh fractionation of a LREE-enriched melt - composition from [1] - and crystallisation of an ol-cpx adcumulate from it at F (fraction of melt remaining) = 0.6. Addition of ~5% trapped melt to this produces REE abundances similar to those of Y00593. Nakhla has trapped slightly lower proportions of melt, or its melt experienced a lower degree of crystal fractionation.




Fig. 1: CI normalised trace elements in Y00593 – see also [2] - and Nakhla. These can be modelled as addition of trapped melt to an ol-cpx adcumulate derived from a LREE-enriched parental melt.

**Siderite-bearing veins in NWA998 and Y00593:** ASEM analyses of veins within the olivine of these 2 nakhlites show that they contain a mixture of Mn-bearing siderite and Fe-rich clay. The chemical and textural similarities between these veins and those of the other nakhlites suggest they have a related origin on Mars [3,4]. The volume of these veins is smaller than those of the 3 other nakhlites we have studied [3]. Using the model of [3] the Fe-Mn dominated composition of the siderite/clay mixtures suggests that they formed at a similar, late stage of brine evaporation as the veins in Nakhla.

**References:** [1] Wadhwa M. and Crozaz G. (2003) *LPS XXXIV*, #2075. [2] Nakamura N. et al. (2002) *Antarct. Meteorites XXVII*, 112. [3] Bridges J. C. and Grady M. M. (2000) *EPSL*, 176, 267–279. [4] Gooding J. L. et al. (1991) *Meteoritics*, 26, 135-143.