

# Open Research Online

---

The Open University's repository of research publications and other research outputs

## A counterexample to a continued fraction conjecture

### Journal Item

How to cite:

Short, Ian (2006). A counterexample to a continued fraction conjecture. Proceedings of the Edinburgh Mathematical Society. Series II, 49(3) pp. 735–737.

For guidance on citations see [FAQs](#).

© 2006 Edinburgh Mathematical Society

Version: Accepted Manuscript

Link(s) to article on publisher's website:  
<http://dx.doi.org/doi:10.1017/S0013091505000581>

---

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

---

[oro.open.ac.uk](http://oro.open.ac.uk)

# A COUNTEREXAMPLE TO A CONTINUED FRACTION CONJECTURE

IAN SHORT

ABSTRACT. It is known that if  $a \in \mathbb{C} \setminus (-\infty, -1/4]$  and  $a_n \rightarrow a$  as  $n \rightarrow \infty$ , then the infinite continued fraction with coefficients  $a_1, a_2, \dots$  converges. A conjecture has been recorded by L. Jacobsen et al, taken from the unorganized portions of Ramanujan's notebooks, that if  $a \in (-\infty, -1/4)$  and  $a_n \rightarrow a$  as  $n \rightarrow \infty$ , then the continued fraction diverges. Counterexamples to this conjecture for each value of  $a$  in  $(-\infty, -1/4)$  are provided. Such counterexamples have already been constructed by Glutsyuk, but the examples given here are significantly shorter and simpler.

## 1. INTRODUCTION

For each  $n \in \mathbb{N}$ , let  $a_n$  be a non-zero complex number and let  $t_n$  be the Möbius transformation  $t_n(z) = a_n/(1+z)$ ; then the continued fraction

$$(1.1) \quad \mathbf{K}(a_n | 1) = \frac{a_1}{1 + \frac{a_2}{1 + \frac{a_3}{1 + \dots}}}$$

is considered to converge if the sequence with  $n$ th term equal to the  $n$ -fold composition  $t_1 \cdots t_n(0)$  converges within the extended complex plane  $\mathbb{C}_\infty$ . We identify the continued fraction (1.1) with the sequence  $t_1, t_2, \dots$  of Möbius transformations. A problem derived from the private notebooks of Ramanujan is posed in [3, page 38], which asks whether, for a given complex number  $a \neq -1/4$  and a sequence  $a_1, a_2, \dots$  that converges to  $a$ , the continued fraction  $\mathbf{K}(a_n | 1)$  diverges if and only if  $a \in (-\infty, -1/4)$ . In this paper it is demonstrated that  $\mathbf{K}(a_n | 1)$  may or may not converge if  $a \in (-\infty, -1/4)$ , thereby proving the conjecture to be false. Glutsyuk has already provided such examples in [4], but our methods are significantly shorter and simpler. Our conclusions are summarized in a theorem, whose proof is postponed until §3.

---

2000 *Mathematics Subject Classification.* Primary 40A15, 30B70.

*Key words and phrases.* Möbius transformations, continued fractions, iteration, dynamics.

This work was supported by Science Foundation Ireland grant 05/RFP/MAT0003.

**Theorem 1.1.** *If  $a \in (-\infty, -1/4)$  then there are sequences  $a_n$  of real numbers that converge to  $a$  for which  $\mathbf{K}(a_n|1)$  converges and there are sequences  $a_n$  of real numbers that converge to  $a$  for which  $\mathbf{K}(a_n|1)$  diverges.*

## 2. ITERATION OF A SINGLE MÖBIUS TRANSFORMATION

To understand the dynamics of the sequence  $t_1 \cdots t_n$ , where  $t_n(z) = a_n/(1+z)$  and  $a_n \rightarrow a$  as  $n \rightarrow \infty$ , one must first understand the dynamics of the sequence formed through iterating the Möbius map  $t(z) = a/(1+z)$ . The theory of iteration of a single Möbius transformation is well known (see, for example, [1] or [5]) and it is independent of continued fractions. We elaborate briefly on this theory.

The conjugacy type of a given Möbius transformation  $f(z) = (Az + B)/(Cz + D)$  may be determined from the conjugation invariant quantity  $T(f) = (A + D)^2/(AD - BC)$ : if  $T(f) \in [0, 4)$  then  $f$  is *elliptic*; if  $T(f) = 4$  then  $f$  is *parabolic*; otherwise  $f$  is *loxodromic*. Therefore  $t$  is elliptic if  $a \in (-\infty, -1/4)$ ; parabolic if  $a = -1/4$ ; and otherwise loxodromic.

If  $t$  is loxodromic and  $a_n \rightarrow a$  as  $n \rightarrow \infty$ , it follows from the general theory (see [2] or [6]) that  $\mathbf{K}(a_n|1)$  converges. If  $t$  is parabolic and  $a_n \rightarrow a$  as  $n \rightarrow \infty$ ,  $\mathbf{K}(a_n|1)$  may converge or it may diverge, and it is easy to construct examples of both circumstances. This leaves the situation of Theorem 1.1, when  $t$  is elliptic. Elliptic maps are by definition conjugate to Möbius maps of the form  $z \mapsto e^{i\theta}z$ , where  $\theta \in (0, 2\pi)$ , hence  $t^n(0)$  diverges (since 0 is not a fixed point of  $t$ ), that is,  $\mathbf{K}(a|1)$  diverges. Thus, for one part of Theorem 1.1 we may choose  $a_n$  to be the constant sequence  $a, a, \dots$ . The other part of Theorem 1.1 is proved in §3.

## 3. PROOF OF THEOREM 1.1

We need a preliminary lemma.

**Lemma 3.1.** *The subset of  $(-\infty, -1/4)$  consisting of those numbers  $a \in (-\infty, -1/4)$  for which  $t(z) = a/(1+z)$  is a map of finite order is a dense subset of  $(-\infty, -1/4)$ .*

*Proof.* Let  $t$  be conjugate to  $g(z) = e^{i\theta}z$ ,  $\theta \in (0, 2\pi)$ ; then

$$(3.1) \quad -1/a = T(t) = T(g) = 4 \cos^2 \frac{1}{2}\theta.$$

The maps  $t$  and  $g$  are of finite order if and only if  $\theta$  is a rational multiple of  $\pi$ , and rational multiples of  $\pi$  are dense in  $(0, 2\pi)$ . The result is assured by continuity of the correspondence (3.1).  $\square$

*Proof of Theorem 1.1.* We construct a sequence  $a_n$  that converges to  $a \in (-\infty, -1/4)$  for which  $\mathbf{K}(a_n|1)$  converges. By Lemma 3.1, we may choose a sequence  $\alpha_1, \alpha_2, \dots$  in  $(-\infty, -1/4)$  that converges to  $a$  for which each map  $s_n(z) = \alpha_n/(1+z)$  is of finite order. Let  $\epsilon_1, \epsilon_2, \dots$  be a sequence in  $(0, 1)$  that converges to 0 for which  $\sum \epsilon_n$  diverges. Define  $t_n(z) = (1 - \epsilon_n)\alpha_n/(1+z)$ , for  $n = 1, 2, \dots$ . One may easily verify that

$$(3.2) \quad t_n s_n^{-2} t_n(z) = z + \epsilon_n.$$

Since  $s_n$  is of finite order, the two equal quantities in equation (3.2) are also equal to the  $m$ -fold composition  $t_n s_n \cdots s_n t_n(z)$ , where  $m = \text{order}(s_n)$ .

For each  $n$ , choose an integer  $N_n$  such that  $N_n \epsilon_n$  is greater than the maximum element from the finite set

$$(3.3) \quad \{|t_{n+1} s_{n+1}^q(0)| : 0 \leq q \leq \text{order}(s_{n+1}) - 2, t_{n+1} s_{n+1}^q(0) \neq \infty\}.$$

Let  $\phi_n$  represent the string of maps  $t_n, s_n, \dots, s_n, t_n$ , in which  $s_n$  occurs  $\text{order}(s_n) - 2$  times. The continued fraction corresponding to the sequence of Möbius maps

$$(3.4) \quad \phi_1, \dots, \phi_1, \phi_2, \dots, \phi_2, \dots$$

is the example we require, where the string  $\phi_n$  occurs in the continued fraction  $N_n$  times. To see that (3.4) provides an example of the required form, notice that the coefficients  $a_n$  arise from the maps  $s_n$  or  $t_n$ , thus certainly  $a_n \rightarrow a$  as  $n \rightarrow \infty$ . It remains to demonstrate that the continued fraction converges (to  $\infty$ ). This is true as

$$(t_1 s_1^{-2} t_1)^{N_1} \cdots (t_n s_n^{-2} t_n)^{N_n}(z) = z + \sum_{i=1}^n N_i \epsilon_i,$$

by equation (3.2), hence

$$\begin{aligned} & (t_1 s_1^{-2} t_1)^{N_1} \cdots (t_n s_n^{-2} t_n)^{N_n} (t_{n+1} s_{n+1}^{-2} t_{n+1})^p t_{n+1} s_{n+1}^q(0) \\ &= t_{n+1} s_{n+1}^q(0) + p \epsilon_{n+1} + \sum_{i=1}^n N_i \epsilon_i \\ &> \sum_{i=1}^{n-1} N_i \epsilon_i, \end{aligned}$$

by (3.3), where  $0 \leq p < N_{n+1}$  and  $0 \leq q \leq \text{order}(s_{n+1}) - 2$ . Therefore, the continued fraction converges to  $\infty$ .  $\square$

## REFERENCES

- [1] A. F. Beardon. *The Geometry of Discrete Groups*. Springer-Verlag, 1983.
- [2] A. F. Beardon. Continued fractions, discrete groups and complex dynamics. *Computational Methods and Function Theory*, 1:535–594, 2001.

- [3] L. Jacobsen R. L. Lamphere G. E. Andrews, B. C. Berndt. The continued fractions found in the unorganized portions of Ramanujan's notebooks - memoirs of the AMS. *Mem. Amer. Math. Soc.*, 99(477), 1992.
- [4] A. A. Glutsyuk. On convergence of generalized continued fractions and Ramanujan's conjecture. *C. R. Acad. Sci. Paris*, 341(7):427–432, 2005.
- [5] G. Jones and D. Singerman. *Complex Functions: an algebraic and geometric viewpoint*. Cambridge University Press, 1987.
- [6] M. Mandell and A. Magnus. On convergence of sequences of linear fractional transformations. *Math. Zeit.*, 115(1):11–17, 1970.

*E-mail address:* `Ian.Short@nuim.ie`

MATHEMATICS DEPARTMENT, NATIONAL UNIVERSITY OF IRELAND, MAYNOOTH,  
COUNTY KILDARE, IRELAND