

# Open Research Online

---

The Open University's repository of research publications and other research outputs

## Norms of Möbius maps

### Journal Item

How to cite:

Beardon, Alan F. and Short, Ian (2010). Norms of Möbius maps. *Bulletin of the London Mathematical Society*, 422(3) pp. 499–505.

For guidance on citations see [FAQs](#).

© 2010 London Mathematical Society

Version: Accepted Manuscript

Link(s) to article on publisher's website:  
<http://dx.doi.org/doi:10.1112/blms/bdq015>

---

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

---

[oro.open.ac.uk](http://oro.open.ac.uk)

# Norms of Möbius maps

Alan F. Beardon and Ian Short

## ABSTRACT

We derive inequalities between the matrix, chordal, hyperbolic, three-point, and unitary norms of a Möbius map. These extend inequalities proved earlier by Gehring and Martin.

## 1. Introduction

A Möbius transformation  $z \mapsto (az + b)/(cz + d)$ , where  $ad - bc \neq 0$ , is a homeomorphism of the extended complex plane  $\mathbb{C}_\infty$  onto itself with the chordal metric  $q$  given by

$$q(z, w) = \frac{2|z - w|}{\sqrt{1 + |z|^2}\sqrt{1 + |w|^2}}, \quad q(z, \infty) = \frac{2}{\sqrt{1 + |z|^2}},$$

and also a conformal isometry of the upper half-space  $\mathbb{H}^3$  of  $\mathbb{R}^3$  endowed with the hyperbolic metric  $ds = |dx|/x_3$ . In [3, 4] Gehring and Martin derived inequalities between the matrix norm, the chordal norm and the hyperbolic norm of a Möbius map (all of which are defined below). Here we introduce two more norms and study the relationships between these five norms. It is known that if a sequence of Möbius transformations converges at three distinct points to three distinct values, then it converges uniformly on  $\mathbb{C}_\infty$  to a Möbius transformation. The work in this paper originated in an attempt to find a proof of this result which exhibits an explicit rate of convergence, and our inequalities provide such an estimate.

The group  $\mathcal{M}$  of Möbius maps is equipped with the supremum metric  $d$ , where

$$d(f, g) = \sup \{q(f(z), g(z)) : z \in \mathbb{C}_\infty\}, \quad f, g \in \mathcal{M},$$

so that  $d(f_n, f) \rightarrow 0$  if and only if  $f_n \rightarrow f$  uniformly on  $\mathbb{C}_\infty$ . Following Gehring and Martin ([3, 4]), we define the *chordal norm* of a Möbius map  $f$  to be  $d(f, I)$ , where  $I$  denotes the identity map: thus

$$d(f, I) = \sup \{q(f(z), z) : z \in \mathbb{C}_\infty\}.$$

Given a Möbius map  $f$ , we can write

$$f(z) = \frac{az + b}{cz + d}, \quad A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad ad - bc = 1, \quad (1.1)$$

where  $A$  is determined to within a factor  $\pm 1$ . The *matrix norm* of  $f$  is  $\|f\|$ , or  $\|A\|$ , where

$$\|f\| = \|A\| = \sqrt{|a|^2 + |b|^2 + |c|^2 + |d|^2},$$

and this is independent of the factor  $\pm 1$ . Gehring and Martin introduce the norm

$$m(f) = \|A - A^{-1}\| = \sqrt{2|a - d|^2 + 4|b|^2 + 4|c|^2},$$

but more often than not use the expression  $m(f)/\|A\|$ . We shall combine these ideas and define

$$M(f) = \frac{\|A - \det(A)A^{-1}\|}{\|A\|},$$

where  $A$  is now any matrix which represents  $f$  as in (1.1), except that we no longer insist that  $ad - bc = 1$ . Note that  $M(f)$  is independent of the choice of  $A$  from  $\text{GL}(2, \mathbb{C})$  which represents  $f$ , and if  $\det(A) = 1$ , then  $M(f) = m(f)/\|A\|$ . In this notation, Gehring and Martin prove (see [4, (3.35)])

$$M(f) \leq d(f, I) \leq \sqrt{2}M(f). \quad (1.2)$$

Both constants in these inequalities are best possible.

Guided by the fact that convergence at three points implies uniform convergence we now introduce the *three-point norm*

$$\varepsilon(f) = \max \{q(f(1), 1), q(f(\omega), \omega), q(f(\omega^2), \omega^2)\},$$

where  $\omega = e^{2\pi i/3}$ . In principle, we could take any three points here, and originally the authors proved an estimate similar to (1.3) below using the points 0, 1 and  $\infty$  instead of 1,  $\omega$  and  $\omega^2$ , and with a constant on the right between 2 and 3 (and not best possible). The authors thank the referee for providing the better inequality  $M(f) \leq \sqrt{2}\varepsilon(f)$  in (1.3) and its proof.

**THEOREM 1.1.** *For any Möbius transformation  $f$ ,*

$$\frac{\varepsilon(f)}{\sqrt{2}} \leq M(f) \leq \sqrt{2}\varepsilon(f), \quad (1.3)$$

and

$$\varepsilon(f) \leq d(f, I) \leq 2\varepsilon(f). \quad (1.4)$$

Further, each constant in each of the four inequalities is best possible.

In the case of parabolic Möbius maps, we can say more than Theorem 1.1. Let  $\varphi$  be the stereographic projection of  $\mathbb{C}_\infty$  onto the unit sphere  $\mathbb{S}$  in  $\mathbb{R}^3$ . Then  $p$  and  $\hat{p}$  in  $\mathbb{C}_\infty$  are *antipodal points* if and only if  $\varphi(p)$  and  $\varphi(\hat{p})$  are the end-points of a diameter of  $\mathbb{S}$  (that is,  $q(p, \hat{p}) = 2$ ). In particular, the points 0 and  $\infty$  are antipodal, and we define

$$\varepsilon_0(f) = \max \{q(f(0), 0), q(f(\infty), \infty)\}.$$

The choice of the pair of antipodal points used to define  $\varepsilon_0$  is insignificant (see Section 2).

**THEOREM 1.2.** *For any parabolic Möbius transformation  $f$ ,*

$$\frac{\varepsilon_0(f)}{\sqrt{2}} \leq M(f) \leq \sqrt{2}\varepsilon_0(f), \quad (1.5)$$

and

$$\varepsilon_0(f) \leq d(f, I) \leq 2\varepsilon_0(f). \quad (1.6)$$

Further, each constant in each of the four inequalities is best possible.

Next, the *hyperbolic norm* of  $f$  is  $\rho(j, f(j))$ , where  $\rho$  is the hyperbolic metric on  $\mathbb{H}^3$ , and  $j = (0, 0, 1)$ , and Gehring and Martin obtained an inequality that is equivalent to

$$2 \tanh \frac{1}{2}\rho(j, f(j)) \leq d(f, I). \quad (1.7)$$

(see [4, (1.19) and Theorem 3.19]).

A Möbius map  $u$  is a *unitary map* if its conjugate  $\varphi u \varphi^{-1}$  is a rotation of the sphere  $\mathbb{S}$  or, equivalently, if  $u(j) = j$ . Now suppose that  $u$  is unitary, and apply (1.7) to  $f u^{-1}$  instead of  $f$ ; this gives

$$2 \tanh \frac{1}{2} \rho(j, f(j)) \leq d(f, u).$$

We let  $\mathcal{U}$  be the subgroup of unitary maps, and we call  $d(f, \mathcal{U})$  the *unitary norm* of  $f$ , where

$$d(f, \mathcal{U}) = \inf \{d(f, u) : u \in \mathcal{U}\};$$

this measures how far  $f$  is from the subgroup  $\mathcal{U}$ . The Gehring Martin inequality (1.7) implies that

$$2 \tanh \frac{1}{2} \rho(j, f(j)) \leq d(f, \mathcal{U}), \tag{1.8}$$

and our last result is that this inequality is, in fact, an equality.

**THEOREM 1.3.** *In the notation above,*

$$d(f, \mathcal{U}) = 2 \tanh \frac{1}{2} \rho(j, f(j)) = 2 \sqrt{\frac{\|f\|^2 - 2}{\|f\|^2 + 2}}.$$

We discuss unitary maps in Section 2. The proofs of Theorems 1.1, 1.2 and 1.3 are given in Sections 3, 4 and 5, respectively. Finally, in Section 6, we return to the original motivation for this paper and obtain a quantitative version of the theorem that says convergence at three points implies uniform convergence.

## 2. Unitary maps

We devote this section to a brief discussion of unitary maps, and the reader is referred to [1] for details. A Möbius map  $u$  is a *unitary map* if and only if  $\varphi u \varphi^{-1}$  is a rotation of the sphere  $\mathbb{S}$ , and a unitary Möbius map is clearly a chordal isometry. Also, a Möbius map is unitary if and only if, in its action on  $\mathbb{H}^3$ , it fixes  $j = (0, 0, 1)$ . Since

$$\|f\|^2 = 2 \cosh \rho(j, f(j)) \tag{2.1}$$

for any Möbius map  $f$ , we also see that  $f$  is unitary if and only if  $\|f\|^2 = 2$ .

The metric  $d$  is *right invariant*: for all Möbius  $f, g$  and  $h$ ,

$$d(fh, gh) = d(f, g).$$

The Möbius map  $h$  induces the left invariance property  $d(hf, hg) = d(f, g)$  for all  $f$  and  $g$  if and only if  $h$  is unitary.

Unitary maps have other useful invariance properties; for example, if  $X$  is any  $2 \times 2$  complex matrix (singular or non-singular), and if  $U$  is a unitary matrix (corresponding to a unitary Möbius map), then  $\|UX\| = \|X\| = \|XU\|$ . Now consider any Möbius map  $f$ , and any unitary Möbius map  $u$ . Then,  $\|u f u^{-1}\| = \|f\|$ . Also, since  $u$  is a chordal isometry,

$$d(u f u^{-1}, I) = d(u f u^{-1}, u u^{-1}) = d(f, I).$$

These facts imply that any relationship between  $d(f, I)$  and  $\|f\|$  is invariant under conjugation by a unitary map, and this leads to a considerable simplification of our arguments. Since a unitary map  $u$  fixes  $j$  and is a hyperbolic isometry, we see that, for any Möbius map  $f$ ,

$$\rho(u f u^{-1}(j), j) = \rho(f(j), j),$$

so similar comments apply to this norm too.

In conclusion, these remarks show that we could replace the three points  $1$ ,  $\omega$  and  $\omega^2$  in the definition of  $\varepsilon$  by any three points equally spaced around a great circle and Theorem 1.1 would remain true. Similarly, Theorem 1.2 would remain true if  $0$  and  $\infty$  are replaced by any pair of antipodal points.

### 3. The proof of Theorem 1.1

To establish the four inequalities in (1.3) and (1.4), it suffices to prove the *right hand* inequality of (1.3) and the *left hand* inequality of (1.4), because the remaining inequalities follow from these two inequalities together with  $d(f, I) \leq \sqrt{2}M(f)$  from (1.2). The left hand inequality of (1.4),  $\varepsilon(f) \leq d(f, I)$ , follows immediately from the definition of  $\varepsilon$ , so we have only to prove the right hand inequality  $M(f) \leq \sqrt{2}\varepsilon(f)$  of (1.3). In fact, we shall prove the following slightly stronger result.

**PROPOSITION 3.1.** *There are positive numbers  $\mu_0$ ,  $\mu_1$  and  $\mu_2$ , with  $\mu_0 + \mu_1 + \mu_2 = 1$ , such that*

$$M(f)^2 \leq 2 \left[ \mu_0 q(f(1), 1)^2 + \mu_1 q(f(\omega), \omega)^2 + \mu_2 q(f(\omega^2), \omega^2)^2 \right] \leq 2\varepsilon(f)^2. \quad (3.1)$$

*Proof.* If  $|z| = 1$  then

$$(|az + b|^2 + |cz + d|^2) = \|A\|^2 + 2\operatorname{Re}[(a\bar{b} + c\bar{d})z],$$

and since  $1 + \omega + \omega^2 = 0$ , this gives

$$\sum_{z^3=1} (|az + b|^2 + |cz + d|^2) = 3\|A\|^2. \quad (3.2)$$

In a similar way we get

$$\sum_{z^3=1} |(az + b) - z(cz + d)|^2 = 3(|a - d|^2 + |b|^2 + |c|^2).$$

Now define

$$\mu_j = \frac{|a\omega^j + b|^2 + |c\omega^j + d|^2}{3\|A\|^2}, \quad j = 0, 1, 2. \quad (3.3)$$

We see from (3.2) that  $\mu_0 + \mu_1 + \mu_2 = 1$ . Observe that

$$2|(a\omega^j + b) - \omega^j(c\omega^j + d)|^2 = 3\mu_j\|A\|^2 q(f(\omega^j), \omega^j)^2,$$

which means that

$$\sum_{j=0}^2 \mu_j q(f(\omega^j), \omega^j)^2 = \frac{2}{\|A\|^2} (|a - d|^2 + |b|^2 + |c|^2),$$

and this gives (3.1) since

$$M(f)^2 = \frac{2}{\|A\|^2} (|a - d|^2 + 2|b|^2 + 2|c|^2).$$

□

To show that the constants in the four inequalities from (1.3) and (1.4) are best possible it suffices to prove that the constants in the *left hand* inequality of (1.3) and the *right hand* inequality of (1.4) are best possible, because then, using the inequality  $d(f, I) \leq \sqrt{2}M(f)$  from (1.2), we see that all four constants are best possible. For example, if we show that the constant

2 in  $d(f, I) \leq 2\varepsilon(f)$  is best possible, then the constants in the two inequalities  $d(f, I) \leq \sqrt{2}M(f)$  and  $M(f) \leq \sqrt{2}\varepsilon(f)$  must also be best possible.

To see that the constant  $\sqrt{2}$  in the left hand inequality of (1.3),  $\varepsilon(f) \leq \sqrt{2}M(f)$ , is best possible, consider the following sequence of Möbius transformations:

$$f_n(z) = \frac{nz - (n-1)}{-(n+1)z + n}, \quad n = 1, 2, \dots$$

Then  $f_n(1) = -1$ , so that  $\varepsilon(f_n) = 2$ , and one can check that  $M(f_n) \rightarrow \sqrt{2}$  as  $n \rightarrow \infty$ .

To see that the constant 2 in the right hand inequality of (1.4),  $d(f, I) \leq 2\varepsilon(f)$ , is best possible, consider the following one parameter group of Möbius transformations with fixed points  $-1$  and  $1$ :

$$f_t(z) = \frac{(1+t)z + (1-t)}{(1-t)z + (1+t)}, \quad t \in \mathbb{R}.$$

Notice that  $f_t(z) \rightarrow -1$  as  $t \rightarrow \infty$  for all points  $z$  other than  $1$ . Therefore

$$\varepsilon(f_t) \rightarrow 1, \quad d(f_t, I) \rightarrow 2,$$

as  $t \rightarrow \infty$ , and this shows that the constant 2 is best possible.

#### 4. The proof of Theorem 1.2

To establish the four inequalities in (1.5) and (1.6), it suffices to prove the *right hand* inequality of (1.5) and the *left hand* inequality of (1.6), because the remaining inequalities follow from these two inequalities together with  $d(f, I) \leq \sqrt{2}M(f)$  from (1.2). The left hand inequality of (1.6),  $\varepsilon_0(f) \leq d(f, I)$ , follows immediately from the definition of  $\varepsilon_0$ , so we have only to prove the right hand inequality of (1.5),  $M(f) \leq \sqrt{2}\varepsilon_0(f)$ .

Since  $f$  is parabolic we may assume, with the notation of (1.1), that  $ad - bc = 1$  and  $a + d = 2$ . This means that

$$4bc = 4(ad - 1) = -4(a - 1)^2 = -(a - d)^2.$$

Hence

$$\begin{aligned} M(f)^2 &= \frac{2|a - d|^2 + 4|b|^2 + 4|c|^2}{|a|^2 + |b|^2 + |c|^2 + |d|^2} \\ &\leq 2 \left( \frac{4|b|^2 + 4|c|^2}{|a|^2 + |b|^2 + |c|^2 + |d|^2} \right) \\ &\leq 2 \max \left\{ \frac{4|b|^2}{|b|^2 + |d|^2}, \frac{4|c|^2}{|a|^2 + |c|^2} \right\} \\ &= 2\varepsilon_0(f)^2, \end{aligned}$$

as required.

To show that the constants in the four inequalities in (1.5) and (1.6) are best possible it suffices to prove that the constants in the *left hand* inequality of (1.5) and the *right hand* inequality of (1.6) are best possible, because then, using the inequality  $d(f, I) \leq \sqrt{2}M(f)$  from (1.2), we see that all four constants are best possible.

To see that the constant  $\sqrt{2}$  in the left hand inequality of (1.3),  $\varepsilon_0(f) \leq \sqrt{2}M(f)$ , is best possible, consider the maps  $f_t(z) = z + t$  for  $t > 0$ . One can check that  $\varepsilon_0(f_t)/M(f_t) \rightarrow \sqrt{2}$  as  $t \rightarrow 0$ . To see that the constant 2 in the right hand inequality of (1.4),  $d(f, I) \leq 2\varepsilon_0(f)$ , is best possible, consider the following one parameter group of parabolic Möbius transformations with fixed point  $i$ :

$$f_t(z) = \frac{(1+it)z + t}{tz + (1-it)}, \quad t \in \mathbb{R}.$$

As  $t \rightarrow 0$  we find that  $q(f_t(0), 0) \sim 2t$ ,  $q(f_t(\infty), \infty) \sim 2t$  and  $q(f_t(-i), -i) \sim 4t$ . This means that  $\limsup_{t \rightarrow 0} d(f_t, I)/\epsilon_0(f_t) \geq 2$ . Therefore the constant 2 in  $d(f, I) \leq 2\epsilon_0(f)$  is best possible.

### 5. The proof of Theorem 1.3

We begin with a decomposition result for a general Möbius  $f$ ; this is a straightforward consequence of the standard results on isometric spheres.

**THEOREM 5.1.** *Each Möbius map  $f$  can be represented in the form  $f = ug$ , where  $u$  is a unitary map, and  $g$  is a hyperbolic map with antipodal fixed points (or  $I$ ).*

*Proof.* We may assume that  $f$  is not unitary (else we take  $u = f$  and  $g = I$ ). Then the action of the conjugate map  $f^* = \varphi f \varphi^{-1}$  on the unit ball is given by  $f^* = \alpha\beta$ , where  $\beta$  is the inversion in the isometric sphere  $\mathcal{S}$  of  $f$ , and  $\alpha$  is some orthogonal map. Let  $\ell$  be the Euclidean line that passes through 0 and the centre of  $\mathcal{S}$ , and let  $\gamma$  be the reflection in the plane through 0 that is orthogonal to  $\ell$ . Then  $f = (\alpha\gamma)(\gamma\beta)$ , where  $\alpha\gamma$  is unitary and  $\gamma\beta$  is hyperbolic with antipodal fixed points.  $\square$

We can now complete the proof of Theorem 1.3, and we first prove this in the case of a hyperbolic map with antipodal fixed points.

**LEMMA 5.2.** *If  $g$  is hyperbolic with antipodal fixed points then*

$$d(g, \mathcal{U}) = 2 \tanh \frac{1}{2} \rho(j, g(j)).$$

*Proof.* We know from (1.8) that

$$2 \tanh \frac{1}{2} \rho(j, g(j)) \leq d(g, \mathcal{U}).$$

Also, Gehring and Martin prove in [4, Theorem 3.19] that equality holds in (1.7) when  $f$  is hyperbolic with antipodal fixed points. Thus  $2 \tanh \frac{1}{2} \rho(j, g(j)) = d(g, I) \geq d(g, \mathcal{U})$ .  $\square$

Now let  $f$  be a general Möbius map, and write  $f = ug$ , where  $u$  is unitary and  $g$  is hyperbolic with antipodal fixed points (or  $I$ ). Then

$$d(f, \mathcal{U}) = d(g, \mathcal{U}) = 2 \tanh \frac{1}{2} \rho(j, g(j)) = 2 \tanh \frac{1}{2} \rho(j, f(j))$$

as required. Finally, from (2.1) we obtain

$$d(f, \mathcal{U}) = 2 \sqrt{\frac{\|f\|^2 - 2}{\|f\|^2 + 2}}.$$

### 6. The convergence theorem

We finish by returning to the original motivation for this paper, namely that if a sequence of Möbius maps converges at three distinct points to three distinct values, then it converges uniformly on  $\mathbb{C}_\infty$  to a Möbius map. Theorem 1.1 implies that if a sequence of Möbius maps converges to  $I$  on  $\{1, \omega, \omega^2\}$ , then it converges to  $I$  uniformly on  $\mathbb{C}_\infty$ . The extension to the general case is easy because, for any Möbius map  $h$ ,

$$\|h\|^{-2} q(z, w) \leq q(h(z), h(w)) \leq \|h\|^2 q(z, w)$$

[2, pages 543–544]. Suppose that  $z_1, z_2$  and  $z_3$  are distinct, and that a sequence  $g_n$  of Möbius maps satisfies  $g_n(z_j) \rightarrow w_j, j = 1, 2, 3$ , where the  $w_j$  are distinct. We can choose Möbius maps  $r$  and  $s$  that map  $1, \omega$  and  $\omega^2$  to  $z_1, z_2$  and  $z_3$ , and  $w_1, w_2$  and  $w_3$ , respectively, and then

$$\begin{aligned} d(g_n, sr^{-1}) &\leq \|s\|^2 d(s^{-1}g_n, s^{-1}sr^{-1}) \\ &= \|s\|^2 d(s^{-1}g_n r, I) \\ &\leq 2 \|s\|^2 \varepsilon(s^{-1}g_n r) \\ &\leq 2 \|s\|^4 \max \{q(g_n(z_j), w_j) : j = 1, 2, 3\}. \end{aligned}$$

We deduce that  $g_n \rightarrow sr^{-1}$  with the given rate of convergence.

### References

1. A. F. BEARDON, *The geometry of discrete groups* (Graduate Texts in Mathematics, 91, Springer-Verlag, New York, 1983).
2. A. F. BEARDON, ‘Continued fractions, discrete groups and complex dynamics’, *Comput. Methods and Funct. Theory* 1 (2001), no. 2, 535–594.
3. F. W. GEHRING AND G. J. MARTIN, ‘The matrix and chordal norms of Möbius transformations’, *Complex Analysis*, 51–59 (Birkhäuser, Basel, 1988).
4. F. W. GEHRING AND G. J. MARTIN, ‘Inequalities for Möbius transformations and discrete groups’, *J. Reine Angew. Math.* 418 (1991), 31–76.

*Alan F. Beardon and Ian Short*  
*Department of Pure Mathematics and*  
*Mathematical Statistics*  
*Centre for Mathematical Sciences*  
*Wilberforce Road*  
*Cambridge CB3 0JB*  
*United Kingdom*

a.f.beardon@dpmms.cam.ac.uk  
 ims25@cam.ac.uk