

Open Research Online

The Open University's repository of research publications and other research outputs

Raman variation of organic matter in large cluster IDPs and meteoritic IOM

Conference or Workshop Item

How to cite:

Starkey, N. A.; Franchi, I. A. and Alexander, C. M. O'D. (2011). Raman variation of organic matter in large cluster IDPs and meteoritic IOM. In: 74th Annual Meeting of the Meteoritical Society, 8-12 Aug 2011, London, UK.

For guidance on citations see [FAQs](#).

© 2011 The Meteoritical Society

Version: Version of Record

Link(s) to article on publisher's website:

<http://www.lpi.usra.edu/meetings/metsoc2011/pdf/5423.pdf>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

RAMAN VARIATION OF ORGANIC MATTER IN LARGE CLUSTER IDPs AND METEORITIC IOM.

N. A. Starkey¹, I. A. Franchi¹ and C. M. O'D. Alexander².
¹PSSRI, Open University, Milton Keynes, MK7 6AA. UK.
Email: n.starkey@open.ac.uk. ²DTM, Carnegie Institution of
Washington, 5241 Broad Branch Road, Washington DC 20015,
USA.

Introduction: CP-IDPs display a range of primitive features indicative of a cometary origin. The organic material (OM) present in IDPs may be the result of formation processes that occurred across a large volume of the protoplanetary disk to small-scale on the cometary body itself. Raman microscopy offers a non-destructive approach for determining general characteristics of the OM in IDPs. Compared to the large number of potential parent bodies or the huge volume of the protoplanetary disk that may have contributed to the formation of comets, the number of IDPs analysed to date remains relatively small. Raman measurements were performed on 9 IDPs (from 5 large cluster particles; collectors L2005 and L2006) offering the opportunity to investigate the variability of the OM on a number of scales. Previous Raman analyses of IDPs and IOM from meteorites have used a number of different wavelengths, predominantly green. While it has been shown that D band position (ω D) is influenced by excitation λ [1], it is rarely quantified. Interestingly, the wavelength dependent shift in the ω G is known to vary with degree of ordering of OM. Multi-wavelength analyses made here offer an independent measure of organic maturity.

Results: Meteoritic IOM (samples as in [2]) and IDPs mapped using 514nm and 473nm lasers showed a reproducible downshift of 10-13 cm^{-1} in ω D. For ω G, the IDPs displayed a larger downshift ($\sim 4 \text{ cm}^{-1}$) than the IOM (negligible). The range of ω D and ω G in any one IDP cluster is limited, with values comparable to meteoritic IOM. For particles within individual clusters, ω D varies by $< 6 \text{ cm}^{-1}$ and ω G by $< 3 \text{ cm}^{-1}$. Each cluster IDP displays distinctive internal peak parameter co-variation.

Discussion: The large Γ D and Γ G, high ω D, low ω G and ID/IG ratios of 0.9-1.3 indicate that the IDPs are all primitive. The samples plot near to the CI, CM and CR field. The dispersion in ω G with excitation λ is related to the degree of disordering of the carbon [1], and therefore the greater ω G dispersion observed for the IDPs suggests that the OM in the IDPs may be more disordered than meteoritic IOM, in contrast to their similar peak parameters. The IDPs selected for this study do not appear to be as primitive as Grigg-Skjellerup Collection (GSC) particles [3], but are comparable to other non-GSC IDPs [4].

While great isotopic and structural variability exists within individual particles at the sub- μm scale (e.g. [3]), these results indicate that the OM within IDPs is reasonably homogeneous at the few to tens of μm -scale, and that there is greater variability between particles than within. The variability between particles most likely reflects accretion from different reservoirs, potentially on the scale of individual comets given the distinctive signatures from GSC-IDPs [3]. Secondary processes on the comet, in interplanetary space or during atmospheric entry may also play a role despite low peak metamorphic temperatures ($< 200^\circ\text{C}$ (after [5])).

References: [1] Ferrari, A.C. & Robertson, J. 2001. *Phys. Rev. B.* 63:075414. [2] Alexander, C.M. O'D. et al. 2007. *GCA* 71, 4380-4403. [3] Busemann H. et al. 2009. *Earth Planet. Sci. Lett.*, 288:44-57. [4] Davidson, J. 2009. *PhD thesis, Open Univ.* [5] Busemann, H. et al. 2007. *Met. Planet. Sci.* 42:1387-1416.