

Open Research Online

The Open University's repository of research publications and other research outputs

Entire functions for which the escaping set is a spider's web

Journal Item

How to cite:

Sixsmith, D. J. (2011). Entire functions for which the escaping set is a spider's web. *Mathematical Proceedings of the Cambridge Philosophical Society*, 151(3) pp. 551–571.

For guidance on citations see [FAQs](#).

© 2011 Cambridge Philosophical Society

Version: Version of Record

Link(s) to article on publisher's website:
<http://dx.doi.org/doi:10.1017/S0305004111000582>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's [data policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

Entire functions for which the escaping set is a spider's web

BY D. J. SIXSMITH

*Department of Mathematics and Statistics, The Open University,
Walton Hall, Milton Keynes MK7 6AA.
e-mail: sixsmiths@hotmail.co.uk*

(Received 11 April 2011; revised 29 June 2011)

Abstract

We construct several new classes of transcendental entire functions, f , such that both the escaping set, $I(f)$, and the fast escaping set, $A(f)$, have a structure known as a spider's web. We show that some of these classes have a degree of stability under changes in the function. We show that new examples of functions for which $I(f)$ and $A(f)$ are spiders' webs can be constructed by composition, by differentiation, and by integration of existing examples. We use a property of spiders' webs to give new results concerning functions with no unbounded Fatou components.



1. Introduction

Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be a transcendental entire function, and denote by f^n , $n \in \mathbb{N}$, the n th iterate of f . The *Fatou set* $F(f)$ is defined as the set of points $z \in \mathbb{C}$ such that $(f^n)_{n \in \mathbb{N}}$ forms a normal family in a neighborhood of z . The *Julia set* $J(f)$ is the complement of $F(f)$. An introduction to the properties of these sets can be found in [6].

This paper concerns the *escaping set* $I(f)$ and the *fast escaping set* $A(f)$, introduced respectively by Eremenko [10], and Bergweiler and Hinkkanen [7]. These sets are defined as follows:

$$I(f) = \{z : f^n(z) \rightarrow \infty \text{ as } n \rightarrow \infty\},$$

and

$$A(f) = \{z : \text{there exists } L \in \mathbb{N} \text{ such that } |f^{n+L}(z)| \geq M^n(R, f), \text{ for } n \in \mathbb{N}\};$$

see [20] for this form of the definition of $A(f)$. Here,

$$M(r, f) = \max_{|z|=r} |f(z)|, \quad \text{for } r > 0,$$

$M^n(r, f)$ denotes repeated iteration of $M(r, f)$ with respect to the variable r , and $R > 0$ can be taken to be any value such that $M(r, f) > r$ for $r \geq R$. For simplicity, we only write down this restriction on R in formal statements of results – elsewhere this should be assumed to be true.

Rippon and Stallard [20] gave a detailed account of many properties of $A(f)$. Their arguments were frequently based on properties of the set

$$A_R(f) = \{z : |f^n(z)| \geq M^n(R, f), \text{ for } n \in \mathbb{N}\}.$$

In particular they showed that $A_R(f)$, $A(f)$ and $I(f)$ can have a structure known as a spider’s web, and that if $A_R(f)$ is a spider’s web then so are $A(f)$ and $I(f)$. They defined a spider’s web as follows:

Definition. A set E is an (infinite) spider’s web if E is connected and there exists a sequence $(G_n)_{n \in \mathbb{N}}$ of bounded simply connected domains with $G_n \subset G_{n+1}$, for $n \in \mathbb{N}$, $\partial G_n \subset E$, for $n \in \mathbb{N}$, and $\bigcup_{n \in \mathbb{N}} G_n = \mathbb{C}$.

Functions for which $A_R(f)$ is a spider’s web have a number of strong dynamical properties. For example, if $A_R(f)$ is a spider’s web then f has no unbounded Fatou components [20, theorem 1.5(b)] and $A(f)^c$ has uncountably many components [17, theorem 1.2]. Hence, it is desirable to determine those functions for which $A_R(f)$ is a spider’s web, and in [20, section 8] several classes of such functions were given. A further class was given by Mihaljević–Brandt and Peter [16].

In this paper we give additional classes of examples. First, in Section 2, we prove several new results concerning regular growth conditions, which we use in later sections. These results may also be of independent interest.

In Section 3, we demonstrate a technique for constructing new transcendental entire functions for which $A_R(f)$ is a spider’s web by taking finite compositions of functions that satisfy a minimum modulus condition and a regularity condition.

In Section 4, we show that in certain circumstances when $A_R(f)$ is a spider’s web, then so is $A_R(P(f(Q(z)), z))$, where P, Q are polynomials, and so also is $A_R(f + h)$, where the entire function h has smaller growth, in some sense, than f . These results allow us to construct a large class of functions for which $A_R(f)$ is a spider’s web. They also show that the property of having an $A_R(f)$ spider’s web can be stable under changes in f , unlike many other dynamical properties.

In Section 5, we establish a technique for constructing a large class of transcendental entire functions of finite order for which $A_R(f)$ is a spider’s web, by modifying the power series of a transcendental entire function of finite order. This technique is a generalisation of the method used to construct some of the examples in [20]. We show that this class of examples can be extended by differentiation or integration. By combining the results of Sections 3, 4 and 5, we give an unexpectedly simple function for which $A_R(f)$ is a spider’s web.

In Section 6, we present a technique for constructing new transcendental entire functions, of infinite order and with large gaps in their power series, for which $A_R(f)$ is a spider’s web.

Finally, in Section 7, we relate our results to previous work on classes of transcendental entire functions which have no unbounded Fatou components.

Throughout the paper we use the following three facts about the maximum modulus function $M(r, f)$ of a transcendental entire function f . The first two are well known, and the third is given in [19, lemma 2.2]:

$$\frac{\log M(r, f)}{\log r} \rightarrow \infty \text{ as } r \rightarrow \infty, \tag{1.1}$$

$$\text{if } k > 1 \text{ then } \frac{M(kr, f)}{M(r, f)} \rightarrow \infty \text{ as } r \rightarrow \infty, \tag{1.2}$$

and there exists $R > 0$ such that

$$M(r^c, f) \geq M(r, f)^c, \quad \text{for } r \geq R, \ c > 1. \tag{1.3}$$

We also use the minimum modulus function defined by

$$m(r, f) = \min_{|z|=r} |f(z)|, \quad \text{for } r > 0. \tag{1.4}$$

Finally we use the following notation for a disc

$$B(z_0, r) = \{z : |z - z_0| < r\}, \quad \text{for } z_0 \in \mathbb{C}, r > 0.$$

2. New results on regularity

In this section we set out conditions which ensure that $A_R(f)$ is a spider's web. Many of these conditions require some form of regularity of growth. We prove several new results concerning forms of regularity of growth, which enable us to construct examples of functions with an $A_R(f)$ spider's web later in the paper.

A pair of conditions that are together necessary and sufficient for $A_R(f)$ to be a spider's web were obtained in [20, theorem 8·1].

THEOREM 2·1. *Let f be a transcendental entire function and let $R > 0$ be such that $M(r, f) > r$ for $r \geq R$. Then $A_R(f)$ is a spider's web if and only if there exists a sequence $(G_n)_{n \geq 0}$ of bounded simply connected domains such that, for all $n \geq 0$,*

$$G_n \supset B(0, M^n(R, f)) \tag{2.1}$$

and

$$G_{n+1} \text{ is contained in a bounded component of } \mathbb{C} \setminus f(\partial G_n). \tag{2.2}$$

This result is very general, and so, in order to construct examples, the following, more readily applicable, sufficient conditions for $A_R(f)$ to be a spider's web were established in [20, corollary 8·3].

LEMMA 2·1. *Let f be a transcendental entire function and let $R > 0$ be such that $M(r, f) > r$ for $r \geq R$. Then $A_R(f)$ is a spider's web if, for some $m > 1$, $m \in \mathbb{R}$,*

(a) *there exists $R_0 > 0$ such that, for all $r \geq R_0$,*

$$\text{there exists } \rho \in (r, r^m) \text{ with } m(\rho, f) \geq M(r, f), \text{ and} \tag{2.3}$$

(b) *f has regular growth in the sense that there exists a sequence $(r_n)_{n \geq 0}$ with*

$$r_n \geq M^n(R, f) \text{ and } M(r_n, f) \geq r_{n+1}^m, \quad \text{for } n \geq 0. \tag{2.4}$$

We use the following condition, which is stronger than the regularity condition of Lemma 2·1(b), in order to construct a new class of functions with an $A_R(f)$ spider's web.

Definition. A transcendental entire function f is ψ -regular if, for each $m > 1$, $m \in \mathbb{R}$, there exist an increasing function ψ_m and $R_m > 0$ such that, for all $r \geq R_m$,

$$\psi_m(r) \geq r \text{ and } M(\psi_m(r), f) \geq (\psi_m(M(r, f)))^m. \tag{2.5}$$

For given $m > 1$, $m \in \mathbb{R}$, we call ψ_m a regularity function for f .

This condition is slightly stronger than one used in [19, theorem 5] in connection with transcendental entire functions with no unbounded Fatou components. That version did not require the regularity function to be increasing. However, all the regularity functions used in

[19, 20] are, in fact, increasing. It was shown in [20, section 8] that if f is ψ -regular, then it satisfies Lemma 2.1(b) for all $m > 1, m \in \mathbb{R}$.

We also use the following condition, which is stronger than ψ -regularity, in order to construct several classes of functions with an $A_R(f)$ spider's web.

Definition. Let $c > 0$. A transcendental entire function f is *log-regular, with constant c* , if the function $\phi(t) = \log M(e^t, f)$ satisfies

$$\frac{\phi'(t)}{\phi(t)} \geq \frac{1 + c}{t}, \quad \text{for large } t. \tag{2.6}$$

This condition was used by Anderson and Hinkkanen in [2, theorem 2], also in connection with transcendental entire functions with no unbounded Fatou components. The name log-regular was suggested by Aimo Hinkkanen in a private communication. The condition was also used in [20, section 8] in order to construct classes of functions with an $A_R(f)$ spider's web.

In [19, section 7] it was shown that if f is log-regular with constant c , then, for all $m > 1, m \in \mathbb{R}, f$ is ψ -regular with regularity function $\psi_m(r) = r^{m^{1/c}}$; see also Lemma 2.2 below. Hence if f is log-regular, then it satisfies Lemma 2.1(b) for all $m > 1, m \in \mathbb{R}$.

We now state three new results concerning ψ -regularity and log-regularity. The first concerns the composition of ψ -regular functions.

THEOREM 2.2. *Let f_1, f_2, \dots, f_k be transcendental entire functions. Suppose that, for all $j \in \{1, 2, \dots, k\}, f_j$ is ψ -regular with regularity function ψ_m for each $m > 1, m \in \mathbb{R}$. Let $g = f_1 \circ f_2 \circ \dots \circ f_k$. Then, for any $c > 1, g$ is ψ -regular with regularity function $c\psi_m$ for each $m > 1, m \in \mathbb{R}$.*

In particular it follows that ψ -regularity is preserved under iteration.

COROLLARY 2.1. *If f is a ψ -regular transcendental entire function, then so is f^n for all $n \in \mathbb{N}$.*

The second result relates to the composition of entire functions, one of which is log-regular.

THEOREM 2.3. *Let f_1, f_2, \dots, f_k be non-constant entire functions such that, for some $j \in \{1, 2, \dots, k\}, f_j$ is a log-regular transcendental entire function. Let $g = f_1 \circ f_2 \circ \dots \circ f_k$. Then g is log-regular.*

In particular it follows that log-regularity is preserved under iteration.

COROLLARY 2.2. *If f is a log-regular transcendental entire function, then so is f^n for all $n \in \mathbb{N}$.*

Note that Theorem 2.2 requires all functions to be ψ -regular transcendental entire functions, whereas Theorem 2.3 requires just one to be a log-regular transcendental entire function and the others only to be entire.

The third result shows that if f is log-regular, then so is any transcendental entire function with similar growth.

THEOREM 2.4. *Let f and g be transcendental entire functions. If f is log-regular and there exist $a_1, a_2 \geq 1$ and $R_0 > 0$ such that*

$$M(r^{a_1}, g) \geq M(r, f) \text{ and } M(r^{a_2}, f) \geq M(r, g), \quad \text{for } r \geq R_0, \tag{2.7}$$

then g is log-regular.

We need three preparatory lemmas to prove these results. The first lemma is from [18], and gives a necessary condition and a sufficient condition for f to be log-regular.

LEMMA 2.2. *Let f be a transcendental entire function.*

(a) *If f is log-regular, with constant c , then there is an $R_0 > 0$ such that, if $k > 1$ and $d = k^c$, then*

$$M(r^k, f) \geq M(r, f)^{kd}, \quad \text{for } r \geq R_0. \tag{2.8}$$

(b) *If (2.8) holds for some $d, k > 1$ and $R_0 > 0$, then f is log-regular.*

The second lemma comes from Wiman–Valiron theory, (see, for example, [13]), which was first used in connection with the escaping set by Eremenko [10]. We first need to introduce some terminology. Let $g(z) = \sum_{n=0}^{\infty} a_n z^n$ be a transcendental entire function. Define

$$\mu(r) = \sup_n |a_n| r^n = |a_N| r^N, \quad r > 0, \tag{2.9}$$

to be the *maximal term* of the power series, and call $N = N(r)$ the *central index*; if (2.9) holds for several N , we take $N(r)$ to be the largest of these. Note that $N(r)$ is increasing and $N(r) \rightarrow \infty$ as $r \rightarrow \infty$. Wiman–Valiron theory uses $\mu(r)$ to give results about the behaviour of g near points $z(r)$, $r > 0$, that satisfy

$$|z(r)| = r \text{ and } |g(z(r))| = M(r, g). \tag{2.10}$$

A key result of Wiman–Valiron theory is the following.

LEMMA 2.3. *Suppose that g is a transcendental entire function and $\alpha > 1/2$. For $r > 0$, let $z(r)$ be a point satisfying (2.10), and define*

$$D(r) = B \left(z(r), \frac{r}{(N(r))^\alpha} \right), \quad r > 0.$$

Then there exists a set $E \subset (0, \infty)$ with $\int_E 1/t \, dt < \infty$ such that, for $r \notin E$ and $z \in D(r)$,

$$g(z) = \left(\frac{z}{z(r)} \right)^{N(r)} g(z(r))(1 + \epsilon), \tag{2.11}$$

where $\epsilon = \epsilon(r, z) \rightarrow 0$ uniformly with respect to z as $r \rightarrow \infty$, $r \notin E$. In particular, if r is sufficiently large and $r \notin E$, then

$$g(D(r)) \supset \{w : |w| = M(r, g)\}. \tag{2.12}$$

We use Lemma 2.3 to prove a result on the behaviour of the maximum modulus of the composite of two entire functions.

LEMMA 2.4. *Suppose that f is a non-constant entire function and g is a transcendental entire function. Then, given $v > 1$, there exist $R_0, R_1 > 0$ such that*

$$M(vr, f \circ g) \geq M(M(r, g), f) \geq M(r, f \circ g), \quad \text{for } r \geq R_0, \tag{2.13}$$

and

$$M(vr, g \circ f) \geq M(M(r, f), g) \geq M(r, g \circ f), \quad \text{for } r \geq R_1. \tag{2.14}$$

Proof. We first prove (2.13). Let $\alpha > 1/2$, and let $N(r)$, E and $D(r)$ be related to g as in Lemma 2.3. Note that E has finite logarithmic measure, and $N(r) \rightarrow \infty$ as $r \rightarrow \infty$. Hence, for sufficiently large r , there exists $r' \in (r, (\nu + 1)r/2) \setminus E$, with

$$D(r') \subset B(0, \nu r) \text{ and } g(D(r')) \supset \{w : |w| = M(r', g)\}. \tag{2.15}$$

Let w be such that $|w| = M(r', g)$ and $|f(w)| = M(M(r', g), f)$. Then, by (2.15), there is a $z \in D(r')$ with $g(z) = w$. Hence

$$|(f \circ g)(z)| = M(M(r', g), f) > M(M(r, g), f).$$

The first part of (2.13) now follows, by (2.15). The second part of (2.13) is immediate.

Equation (2.14) follows in the same way if f is transcendental. Otherwise, suppose that f is a polynomial. Then, for sufficiently large r ,

$$f(B(0, \nu r)) \supset \{w : |w| = M(r, f)\}. \tag{2.16}$$

Let w be such that $|w| = M(r, f)$ and $|g(w)| = M(M(r, f), g)$. Then, by (2.16), there is a $z \in B(0, \nu r)$ with $f(z) = w$. Hence

$$|(g \circ f)(z)| = M(M(r, f), g).$$

The first part of (2.14) follows. The second part of (2.14) is immediate.

In passing, we note a related result discussed by Bergweiler and Hinkkanen [7, lemma 1] that, if we also have $g(0) = 0$, then

$$M(6r, f \circ g) \geq M(M(r, g), f), \quad \text{for } r > 0.$$

Now we are ready to prove Theorems 2.2, 2.3 and 2.4.

Proof of Theorem 2.2. Suppose that $m > 1$, $m \in \mathbb{R}$. We note first a general result. Suppose that f is ψ -regular with regularity function ψ_m , and let $\lambda > 1$. Then, for sufficiently large r , by (1.2) and (2.5),

$$M(\lambda \psi_m(r), f) \geq \lambda^m M(\psi_m(r), f) \geq (\lambda \psi_m(M(r, f)))^m. \tag{2.17}$$

Hence $\lambda \psi_m$ is also a regularity function for f .

Now, let $a = c^{1/(k-1)} > 1$. Suppose that $k = 2$. Then, for sufficiently large r ,

$$\begin{aligned} M(a\psi_m(r), f_1 \circ f_2) &\geq M(M(\psi_m(r), f_2), f_1) && \text{by Lemma 2.4} \\ &\geq M((\psi_m(M(r, f_2)))^m, f_1) && \text{by (2.5)} \\ &\geq M(\psi_m(M(r, f_2)), f_1)^m && \text{by (1.3)} \\ &\geq (\psi_m(M(M(r, f_2)), f_1))^{m^2} && \text{by (2.5)} \\ &\geq (\psi_m(M(r, f_1 \circ f_2)))^{m^2} && \text{since } \psi_m \text{ is increasing} \\ &\geq (a\psi_m(M(r, f_1 \circ f_2)))^m. \end{aligned}$$

Hence g is ψ -regular with regularity function $a\psi_m$. Finally, $c\psi_m = a\psi_m$, since $k = 2$. A similar argument with $f_1 \circ f_2$ and f_3 , both of which are ψ -regular with regularity function $a\psi_m$, by (2.17), gives the result for $k = 3$. The proof follows similarly for larger values of k .

Proof of Theorem 2.3. It is sufficient to prove the result for $k = 2$. Suppose then that $k = 2$ and that f_2 is log-regular. By Lemma 2.2(a) applied to f_2 , there are $k, d > 1$ and

$r_1 > 0$ such that

$$M(r^k, f_2) \geq M(r, f_2)^{kd}, \quad \text{for } r \geq r_1. \quad (2.18)$$

Choose ν such that $1 < \nu < d$, put $k' = k\nu > 1$ and $d' = d/\nu > 1$. Then, for sufficiently large r ,

$$\begin{aligned} M(r^{k'}, f_1 \circ f_2) &\geq M(\nu r^k, f_1 \circ f_2) \\ &\geq M(M(r^k, f_2), f_1) && \text{by Lemma 2.4} \\ &\geq M(M(r, f_2)^{kd}, f_1) && \text{by (2.18)} \\ &\geq M(M(r, f_2), f_1)^{kd} && \text{by (1.3)} \\ &\geq M(r, f_1 \circ f_2)^{k'd'} && \text{by choice of } k', d'. \end{aligned}$$

Thus $f_1 \circ f_2$ is log-regular by Lemma 2.2(b). If f_1 is log-regular but f_2 is not, then the proof that $f_1 \circ f_2$ is log-regular is very similar.

Proof of Theorem 2.4. Suppose that f is log-regular with constant c , and a_1, a_2 are as in (2.7). Choose $k > 1$ sufficiently large that $k^c > a_1 a_2$, put $d = k^c$, $k' = a_1 a_2 k > 1$, and $d' = d/a_1 a_2 > 1$. Then, for sufficiently large r ,

$$\begin{aligned} M(r^{k'}, g) &= M(r^{a_1 a_2 k}, g) \\ &\geq M(r^{a_2 k}, f) && \text{by (2.7)} \\ &\geq M(r^{a_2}, f)^{kd} && \text{by (2.8)} \\ &\geq M(r, g)^{kd} && \text{by (2.7)} \\ &= M(r, g)^{k'd'} && \text{by choice of } k', d'. \end{aligned}$$

Hence g is log-regular by Lemma 2.2(b).

We now prove several useful corollaries of Theorem 2.4. The first relates to the derivatives and integrals of log-regular functions.

COROLLARY 2.3. *Let f be a transcendental entire function. Then f is log-regular if and only if f' is log-regular.*

Proof. By integration and (1.3), for sufficiently large r ,

$$M(r^2, f') \geq rM(r, f') + |f(0)| \geq M(r, f). \quad (2.19)$$

On the other hand, by Cauchy's estimates, for sufficiently large r ,

$$M(r^2, f) \geq M(2r, f)/r \geq M(r, f'). \quad (2.20)$$

The result follows by Theorem 2.4, with $a_1 = a_2 = 2$.

The remaining corollaries of Theorem 2.4 are used later to give stability results about $A_R(f)$ spiders' webs. While they could be combined, they are stated separately for clarity. The first concerns addition of a function to a log-regular function.

COROLLARY 2.4. *Let f be a log-regular transcendental entire function, and let h be an entire function. Suppose that there exist $a \in (0, 1)$ and $R_0 > 0$ such that*

$$aM(r, f) \geq M(r, h), \quad \text{for } r \geq R_0. \quad (2.21)$$

Then $g = f + h$ is log-regular.

Proof. We observe that

$$(1 + a)M(r, f) \geq M(r, g) \geq (1 - a)M(r, f), \quad \text{for } r \geq R_0. \tag{2.22}$$

The result now follows by (1.3), and Theorem 2.4 with $a_1 = a_2 = 2$.

Note that, by (1.1), if h is a polynomial, then (2.21) is satisfied for any transcendental entire function f and any $a \in (0, 1)$.

The second corollary concerns a case where log-regularity is preserved under multiplication.

COROLLARY 2.5. *Let f be a log-regular transcendental entire function. Then $g(z) = zf(z)$ is log-regular.*

Proof. By (1.3) and (1.1), for sufficiently large r , $M(r^2, f) \geq M(r, f)^2 \geq M(r, g)$. Also, for sufficiently large r , $M(r, g) \geq M(r, f)$. The result follows, by Theorem 2.4 with $a_1 = 1$ and $a_2 = 2$.

Our final corollary is quite general.

COROLLARY 2.6. *Let f be a log-regular transcendental entire function. Let $P(w, z)$ be a polynomial, which is of degree at least one in w , and let $Q(z)$ be a polynomial of degree at least one. Then $g(z) = P(f(Q(z)), z)$ is log-regular.*

Proof. Suppose that

$$P(f(Q(z)), z) = af(Q(z))^{N_1}z^{N_2} + h(z) = g_0(z) + h(z),$$

where N_1 is the highest power of w in $P(w, z)$ and N_2 is the highest power of z corresponding to $f(Q(z))^{N_1}$. By Theorem 2.3, the function $z \mapsto af(Q(z))^{N_1}$ is log-regular. By Corollary 2.5, g_0 is log-regular. Since, by (1.1), we have

$$\frac{1}{2}M(r, g_0) \geq M(r, h), \quad \text{for large } r,$$

the result follows by Corollary 2.4.

3. Using composition to give functions for which $A_R(f)$ is a spider's web

In this section we demonstrate that $A_R(g)$ is a spider's web if $g = f_1 \circ f_2 \circ \dots \circ f_k$, and the entire functions $f_j, j \in \{1, 2, \dots, k\}$, satisfy certain conditions. We need a preparatory lemma before we can state the results. This lemma is a generalisation of Lemma 2.1, in which condition (a) is relaxed slightly. This condition was also used, independently, in [16].

LEMMA 3.1. *Let f be a transcendental entire function and let $R > 0$ be such that $M(r, f) > r$ for $r \geq R$. Then $A_R(f)$ is a spider's web if, for some $m > 1, m \in \mathbb{R}$,*

- (a) *there exists $R_0 > 0$ such that, for all $r \geq R_0$, there is a simply connected domain $G = G(r)$ with*

$$B(0, r) \subset G \subset B(0, r^m) \text{ and } |f(z)| \geq M(r, f), \quad \text{for } z \in \partial G, \tag{3.1}$$

and

- (b) *f has regular growth in the sense that there exists a sequence $(r_n)_{n \geq 0}$ with*

$$r_n \geq M^n(R, f) \text{ and } M(r_n, f) \geq r_{n+1}^m, \quad \text{for } n \geq 0. \tag{3.2}$$

Proof. Let m and R_0 be as in (a), and choose $(r_n)_{n \geq 0}$ satisfying (3.2) with $r_n > R_0$ for $n \geq 0$. For each $n \geq 0$, let $G_n = G(r_n)$.

First, by (3.1) and (3.2),

$$G_n \supset B(0, r_n) \supset B(0, M^n(R, f)), \quad \text{for } n \geq 0, \tag{3.3}$$

and so (G_n) satisfies (2.1).

Second, by (3.1) and (3.2), if $z \in \partial G_n$ then $|f(z)| \geq M(r_n, f) \geq r_{n+1}^m$. Thus $f(G_n)$ contains $B(0, r_{n+1}^m)$, since f maps points of $B(0, M^n(R, f))$ into $B(0, M^{n+1}(R, f)) \subset B(0, r_{n+1}^m)$. Now G_{n+1} is contained in $B(0, r_{n+1}^m)$ and so is contained in a bounded component of $\mathbb{C} \setminus f(\partial G_n)$. Thus (G_n) satisfies (2.2). Hence, by Theorem 2.1, $A_R(f)$ is a spider's web.

We note that if P is a non-constant polynomial, then P satisfies Lemma 3.1(a) for every $m > 1$, $m \in \mathbb{R}$, taking $G(r) = B(0, r^\alpha)$, where $\alpha \in (1, m)$, and a suitable R_0 .

We now state the main results of this section. The first relates to the composition of ψ -regular functions, and the second relates to the composition of entire functions, one of which is log-regular.

THEOREM 3.1. *Let f_1, f_2, \dots, f_k be transcendental entire functions. Suppose that, for all $j \in \{1, 2, \dots, k\}$, f_j satisfies Lemma 3.1(a) with $m = m_j > 1$, $m_j \in \mathbb{R}$. Suppose also that, for all $j \in \{1, 2, \dots, k\}$, f_j is ψ -regular, with regularity function ψ_m for each $m > 1$, $m \in \mathbb{R}$. Let $g = f_1 \circ f_2 \circ \dots \circ f_k$. Then $A_R(g)$ is a spider's web, where $R > 0$ is such that $M(r, g) > r$ for $r \geq R$.*

THEOREM 3.2. *Let f_1, f_2, \dots, f_k be entire functions. Suppose that, for all $j \in \{1, 2, \dots, k\}$, f_j satisfies Lemma 3.1(a) with $m = m_j > 1$, $m_j \in \mathbb{R}$. Suppose also that, for some $j \in \{1, 2, \dots, k\}$, f_j is a log-regular transcendental entire function. Let $g = f_1 \circ f_2 \circ \dots \circ f_k$. Then $A_R(g)$ is a spider's web, where $R > 0$ is such that $M(r, g) > r$ for $r \geq R$.*

We need one further lemma before we can prove these results. This lemma also concerns the composition of entire functions.

LEMMA 3.2. *Let f_1, f_2, \dots, f_k be entire functions. Suppose that, for all $j \in \{1, 2, \dots, k\}$, f_j satisfies Lemma 3.1(a) with $m = m_j > 1$, $m_j \in \mathbb{R}$. Let $g = f_1 \circ f_2 \circ \dots \circ f_k$. Then g satisfies Lemma 3.1(a) with $m = m_1 m_2 \dots m_k$.*

Proof. It is sufficient to prove the result for $k = 2$. The result is immediate if f_1 and f_2 are both polynomials. Otherwise, let m_1 and m_2 be as given. Consider first the case that f_2 is a transcendental entire function. For sufficiently large r , let $G_1 = G_1(r)$ be a simply connected domain such that

$$B(0, M(r, f_2)) \subset G_1 \subset B(0, M(r, f_2)^{m_1}), \tag{3.4}$$

and

$$|f_1(z)| \geq M(M(r, f_2), f_1), \quad \text{for } z \in \partial G_1. \tag{3.5}$$

For sufficiently large r , let $G_2 = G_2(r)$ be a simply connected domain such that

$$B(0, r^{m_1}) \subset G_2 \subset B(0, r^{m_1 m_2}), \tag{3.6}$$

and

$$|f_2(z)| \geq M(r^{m_1}, f_2), \quad \text{for } z \in \partial G_2. \tag{3.7}$$

If $f_2(z) \in \partial G_1$ then, by (3.4), $|z| \geq r$, and so there is a component G_3 of $f_2^{-1}(G_1)$ which contains $B(0, r)$. Note that G_3 is simply connected, and f_2 is a proper map of G_3 to G_1 .

If $z \in \partial G_2$ then, by (3.7) and (1.3),

$$|f_2(z)| \geq M(r^{m_1}, f_2) \geq M(r, f_2)^{m_1}, \quad \text{for large } r. \tag{3.8}$$

If $z \in \partial G_3$ then, by (3.4), $|f_2(z)| \leq M(r, f_2)^{m_1}$. Hence, by the maximum principle, if $z \in G_3$ then $|f_2(z)| < M(r, f_2)^{m_1}$. Thus $\partial G_2 \cap G_3 = \emptyset$, by (3.8), and so $B(0, r) \subset G_3 \subset B(0, r^{m_1 m_2})$, by (3.6). Also, if $z \in \partial G_3$ then, by (3.5),

$$|(f_1 \circ f_2)(z)| \geq M(M(r, f_2), f_1) \geq M(r, f_1 \circ f_2). \tag{3.9}$$

Hence $f_1 \circ f_2$ satisfies Lemma 3.1(a), with $m = m_1 m_2$.

Secondly, we consider the case where f_2 is a polynomial. Choose m' such that $m' > m_1$. For sufficiently large r , let G_1 and G_3 be the domains from the first part of the proof, and let $G_2 = B(0, r^{m'})$. Since f_2 is a polynomial, for sufficiently large r ,

$$|f_2(z)| \geq M(r, f_2)^{m'}, \quad \text{for } z \in \partial G_2.$$

As in the first part of the proof, $\partial G_2 \cap G_3 = \emptyset$, and so $B(0, r) \subset G_3 \subset B(0, r^{m'})$. Also, if $z \in \partial G_3$ then $|(f_1 \circ f_2)(z)| \geq M(r, f_1 \circ f_2)$. Hence $f_1 \circ f_2$ satisfies Lemma 3.1(a), with $m = m' > m_1$, in particular with $m = m_1 m_2$.

In particular it follows from Lemma 3.2 that the property of satisfying Lemma 3.1(a) for some $m > 1$, $m \in \mathbb{R}$, is preserved under iteration.

COROLLARY 3.1. *If f is a transcendental entire function that satisfies Lemma 3.1(a) for some $m > 1$, $m \in \mathbb{R}$, then so is f^n for all $n \in \mathbb{N}$.*

We are now able to prove Theorems 3.1 and 3.2.

Proof of Theorem 3.1. By Lemma 3.2, g satisfies Lemma 3.1(a) for some $m > 1$, $m \in \mathbb{R}$. By Theorem 2.2, g is ψ -regular, and so satisfies Lemma 3.1(b) for all $m > 1$, $m \in \mathbb{R}$. The result follows by Lemma 3.1.

Proof of Theorem 3.2. As in the proof of Theorem 3.1, g satisfies Lemma 3.1(a) for some $m > 1$, $m \in \mathbb{R}$. By Theorem 2.3, g is log-regular, and so satisfies Lemma 3.1(b) for all $m > 1$, $m \in \mathbb{R}$. The result follows by Lemma 3.1.

Rippon and Stallard [18] show that there are examples of ψ -regular functions that are not log-regular. Some of these examples have order less than $1/2$, and so satisfy Lemma 3.1(a) for some $m > 1$, $m \in \mathbb{R}$; see Lemma 5.2. This shows that there are situations in which Theorem 3.1 can be applied, but not Theorem 3.2.

Finally, we note that the conditions of Theorem 3.2 are satisfied by many of the examples in [20, section 8], and all the examples in this paper (see Sections 5 and 6).

4. Stability of $A_R(f)$ spiders' webs

Many known dynamical properties of a transcendental entire function f are unstable under relatively small changes in f . For example, the functions $f_1(z) = \exp(-z)$, $f_2(z) = f_1(z) + z + 2\pi i - 1$ and $f_3(z) = f_1(z) + z + 1$ all have very different Fatou sets (see, for

example, [6, section 4]). In this section we prove results which show that, in certain circumstances, $A_R(f)$ spiders' webs can be very stable. The first result concerns composition with polynomials.

THEOREM 4.1. *Suppose that f is a log-regular transcendental entire function which satisfies Lemma 3.1(a) for some $m > 1$, $m \in \mathbb{R}$. Let $P(w, z)$ be a polynomial, which is of degree at least one in w , and let $Q(z)$ be a polynomial of degree at least one.*

Let $g(z) = P(f(Q(z)), z)$. Then $A_R(g)$ is a spider's web, where $R > 0$ is such that $M(r, g) > r$ for $r \geq R$.

Proof. By Corollary 2.6, g is log-regular and so satisfies Lemma 3.1(b) for all $m > 1$, $m \in \mathbb{R}$. Hence we need only prove that g satisfies Lemma 3.1(a) for some $m > 1$, $m \in \mathbb{R}$.

As in the proof of Corollary 2.6, let

$$g(z) = af(Q(z))^{N_1}z^{N_2} + \dots,$$

where N_1 is the highest power of w in $P(w, z)$, and N_2 is the highest power of z corresponding to $f(Q(z))^{N_1}$. By Lemma 3.2, $f \circ Q$ satisfies Lemma 3.1(a). Hence, there is an $m > 1$, $m \in \mathbb{R}$ such that, for sufficiently large r , there is a simply connected domain $G = G(r)$ with $B(0, r^m) \subset G \subset B(0, r^{m^2})$ and

$$|f(Q(z))| \geq M(r^m, f \circ Q), \quad \text{for } z \in \partial G. \tag{4.1}$$

Hence, when $z \in \partial G$, for sufficiently large r ,

$$\begin{aligned} |g(z)| &\geq \frac{1}{2}|a|M(r^m, f \circ Q)^{N_1}r^{N_2} && \text{by (4.1) and (1.1)} \\ &\geq 2|a|M(r, f \circ Q)^{N_1}r^{N_2} && \text{by (1.3)} \\ &\geq M(r, g) && \text{by (1.1)}. \end{aligned}$$

Thus g satisfies Lemma 3.1(a) with m replaced by m^2 , so the proof is complete.

The second result concerns addition of an entire function to a transcendental entire function with an $A_R(f)$ spider's web.

THEOREM 4.2. *Suppose that f is a log-regular transcendental entire function which satisfies Lemma 3.1(a) for some $m > 1$, $m \in \mathbb{R}$, and that h is an entire function. Suppose also that there exist $a \in (0, 1)$ and $R_0 > 0$ such that*

$$aM(r, f) \geq M(r^m, h), \quad \text{for } r \geq R_0. \tag{4.2}$$

Let $g = f + h$. Then $A_R(g)$ is a spider's web, where $R > 0$ is such that $M(r, g) > r$ for $r \geq R$.

Proof. First we note that, for sufficiently large r , $aM(r, f) \geq M(r, h)$. Hence, by Corollary 2.4, g is log-regular and so satisfies Lemma 3.1(b) for all $m > 1$, $m \in \mathbb{R}$. Thus, by Lemma 3.1, it remains to prove that g satisfies Lemma 3.1(a) for some $m > 1$, $m \in \mathbb{R}$.

By hypothesis, for sufficiently large r , there is a simply connected domain $G = G(r)$ with $B(0, r^m) \subset G \subset B(0, r^{m^2})$ and

$$|f(z)| \geq M(r^m, f), \quad \text{for } z \in \partial G. \tag{4.3}$$

Thus, when $z \in \partial G$, for sufficiently large r ,

$$\begin{aligned} |g(z)| &\geq |f(z)| - |h(z)| \\ &\geq (1 - a)M(r^m, f) && \text{by (4.2), and (4.3)} \\ &\geq (1 + a)M(r, f) && \text{by (1.3)} \\ &\geq M(r, g) \end{aligned}$$

Hence g satisfies Lemma 3.1(a) with m replaced by m^2 , so the proof is complete.

Remark. Using the same method of proof it can be shown that in Theorem 4.2 the function h can also be of the form $h(z) = f(z)/(z - c)$, where $f(c) = 0$.

Finally, we note that the conditions on f in Theorems 4.1 and 4.2 are satisfied by many of the examples in [20, section 8], and all the examples in this paper. It can be shown that these conditions are also satisfied by the functions in [16]. So we can produce new functions for which $A_R(f)$ is a spider’s web by taking these known examples and applying Theorems 4.1 and 4.2.

5. Functions of finite order for which $A_R(f)$ is a spider’s web

In this section we develop a technique which enables us to take a transcendental entire function of finite order, modify its power series, and produce a class of transcendental entire functions of finite order for which $A_R(f)$ is a spider’s web. From the exponential function we obtain a class of such functions (Example 1) which contains the function

$$f(z) = \frac{1}{2}(\cos z^{\frac{1}{4}} + \cosh z^{\frac{1}{4}}) = \sum_{n=0}^{\infty} \frac{z^n}{(4n)!} \tag{5.1}$$

given in [20, section 8], together with the related functions

$$f(z) = \sum_{n=0}^{\infty} \frac{z^{pn}}{(qn)!}, \quad p, q \in \mathbb{N}, \quad p/q < \frac{1}{2}, \tag{5.2}$$

suggested by Halburd and also mentioned in [20, section 8]. We obtain another class (Example 3) from the error function (see [1, p. 297])

$$\operatorname{erf}(z) = \frac{2}{\sqrt{\pi}} \sum_{n=0}^{\infty} \frac{(-1)^n}{n!(2n + 1)} z^{2n+1}. \tag{5.3}$$

We define the order $\rho(f)$ and lower order $\lambda(f)$ of a transcendental entire function f by

$$\rho(f) = \limsup_{r \rightarrow \infty} \frac{\log \log M(r, f)}{\log r} \quad \text{and} \quad \lambda(f) = \liminf_{r \rightarrow \infty} \frac{\log \log M(r, f)}{\log r}. \tag{5.4}$$

We note from, for example, [15] that if $f(z) = \sum_{n=0}^{\infty} a_n z^n$, then

$$\rho(f) = \limsup_{n \rightarrow \infty} \frac{n \log n}{\log |a_n|^{-1}} \tag{5.5}$$

and

$$\lambda(f) = \max_{(n_p)} \liminf_{p \rightarrow \infty} \frac{n_p \log n_{p-1}}{\log |a_{n_p}|^{-1}}. \tag{5.6}$$

We use the following three lemmas, all discussed in [20, corollary 8.3 and the following

remarks]. The first is from [14, p. 205], and gives a sufficient condition for a transcendental entire function to be log-regular.

LEMMA 5.1. *If f is a transcendental entire function of finite order and positive lower order, then f is log-regular.*

The second is from, for example, [3, Satz 1].

LEMMA 5.2. *If f is a transcendental entire function of order less than $\frac{1}{2}$, then f satisfies Lemma 2.1(a) for some $m > 1$, $m \in \mathbb{R}$.*

The third follows from Lemma 5.1, Lemma 5.2 and Lemma 2.1.

LEMMA 5.3. *If f is a transcendental entire function of order less than $1/2$ and positive lower order, then $A_R(f)$ is a spider's web, where $R > 0$ is such that $M(r, f) > r$ for $r \geq R$.*

We use the following operator to produce classes of functions which satisfy the conditions of Lemma 5.3.

Definition. For $n, m \in \mathbb{N}$, let $T_{\frac{n}{m}}$ be defined by

$$T_{\frac{n}{m}}(f(z)) = \frac{1}{m} \sum_{k=1}^m f(e^{\frac{2\pi ik}{m}} z^{\frac{n}{m}}), \tag{5.7}$$

where f is an entire function, and we choose a consistent branch of the m th root for each term in the sum.

If f is a transcendental entire function, then the $T_{\frac{n}{m}}$ operator extracts from the power series of f only those terms with exponents which are multiples of m , and these exponents are multiplied by n/m (see (5.8) below). For example, if $f(z) = e^z$, then

$$T_{\frac{2}{3}}(f(z)) = 1 + \frac{z^2}{3!} + \frac{z^4}{6!} + \dots$$

We note in passing that the $T_{\frac{n}{m}}$ operator has some appealing properties; for example, $T_{\frac{1}{n}} \circ T_{\frac{1}{m}} = T_{\frac{1}{nm}}$ and also $T_{\frac{n}{m}}(f(z^m)) = f(z^n)$.

The following result concerns a key property of this operator, namely its effect on the order of a function.

THEOREM 5.1. *If f is a transcendental entire function of order $\rho(f)$ and $n, m \in \mathbb{N}$, then $T_{\frac{n}{m}}(f)$ is a well-defined entire function of order at most $\frac{n}{m}\rho(f)$.*

Proof. First, we consider the action of $T_{\frac{n}{m}}$ on the power series $f(z) = \sum_{l=0}^{\infty} a_l z^l$. Since we have a consistent choice of the m th root, the sum of the complex roots of unity is zero, and with $p = l/m$, we obtain

$$T_{\frac{n}{m}}(f(z)) = \frac{1}{m} \sum_{k=1}^m \sum_{l=0}^{\infty} a_l e^{\frac{2\pi ikl}{m}} z^{\frac{ln}{m}} = \sum_{l=0}^{\infty} a_l z^{\frac{ln}{m}} \sum_{k=1}^m \frac{1}{m} e^{\frac{2\pi ikl}{m}} = \sum_{p=0}^{\infty} a_{pm} z^{pn}. \tag{5.8}$$

Hence the value of $T_{\frac{n}{m}}(f)$ is independent of the choice of the m th root, and this power series has infinite radius of convergence.

We deduce from (5.5), with $k = pm$, that

$$\begin{aligned} \rho(T_{\frac{n}{m}}(f)) &= \limsup_{p \rightarrow \infty} \frac{pn \log pn}{\log |a_{pm}|^{-1}} \\ &\leq \limsup_{k \rightarrow \infty} \frac{(kn/m) \log(kn/m)}{\log |a_k|^{-1}} \\ &= \frac{n}{m} \limsup_{k \rightarrow \infty} \frac{k \log k}{\log |a_k|^{-1}} \\ &= \frac{n}{m} \rho(f), \end{aligned}$$

as required.

We now seek to use this operator, together with Lemma 5.3, to generate transcendental entire functions for which $A_R(f)$ is a spider’s web. It is possible for the function $T_{\frac{n}{m}}(f)$ to be simply a polynomial when f is a transcendental entire function. For example, if $f(z) = z \exp(z^2)$ then $T_{\frac{1}{2}}(f(z)) = 0$, because the power series of f has only odd powers of z which are eliminated by the $T_{\frac{1}{2}}$ operator.

Even if $T_{\frac{n}{m}}(f)$ is transcendental, $T_{\frac{n}{m}}(f)$ may not have positive lower order when f does. For example, if g is a transcendental entire function of order less than 1 and lower order zero, then $f(z) = g(z^2) + z \exp(z^2)$ has both order and lower order 2, but $T_{\frac{1}{2}}(f(z)) = T_{\frac{1}{2}}(g(z^2)) = g(z)$ has order less than 1 and lower order zero, reasoning as in the previous paragraph.

The following lemma gives two sufficient conditions for $T_{\frac{n}{m}}(f)$ to have positive lower order.

LEMMA 5.4. *Let $f(z) = \sum_{p=0}^{\infty} a_p z^p$ be a transcendental entire function, and let $n, m \in \mathbb{N}$.*

(a) *If*

$$\liminf_{p \rightarrow \infty} \frac{p \log p}{\log |a_{pm}|^{-1}} > 0, \tag{5.9}$$

then $T_{\frac{n}{m}}(f)$ has positive lower order.

(b) *If $T_{\frac{n}{m}}(f)$ has positive lower order, and $g(z) = \sum_{p=0}^{\infty} b_p z^p$ is a transcendental entire function with $|b_p| \geq |a_p|$ for p sufficiently large, then $T_{\frac{n}{m}}(g)$ has positive lower order.*

Proof. For part (a) we note, by (5.8) and with $n_p = np$ in (5.6), that

$$\lambda(T_{\frac{n}{m}}(f)) \geq \liminf_{p \rightarrow \infty} \frac{np \log(n(p-1))}{\log |a_{pm}|^{-1}} = n \liminf_{p \rightarrow \infty} \frac{p \log p}{\log |a_{pm}|^{-1}} > 0.$$

Part (b) follows immediately from (5.6).

We now give some explicit examples of classes of functions for which $A_R(f)$ is a spider’s web. The first example includes (5.1) as a special case.

Example 1. Let $f = T_{\frac{n}{m}}(g)$, where $g(z) = \exp(z)$ and where $m > 2n$. Then $A_R(f)$ is a spider’s web, where $R > 0$ is such that $M(r, f) > r$ for $r \geq R$.

Proof. The exponential function has order 1, and satisfies (5.9) for all $m > 1$, $m \in \mathbb{R}$. Thus f has order less than 1/2 by Theorem 5.1, and the result follows by Lemma 5.4(a) and Lemma 5.3.

The second example illustrates the use of both parts of Lemma 5.4. This result can also be justified by the results of Sections 3 and 4.

Example 2. Let $f = T_{\frac{n}{m}}(g)$, where $g(z) = z \exp(z^2) + \exp(z)$, $m > 4n$ and m is odd. Then $A_R(f)$ is a spider's web, where $R > 0$ is such that $M(r, f) > r$ for $r \geq R$.

Proof. The function $z \mapsto z \exp(z^2)$ has order 2, and satisfies (5.9) when m is odd. Thus f has order less than $1/2$ by Theorem 5.1, and the result follows by Lemma 5.4 and Lemma 5.3.

The technique of this section can be applied any transcendental entire function of finite order, provided its power series satisfies (5.9) for some $m \in \mathbb{N}$. We illustrate this with the error function.

Example 3. Let $f = T_{\frac{n}{m}}(g)$, where $g(z) = \operatorname{erf}(z)$, $m > 4n$ and m is odd. Then $A_R(f)$ is a spider's web, where $R > 0$ is such that $M(r, f) > r$ for $r \geq R$.

Proof. By (5.3) and (5.5), g has order 2, and satisfies (5.9) when m is odd. Thus f has order less than $1/2$ by Theorem 5.1, and the result follows by Lemma 5.4(a) and Lemma 5.3.

Our final example combines earlier results to give an unexpectedly simple function with an $A_R(f)$ spider's web.

Example 4. Let $f(z) = \cos z + \cosh z$. Then $A_R(f)$ is a spider's web, where $R > 0$ is such that $M(r, f) > r$ for $r \geq R$.

Proof. This follows from Theorem 3.2 and the function in (5.1).

Our goal in this section has been to produce a class of log-regular transcendental entire functions of order less than $1/2$, which, by Lemmas 5.2 and 2.1, have an $A_R(f)$ spider's web. Finally, we show that this class can be extended by differentiation or integration, thus giving a further method of constructing $A_R(f)$ spiders' webs.

THEOREM 5.2. *Let f be a log-regular transcendental entire function of order less than $1/2$, and let g be the derivative of f or an integral of f . Then $A_R(g)$ is a spider's web, where $R > 0$ is such that $M(r, g) > r$ for $r \geq R$.*

Proof. Since g has the same order as f , and is log-regular by Corollary 2.3, the result follows by Lemmas 5.2 and 2.1.

6. A function of infinite order with gaps for which $A_R(f)$ is a spider's web

We recall that a transcendental entire function f has Fabry gaps if

$$f(z) = \sum_{k=1}^{\infty} a_k z^{n_k}$$

and $n_k/k \rightarrow \infty$ as $k \rightarrow \infty$. By a result of Fuchs [11], an entire function f of finite order with Fabry gaps satisfies Lemma 2.1(a) for $m > 1$, $m \in \mathbb{R}$. This fact was used by Wang in [22, theorem 1] to describe a class of entire functions with no unbounded Fatou components. Thus if f is also log-regular then, by Lemma 2.1, $A_R(f)$ is a spider's web. (As noted earlier, a log-regular transcendental entire function satisfies Lemma 2.1(b) for all $m > 1$, $m \in \mathbb{R}$.) This fact was pointed out by Rippon and Stallard [20, theorem 1.9(d)],

who gave an example of such a function [20, example 1], shown to be log-regular by using Lemma 5.1.

It was also pointed out in [22] and in [20, section 8] that, by a result of Hayman [12], Lemma 2.1(a) holds in the case of certain functions of infinite order with gaps:

LEMMA 6.1. *Let $f(z) = \sum_{k=1}^{\infty} a_k z^{n_k}$ be a transcendental entire function where, for some $\alpha > 2$,*

$$n_k > k \log k(\log \log k)^\alpha, \quad \text{for large } k. \tag{6.1}$$

Then f satisfies Lemma 2.1(a) for $m > 1$, $m \in \mathbb{R}$.

Wang [22, theorem 2] used this result to show that if f satisfies (6.1) and has a property equivalent to log-regularity, then f has no unbounded Fatou components.

Suppose that g is a transcendental entire function of infinite order generated by omitting terms from the power series of another transcendental entire function, f say, and g satisfies (6.1). If g is also log-regular, then $A_R(g)$ is a spider’s web, by Lemma 2.1. If f has infinite order, then it does not seem straightforward to check that such a function g is log-regular. In this section we demonstrate a method for achieving this, and then give an explicit example of such a function.

We start with a general result.

THEOREM 6.1. *Suppose that $f(z) = \sum_{n=0}^{\infty} a_n z^n$ is a log-regular transcendental entire function and there exists $N_0 \in \mathbb{N}$ such that*

$$0 < a_{n+1} \leq a_n, \quad \text{for } n \geq N_0. \tag{6.2}$$

Suppose also that g is a transcendental entire function with

$$g(z) = \sum_{k=1}^{\infty} a_{n_k} z^{n_k}, \tag{6.3}$$

where, for some $M > 1$ and $\alpha > 2$,

$$1 < \frac{n_{k+1}}{n_k} < M, \quad \text{for large } k, \tag{6.4}$$

and

$$n_k > k \log k(\log \log k)^\alpha, \quad \text{for large } k. \tag{6.5}$$

Then g is log-regular and $A_R(g)$ is a spider’s web, where $R > 0$ is such that $M(r, g) > r$ for $r \geq R$.

Proof. By Lemma 6.1, g satisfies Lemma 2.1(a) for $m > 1$, $m \in \mathbb{R}$. To complete the proof, we use Theorem 2.4 to show that g is log-regular.

Without loss of generality, by adding a polynomial, we can assume by Corollary 2.6 that $N_0 = 0$ and (6.4) holds for $k \geq 1$. Because $a_n > 0$ for $n \geq 0$,

$$M(r, f) = f(r) > g(r) = M(r, g), \quad \text{for } r > 0.$$

Thus it remains to show that there exist $a > 1$ and $R_0 > 0$ such that

$$M(r^a, g) \geq M(r, f), \quad \text{for } r \geq R_0. \tag{6.6}$$

Choose $a' > 1$ and $K > 1$ sufficiently large such that

$$\frac{n_{k+1}}{n_k} < \frac{1}{2}(1 + a') < a', \quad \text{and} \quad K^{n_k} > n_{k+1} - n_k, \quad \text{for } k \geq 1. \tag{6.7}$$

Now let $\mu = (a' - 1)/2 > 0$, and define

$$M(r^{a'}, g) = \sum_{k=1}^{\infty} A_k, \quad A_k = a_{n_k} r^{a' n_k}, \tag{6.8}$$

$$M(r, f) = \sum_{n=0}^{a_{n_1}-1} a_n r^n + \sum_{k=1}^{\infty} B_k, \quad B_k = a_{n_k} r^{n_k} + \dots + a_{n_{k+1}-1} r^{n_{k+1}-1}. \tag{6.9}$$

Because the a_n are decreasing,

$$B_k < (n_{k+1} - n_k) a_{n_k} r^{n_{k+1}}, \quad \text{for } r > 1 \text{ and } k \geq 1. \tag{6.10}$$

Thus, if $r > \max\{1, K^{\frac{1}{\mu}}\}$, then, by (6.8) and (6.7),

$$B_k < (n_{k+1} - n_k) r^{n_{k+1}-a' n_k} A_k < K^{n_k} r^{-n_k \mu} A_k < A_k, \quad \text{for } k \geq 1. \tag{6.11}$$

Thus, by (6.8) and (6.9),

$$M(r^{a'}, g) > M(r, f) - \sum_{n=0}^{a_{n_1}-1} a_n r^n, \quad \text{for } r > \max\{1, K^{\frac{1}{\mu}}\}. \tag{6.12}$$

Finally, for any $a > a'$ we can choose r sufficiently large such that

$$M(r^a, g) \geq 2M(r^{a'}, g) \tag{6.13}$$

by (1.2)

$$> 2M(r, f) - 2 \sum_{n=0}^{a_{n_1}-1} a_n r^n \tag{6.14}$$

by (6.12)

$$\geq M(r, f) \tag{6.15}$$

by (1.1).

This proves (6.6) as required.

In the rest of this section we construct an explicit example of a transcendental entire function f of infinite order, defined by a gap series, for which $A_R(f)$ is a spider's web. First we need a simple result about functions of infinite order.

LEMMA 6.2. *Let f and g be transcendental entire functions, and suppose that f has infinite order. If there exist $a, R_0 > 0$ such that*

$$M(r^a, g) \geq M(r, f), \quad \text{for } r \geq R_0,$$

then g has infinite order.

Proof. By (5.4),

$$\rho(g) = \limsup_{r \rightarrow \infty} \frac{\log \log M(r^a, g)}{\log r^a} \geq \frac{1}{a} \limsup_{r \rightarrow \infty} \frac{\log \log M(r, f)}{\log r} = \frac{1}{a} \rho(f),$$

and the result follows.

The next lemma is needed in the construction of our example.

LEMMA 6.3. Let $g(z) = \sum_{n=0}^{\infty} a_n z^n$ be a transcendental entire function, with $a_n \in \mathbb{R}$ for $n \geq 0$, $a_1 \leq 1$, and

$$0 < (n + 1)a_{n+1} \leq na_n, \quad \text{for } n \geq 1. \tag{6-16}$$

Then $f(z) = \exp(g(z))$ has power series $f(z) = \sum_{n=0}^{\infty} b_n z^n$, where

$$0 < b_{n+1} \leq b_n, \quad \text{for } n \geq 1. \tag{6-17}$$

Proof. Clearly $b_n > 0$ for $n \geq 0$. Since $f'(z) = g'(z)f(z)$ we have

$$\sum_{n=0}^{\infty} (n + 1)b_{n+1}z^n = \sum_{k=0}^{\infty} (k + 1)a_{k+1}z^k \sum_{l=0}^{\infty} b_l z^l. \tag{6-18}$$

Equating powers of z gives

$$(n + 1)b_{n+1} = \sum_{l=0}^n (n + 1 - l)a_{n+1-l}b_l, \quad \text{for } n \geq 0. \tag{6-19}$$

Hence, for $n \geq 1$,

$$(n + 1)b_{n+1} = \sum_{l=0}^{n-1} (n + 1 - l)a_{n+1-l}b_l + a_1 b_n \tag{6-20}$$

$$\leq \sum_{l=0}^{n-1} (n - l)a_{n-l}b_l + b_n, \quad \text{by (6-16) and as } a_1 \leq 1 \tag{6-21}$$

$$= nb_n + b_n, \quad \text{by (6-19),} \tag{6-22}$$

which proves that (6-17) holds.

Finally, as promised, we give our explicit example.

THEOREM 6.4. Let

$$f(z) = \exp(e^z - 1) = \sum_{n=0}^{\infty} b_n z^n \quad \text{and} \quad g(z) = \sum_{n=0}^{\infty} b_n z^{n^2}.$$

Then g is a log-regular transcendental entire function of infinite order, and $A_R(g)$ is a spider's web, where $R > 0$ is such that $M(r, g) > r$ for $r \geq R$.

Proof. We can see that f is log-regular because $\phi(t) = \log M(e^t, f) = \exp(e^t) - 1$ and

$$\frac{\phi'(t)}{\phi(t)} > e^t \geq \frac{2}{t}, \quad \text{for } t \geq 1.$$

Conditions (6-4) and (6-5) are satisfied, and the coefficients b_n are decreasing because the function $z \mapsto e^z - 1$ satisfies the conditions of Lemma 6.3. Hence, by Theorem 6-1, g is log-regular and $A_R(g)$ is a spider's web.

Finally, f has infinite order. We see from the proof of Theorem 6-1 that f and g satisfy (6-6). Hence, by Lemma 6-2, g has infinite order.

Clearly this approach can be used with the function f of Theorem 6-2 to give a class of functions with $A_R(f)$ spiders' webs, by suitably selecting terms from the power series of f . We can also use Lemma 6-3 to find other transcendental entire functions which can be manipulated in this way to give further classes of examples.

Remark. We note in passing that, in Theorem 6.2, $b_n = B_n/n!$, where (B_n) are the Bell numbers (see, for example, [5]). Thus, by (6.17), we have

$$B_{n+1} \leq (n + 1)B_n, \quad \text{for } n \geq 1.$$

In fact the more precise estimates

$$2B_n < B_{n+1} < (n + 1)B_n, \quad \text{for } n \geq 2,$$

hold (see [8, corollary 8]). These can be deduced in a straightforward way from the identity

$$B_{n+1} = \sum_{k=0}^n \binom{n}{k} B_k, \quad \text{for } n \geq 0, \tag{6.23}$$

which follows from (6.19).

7. Transcendental entire functions with no unbounded Fatou components

Baker [4] posed the question of whether the Fatou set of a transcendental entire function of sufficiently small growth can have any unbounded components; see the survey article on this question by Hinkkanen [14]. By [20, theorem 1.5(b)], when $A_R(f)$ is a spider's web, $F(f)$ has no unbounded components. Hence all the examples in this paper have no unbounded Fatou components. In this section we give two results on functions with no unbounded Fatou components, which generalise existing results of this type.

Our first class of functions with no unbounded Fatou components consists of functions formed by composition of ψ -regular functions.

THEOREM 7.1. *Let f_1, f_2, \dots, f_k be transcendental entire functions. Suppose that, for all $j \in \{1, 2, \dots, k\}$, f_j satisfies Lemma 3.1(a) with $m = m_j > 1$, $m_j \in \mathbb{R}$. Suppose also that, for all $j \in \{1, 2, \dots, k\}$, f_j is ψ -regular, with regularity function ψ_m for each $m > 1$, $m \in \mathbb{R}$. Let $g = f_1 \circ f_2 \circ \dots \circ f_k$. Then every component of $F(g)$ is bounded.*

Proof. By Theorem 3.1, $A_R(g)$ is a spider's web, and the result follows by [20, theorem 1.5(b)].

To compare Theorem 7.1 to previous results, we need the following lemma, part of [19, theorem 6]. This gives a sufficient condition for a transcendental entire function to be ψ -regular. We note that although, for other reasons, the full statement of [19, theorem 6] supposes order less than $1/2$, finite order is sufficient for the proof of this part of the result.

LEMMA 7.1. *Let f be a transcendental entire function of finite order. Suppose that there exist $n \in \mathbb{N}$ and $q \in (0, 1)$ such that*

$$M(r, f) \geq \exp^{n+1}((\log^n r)^q), \quad \text{for large } r. \tag{7.1}$$

Then f is ψ -regular with regularity function given, for all $m > 1$, $m \in \mathbb{R}$, by

$$\psi_m(r) = \exp^n((\log r)^p), \quad \text{where } pq > 1.$$

The next result now follows from Lemma 7.1 and Theorem 7.1.

COROLLARY 7.1. *Let f_1, f_2, \dots, f_k be transcendental entire functions of finite order which satisfy Lemma 3.1(a) for some $m > 1$, $m \in \mathbb{R}$. Suppose that there exist $n \in \mathbb{N}$*

and $q \in (0, 1)$ such that, for all $j \in \{1, 2, \dots, k\}$,

$$M(r, f_j) \geq \exp^{n+1}((\log^n r)^q), \quad \text{for large } r. \quad (7.2)$$

Let $g = f_1 \circ f_2 \circ \dots \circ f_k$. Then every component of $F(g)$ is bounded.

Rippon and Stallard, in [19, theorem 6], showed that if f is a transcendental entire function of order less than $1/2$, which satisfies (7.1) for some $n \in \mathbb{N}$ and $q \in (0, 1)$, then f has no unbounded Fatou components. By Lemma 5.2 this is included in Corollary 7.1, with $k = 1$.

Corollary 7.1, with $n = 1$, includes a result of Singh in [21, theorem 1]. (We note that the statement of [21, theorem 1] omits the requirement of finite order, but this was assumed in the proof of [21, lemma 1].)

Our second class of functions with no unbounded Fatou components consists of functions formed by composition of entire functions, one of which is log-regular.

THEOREM 7.2. *Let f_1, f_2, \dots, f_k be entire functions. Suppose that, for all $j \in \{1, 2, \dots, k\}$, f_j satisfies Lemma 3.1(a) with $m = m_j > 1$, $m_j \in \mathbb{R}$. Suppose also that, for some $j \in \{1, 2, \dots, k\}$, f_j is a log-regular transcendental entire function. Let $g = f_1 \circ f_2 \circ \dots \circ f_k$. Then every component of $F(g)$ is bounded.*

Proof. By Theorem 3.2, $A_R(g)$ is a spider's web, and the result follows by [20, theorem 1.5(b)].

As noted in Section 2, this result differs from Theorem 7.1 in that only one function in the composition needs to satisfy the regularity condition and be transcendental.

The final result follows from Theorem 7.2 and Lemma 5.2.

COROLLARY 7.2. *Let f_1, f_2, \dots, f_k be transcendental entire functions of order less than $1/2$. Suppose that, for some $j \in \{1, 2, \dots, k\}$, f_j is log-regular. Let $g = f_1 \circ f_2 \circ \dots \circ f_k$. Then every component of $F(g)$ is bounded.*

This corollary generalises a result of Anderson and Hinkkanen in [2, theorem 2], which states that if a log-regular function has order less than $1/2$, then it has no unbounded Fatou components. Anderson and Hinkkanen's result is included in Corollary 7.2 with $k = 1$.

Cao and Wang [9] developed a similar result to Corollary 7.2, concerning composition of transcendental entire functions. They showed that if $g = f_1 \circ f_2 \circ \dots \circ f_k$, where f_1, f_2, \dots, f_k are transcendental entire functions of order less than $1/2$, at least one of which has positive lower order, then g has no unbounded Fatou components. By Lemma 5.1, Cao and Wang's result is included in Corollary 7.2. We note that it is possible to construct a class of log-regular functions of lower order zero and any given finite order, in particular order less than $1/2$. This shows that there are situations in which Corollary 7.2 can be applied but not the result of [9].

Acknowledgements. The author is grateful to Phil Rippon and Gwyneth Stallard for their patient, invaluable help with this paper, and also to the referee for a number of helpful suggestions.

REFERENCES

- [1] M. ABRAMOWITZ and I. A. STEGUN. Eds. *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*. National Bureau of Standards, 1972.
- [2] J. M. ANDERSON and A. HINKKANEN. Unbounded domains of normality. *Proc. Amer. Math. Soc.* **126**, 11 (1998), 3243–3252.

- [3] I. N. BAKER. Zusammensetzungen ganzer Funktionen. *Math. Z.* **69** (1958), 121–163.
- [4] I. N. BAKER. The iteration of polynomials and transcendental entire functions. *J. Austral. Math. Soc. Ser. A* **30**, 4 (1980/81), 483–495.
- [5] E. T. BELL. Exponential numbers. *Amer. Math. Monthly* **41**, 7 (1934), 411–419.
- [6] W. BERGWELER. Iteration of meromorphic functions. *Bull. Amer. Math. Soc. (N.S.)* **29**, 2 (1993), 151–188.
- [7] W. BERGWELER and A. HINKKANEN. On semiconjugation of entire functions. *Math. Proc. Camb. Phil. Soc.* **126**, 3 (1999), 565–574.
- [8] S. BOUROUBI. Bell numbers and Engel's conjecture. *Rostock. Math. Kolloq.* **62** (2007), 61–70.
- [9] C. CAO and Y. WANG. Boundedness of Fatou components of holomorphic maps. *J. Dynam. Differential Equations* **16**, 2 (2004), 377–384.
- [10] A. E. EREMENKO. On the iteration of entire functions. *Dynamical systems and ergodic theory (Warsaw 1986)* **23** (1989), 339–345.
- [11] W. H. J. FUCHS. Proof of a conjecture of G. Pólya concerning gap series. *Illinois J. Math.* **7** (1963), 661–667.
- [12] W. K. HAYMAN. Angular value distribution of power series with gaps. *Proc. London Math. Soc.* (3) **24** (1972), 590–624.
- [13] W. K. HAYMAN. The local growth of power series: a survey of the Wiman–Valiron method. *Canad. Math. Bull.* **17**, 3 (1974), 317–358.
- [14] A. HINKKANEN. Entire functions with bounded Fatou components. In *Transcendental dynamics and complex analysis*, vol. 348 of *London Math. Soc. Lecture Note Ser.* (Cambridge University Press, 2008), pp. 187–216.
- [15] O. P. JUNEJA. On the coefficients of an entire series. *J. Anal. Math.* **24** (1971), 395–401.
- [16] H. MIHALJEVIĆ–BRANDT and J. PETER. Poincaré functions with spiders' webs. *Preprint*, arXiv:1010.0299v1.
- [17] J. W. OSBORNE. The structure of spider's web fast escaping sets. *Preprint*, arXiv:1011.0359v1.
- [18] P. J. RIPPON and G. M. STALLARD. Regularity and fast escaping points of entire functions. In *preparation*.
- [19] P. J. RIPPON and G. M. STALLARD. Functions of small growth with no unbounded Fatou components. *J. Anal. Math.* **108** (2009), 61–86.
- [20] P. J. RIPPON and G. M. STALLARD. Fast escaping points of entire functions. *Preprint*, arXiv:1009.5081v1 (2010).
- [21] A. P. SINGH. Composite entire functions with no unbounded Fatou components. *J. Math. Anal. Appl.* **335**, 2 (2007), 907–914.
- [22] Y. WANG. On the Fatou set of an entire function with gaps. *Tohoku Math. J. (2)* **53**, 1 (2001), 163–170.