

Open Research Online

The Open University's repository of research publications and other research outputs

On questions of Fatou and Eremenko

Journal Item

How to cite:

Rippon, P.J. and Stallard, G.M. (2005). On questions of Fatou and Eremenko. Proceedings of the American Mathematical Society, 133(4) pp. 1119–1126.

For guidance on citations see [FAQs](#).

© [not recorded]

Version: Version of Record

Link(s) to article on publisher's website:

<http://dx.doi.org/doi:10.1090/S0002-9939-04-07805-0>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

ON QUESTIONS OF FATOU AND EREMENKO

P. J. RIPPON AND G. M. STALLARD

(Communicated by Michael Handel)

This paper is dedicated to the memory of Professor Noel Baker

ABSTRACT. Let f be a transcendental entire function and let $I(f)$ be the set of points whose iterates under f tend to infinity. We show that $I(f)$ has at least one unbounded component. In the case that f has a Baker wandering domain, we show that $I(f)$ is a connected unbounded set.

1. INTRODUCTION

Let f be a transcendental entire function and denote by $f^n, n \in \mathbf{N}$, the n th iterate of f . The *Fatou set*, $F(f)$, is defined to be the set of points, $z \in \mathbf{C}$, such that $(f^n)_{n \in \mathbf{N}}$ forms a normal family in some neighbourhood of z . The complement, $J(f)$, of $F(f)$ is called the *Julia set* of f . An introduction to the basic properties of these sets can be found in, for example, [3].

The first results on the iteration of transcendental entire functions were given by Fatou in [8]. There are several general questions in [8], most of which have now been solved. One question that remains open is based on the following observation of Fatou: for the functions $f(z) = z + 1 + e^{-z}$ and $f(z) = h \sin z$ (where $0 < h < 1$), there are infinitely many unbounded curves $\gamma_k, k \in \mathbf{Z}$, with the property that, for $z \in \gamma_k$, we have $f^n(z) \rightarrow \infty$ as $n \rightarrow \infty$. Fatou asked whether this property holds more generally. It is known that this property does hold for certain families of functions (see [6] and [12]), but it is still not known whether this property holds in general.

This question of Fatou concerns the so-called ‘escaping set’

$$I(f) = \{z : f^n(z) \rightarrow \infty \text{ as } n \rightarrow \infty\},$$

which was first studied for a general transcendental entire function f by Eremenko [7]. He proved that

$$(1.1) \quad I(f) \neq \emptyset,$$

$$(1.2) \quad J(f) = \partial I(f),$$

$$(1.3) \quad I(f) \cap J(f) \neq \emptyset,$$

$$(1.4) \quad \overline{I(f)} \text{ has no bounded components.}$$

Received by the editors April 4, 2003 and, in revised form, November 28, 2003.
2000 *Mathematics Subject Classification*. Primary 37F10; Secondary 37F45.

Eremenko conjectured that it may be possible to replace $\overline{I(f)}$ with $I(f)$ in (1.4), a problem that still remains open; see [12] for a proof in the case of exponential maps.

In this paper we prove results that give partial answers to both Fatou's and Eremenko's questions concerning the set $I(f)$. In order to do this we consider the following subset of $I(f)$, which was introduced by Bergweiler and Hinkkanen in [5]:

$$A(f) = \{z : \text{there exists } L \in \mathbf{N} \text{ such that } |f^n(z)| > M(R, f^{n-L}), \text{ for } n > L\}.$$

Here, $M(r, f) = \max_{|z|=r} |f(z)|$ and R can be taken to be any value such that $R > \min_{z \in J(f)} |z|$. Properties (1.1), (1.2) and (1.3) also hold for $A(f)$ (see [5] and the remarks at the end of this paper), and both $A(f)$ and $I(f)$ are completely invariant.

Our main result is the following.

Theorem 1. *Let f be a transcendental entire function. Then each $z_0 \in A(f)$ lies in an unbounded closed connected subset of $A(f)$; in particular, $A(f)$ has no bounded components.*

This goes some way to answering Fatou's question as it shows that $I(f)$ has at least one unbounded component. It also goes some way to answering Eremenko's question as it shows that $\overline{I(f)}$ can be replaced with $A(f)$ in (1.4).

When $F(f)$ has a multiply connected component, we can show much more than this; here, $I(f)$ has *exactly* one unbounded component and no bounded components, showing that Eremenko's conjecture is true in this case.

Theorem 2. *Let f be a transcendental entire function and suppose that $F(f)$ has a multiply connected component. Then*

- (a) $A(f)$ is connected and unbounded and contains the closure of every multiply connected component of $F(f)$;
- (b) $I(f)$ is connected and unbounded.

Baker [1] showed that it is possible for $F(f)$ to have a multiply connected component U . He also showed that $U \subset I(f)$ and, for $n \geq n_0$, the sets $f^n(U)$ lie in bounded multiply connected components U_n of $F(f)$ surrounding 0, with $U_n \rightarrow \infty$; see [2, Theorem 3.1]. For this reason, in [13], we introduced the name *Baker wandering domain* for such a component of $F(f)$.

2. BASIC PROPERTIES OF $A(f)$

In this section we prove a number of basic properties of $A(f)$ including, for completeness, some that were stated but not proved in [5]. Several of these properties are used in the proofs of Theorems 1 and 2. To do this, it is helpful to introduce the following alternative definition of $A(f)$:

$$B(f) = \{z : \text{there exists } L \in \mathbf{N} \text{ such that } f^{n+L}(z) \notin \widetilde{f^n(D)}, \text{ for } n \in \mathbf{N}\},$$

where D is any open disc meeting $J(f)$ and \widetilde{U} denotes the union of U and its bounded complementary components. We prove several basic properties of the set $B(f)$ and then show that $B(f) = A(f)$.

In what follows we often use the property that if G is a bounded region, then

$$(2.1) \quad f(\widetilde{G}) \subset \widetilde{f(G)},$$

which holds because if γ is any simple closed curve in G , then the image under f of the inside of γ lies inside $f(\gamma)$, and hence in $\widetilde{f(G)}$.

Lemma 2.1. *The set $B(f)$ has the following properties:*

- (a) $B(f)$ is independent of D ;
- (b) $B(f)$ is completely invariant;
- (c) $B(f^p) = B(f)$, for $p \in \mathbf{N}$;
- (d) if $g = h^{-1}fh$, where $h(z) = az + b$, $a \neq 0$, then $B(f) = h(B(g))$.

Proof. Properties (a), (b) and (c) follow easily from the blowing-up property of $J(f)$. We prove just property (c).

Let D be an open disc meeting $J(f)$. If $z \in B(f)$, then there exists $L \in \mathbf{N}$ such that

$$f^{n+L}(z) \notin \widetilde{f^n(D)}, \text{ for } n \in \mathbf{N}.$$

By the blowing-up property, $\widetilde{f^K(D)} \supset D$ for all large K . So $\widetilde{f^{n+K}(D)} \supset \widetilde{f^n(D)}$, for $n \in \mathbf{N}$, by (2.1), and hence

$$f^{pn+L+K}(z) \notin \widetilde{f^{pn}(D)}, \text{ for } n \in \mathbf{N}.$$

Since it is possible to choose K such that p divides $L+K$, it follows that $z \in B(f^p)$.

On the other hand, if $z \in B(f^p)$, then there exists $l \in \mathbf{N}$ such that

$$(f^p)^{n+l}(z) \notin \widetilde{(f^p)^n(D)}, \text{ for } n \in \mathbf{N}.$$

Thus, by (2.1),

$$f^{pn+pl-k}(z) \notin \widetilde{f^{pn-k}(D)}, \text{ for } n \in \mathbf{N}, k = 1, 2, \dots, p-1,$$

and so $z \in B(f)$.

Property (d) is true because the corresponding orbits, discs, image sets and Julia sets are congruent to each other under h .

To prove that $B(f) = A(f)$, we use a generalisation of Bohr's theorem [9, page 170]. We write $B(0, R) = \{z : |z| < R\}$ and $\overline{B}(0, R) = \{z : |z| \leq R\}$.

Lemma 2.2. *Let f be entire with a 2-cycle z_1, z_2 (possibly coincident). There is an absolute constant c , $0 < c < 1$, such that if*

$$|z_1| < 2R/5 \text{ and } |z_2| < cM(R/2, f),$$

then

$$f(B(0, R)) \supset \{w : |w| = r'\}, \text{ where } r' \geq cM(R/2, f).$$

Proof. Suppose that f omits in $B(0, R)$ two values w_1 and w_2 such that

$$(2.2) \quad |w_1| = cM(R/2, f) \quad \text{and} \quad |w_2| = 2cM(R/2, f),$$

where c has yet to be chosen. Put

$$\phi(z) = \frac{f(\mu(z)) - w_1}{w_2 - w_1}, \quad \text{where } \mu(z) = \frac{R^2(z + z_1)}{R^2 + \overline{z_1}z}.$$

Then μ maps $B(0, R)$ onto itself and

$$|\phi(0)| = \left| \frac{z_2 - w_1}{w_2 - w_1} \right| \leq \frac{2cM(R/2, f)}{cM(R/2, f)} = 2.$$

Also, ϕ omits 0 and 1 in $B(0, R)$, so Schottky's theorem [9, page 169], applied to the function $\phi(Rt)$, $|t| < 1$, gives

$$\log |\phi(z)| \leq (\log^+ |\phi(0)| + C) \left(\frac{1 + |z|/R}{1 - |z|/R} \right), \quad \text{for } |z| < R,$$

where C is an absolute positive constant. Thus, if $z \in \overline{B}(0, 3R/4)$, then we have $|\phi(z)| \leq A = \exp(7(\log 2 + C))$ and so, by (2.2),

$$\begin{aligned} |f(\mu(z))| &\leq |w_1| + |w_2 - w_1| |\phi(z)| \\ &\leq c(1 + 3A)M(R/2, f) \\ &< M(R/2, f), \end{aligned}$$

provided that $c < (1 + 3A)^{-1}$. But this contradicts the definition of $M(R/2, f)$ because $|z_1| < 2R/5$ implies that $\mu(\overline{B}(0, 3R/4)) \supset \overline{B}(0, R/2)$. Therefore, if $c < (1 + 3A)^{-1}$, then f cannot omit two values satisfying (2.2). The lemma follows.

We also use the following variant of Schwarz's lemma.

Lemma 2.3. *Let f be entire with a 2-cycle z_1, z_2 (possibly coincident), and let C be an arbitrary constant, $0 < C < 1$. If*

$$|z_1| < CR/8 \text{ and } |z_2| < CM(R, f)/8,$$

then

$$M(CR/8, f) \leq CM(R, f).$$

Proof. Let $d(z, z'; G)$ denote the hyperbolic metric in a domain G . In particular,

$$d(z, 0; B(0, R)) = \log \left(\frac{1 + |z|/R}{1 - |z|/R} \right),$$

or, equivalently, $|z|/R = \tanh(\frac{1}{2}d(z, 0; B(0, R)))$. Then, by the contraction property of the hyperbolic metric, we have, for $z \in \overline{B}(0, CR/8)$,

$$\begin{aligned} d(f(z), z_2; B(0, M(R, f))) &\leq d(f(z), z_2; f(B(0, R))) \\ &\leq d(z, z_1; B(0, R)) \\ &\leq d(z, 0; B(0, R)) + d(z_1, 0; B(0, R)) \\ &\leq 2 \log \left(\frac{1 + C/8}{1 - C/8} \right) \leq C. \end{aligned}$$

Hence, for $z \in \overline{B}(0, CR/8)$,

$$\begin{aligned} d(f(z), 0; B(0, M(R, f))) &\leq C + d(z_2, 0; B(0, M(R, f))) \\ &\leq C + \log \left(\frac{1 + C/8}{1 - C/8} \right) \leq 2C, \end{aligned}$$

so

$$|f(z)| \leq M(R, f) \tanh C \leq CM(R, f),$$

as required.

Lemma 2.4. $B(f) = A(f)$.

Proof. Clearly $A(f) \subset B(f)$. To prove the reverse inclusion, we use the fact that f is known to have a 2-cycle z_1, z_2 which must also be a 2-cycle (possibly coincident) of f^n , for each $n \in \mathbf{N}$. We now choose $R > 0$ such that

$$(2.3) \quad B(0, cR/16) \cap J(f) \neq \emptyset,$$

$$(2.4) \quad |z_i| < cR/16, \text{ for } i = 1, 2,$$

and

$$(2.5) \quad |z_i| < cM(R/2, f^n)/8, \text{ for } i = 1, 2, n \in \mathbf{N},$$

where c is the constant in Lemma 2.2. Note that (2.5) can be satisfied because $M(R/2, f^n) \rightarrow \infty$ as $n \rightarrow \infty$, by (2.3) and the blowing-up property of $J(f)$.

Now put $D = B(0, R)$. If $z \in B(f)$, then it follows from (2.3) that there exists $L \in \mathbf{N}$ such that

$$f^{n+L}(z) \notin \widetilde{f^n(D)}, \text{ for } n \in \mathbf{N}.$$

The hypotheses of Lemma 2.2 are satisfied for the functions f^n , $n \in \mathbf{N}$, by (2.4) and (2.5), and so

$$f^n(D) \supset \{w : |w| = r_n\}, \text{ where } r_n \geq cM(R/2, f^n), \text{ for } n \in \mathbf{N}.$$

Thus

$$|f^{n+L}(z)| > cM(R/2, f^n), \text{ for } n \in \mathbf{N}.$$

The hypotheses of Lemma 2.3 are satisfied for the functions f^n , $n \in \mathbf{N}$, with $C = c$ and R replaced by $R/2$ (by (2.4) and (2.5)), and so

$$|f^{n+L}(z)| > cM(R/2, f^n) > M(cR/16, f^n), \text{ for } n \in \mathbf{N}.$$

Thus $z \in A(f)$, by (2.3).

3. PROOF OF THEOREM 1

We first choose $R > 0$ so that

$$(3.1) \quad B(0, R) \cap J(f) \neq \emptyset.$$

Then put

$$D_n = f^n(B(0, R)), \text{ for } n = 0, 1, 2, \dots,$$

and let E_n be the complement of \widetilde{D}_n .

Now suppose that z_0 satisfies

$$f^n(z_0) \in E_n, \text{ for } n \in \mathbf{N}.$$

Let L_n be the component of $f^{-n}(E_n)$ which contains z_0 . Then L_n is closed and also unbounded, since f^n is analytic. Moreover,

$$(3.2) \quad L_n \supset L_{n+1}, \text{ for } n \in \mathbf{N}.$$

Indeed, if $L_{n+1} \setminus L_n \neq \emptyset$, then (because $L_{n+1} \cap L_n \neq \emptyset$) we can choose z' in L_{n+1} so that $f^n(z') \notin E_n$. Hence $f^{n+1}(z') \notin E_{n+1}$, by (2.1), a contradiction.

Now, for $n \in \mathbf{N}$,

$$K_n = L_n \cup \{\infty\}$$

is a closed connected subset of $\hat{\mathbf{C}}$. It follows from (3.2) that $K_n \supset K_{n+1}$, for $n \in \mathbf{N}$, and so

$$K = \bigcap_{n=1}^{\infty} K_n$$

is also a closed connected subset of $\hat{\mathbf{C}}$, which contains z_0 and ∞ . Let Γ be the component of $K \setminus \{\infty\}$ which contains z_0 . Then Γ is closed in \mathbf{C} and unbounded; see [11, page 84].

We claim that $\Gamma \subset A(f)$. For if $z \in \Gamma$, then $f^n(z) \in E_n$, for $n \in \mathbf{N}$, and so

$$(3.3) \quad f^n(z) \notin \widetilde{D}_n = \widetilde{f^n(D_0)}, \text{ for } n \in \mathbf{N}.$$

Thus $z \in B(f)$, by (3.1), so $z \in A(f)$, by Lemma 2.4.

Now suppose that $z' \in A(f) = B(f)$. Then, by (3.1) and the definition of $B(f)$, there is a positive integer $L = L(z')$ such that

$$(3.4) \quad f^{n+L}(z') \notin \widetilde{D}_n, \text{ for } n \in \mathbf{N}.$$

Thus $z'' = f^L(z')$ satisfies

$$f^n(z'') \notin \widetilde{D}_n, \text{ for } n \in \mathbf{N}.$$

By the previous argument, z'' lies in an unbounded closed connected subset, Γ'' say, of $A(f)$. Let Γ' denote the component of $f^{-L}(\Gamma'')$ which contains z' . Then Γ' is closed and also unbounded, since f is analytic. Since $A(f)$ is completely invariant, by Lemma 2.1(b) and Lemma 2.4, we deduce that $\Gamma' \subset A(f)$. Thus the proof of Theorem 1 is complete. (Note that if Γ' is constructed in this way, then by (3.3) each $z \in \Gamma'$ satisfies (3.4).)

4. PROOF OF THEOREM 2

As in the proof of Theorem 1, we choose $R > 0$ so that

$$(4.1) \quad B(0, R) \cap J(f) \neq \emptyset$$

and put

$$D_n = f^n(B(0, R)), \text{ for } n = 0, 1, 2, \dots$$

Now let U_0 be a Baker wandering domain, that is, a multiply connected component of $F(f)$. As shown by Baker [2, Theorem 3.1], the sets $f^n(U_0)$, $n = 1, 2, \dots$, lie in bounded multiply connected components U_n of $F(f)$ and $U_n \rightarrow \infty$ as $n \rightarrow \infty$ in such a way that, for $n \geq N$ say, we have $U_n \subset \widetilde{U_{n+1}}$. We can assume that $D_0 \subset \widetilde{U_N}$.

Now $f^n(\widetilde{U_N}) \subset \widetilde{U_{N+n}}$, by (2.1). Thus

$$(4.2) \quad D_n = f^n(D_0) \subset \widetilde{U_{N+n}}, \text{ for } n \in \mathbf{N}.$$

Now suppose that $z \in \overline{U_N}$. Then $f^{n+1}(z) \in \overline{U_{N+n+1}}$, and so $f^{n+1}(z) \notin \widetilde{U_{N+n}}$, for $n \in \mathbf{N}$. Thus, by (4.1) and (4.2), we deduce that $z \in B(f) = A(f)$. Hence $\overline{U_N} \subset A(f)$, and so $\overline{U_0} \subset A(f)$, by the complete invariance of $A(f)$; see Lemma 2.1(b) and Lemma 2.4.

Thus the closure of any Baker wandering domain lies in $A(f)$. Since any unbounded connected set must meet U_n for large n , we deduce from Theorem 1 that $A(f)$ is connected. This proves part (a).

In [5, Lemma 3] it was proved that $J(f) \subset \overline{A(f)}$. Hence

$$A(f) \cup (I(f) \cap J(f)) \subset \overline{A(f)},$$

and so $A(f) \cup (I(f) \cap J(f))$ is a connected subset of $I(f)$.

If $z \in I(f) \cap F(f)$, then z belongs to a Baker domain or a wandering domain, by the classification of components of $F(f)$; see, for example, [3]. Baker domains are unbounded and so cannot exist in this case as they would meet the Baker wandering domains. Similarly, there can be no unbounded wandering domains and so it remains to consider any bounded wandering domains in $I(f)$. Let V_0 be such a wandering domain and put $V_n = f^n(V_0)$. Since $V_0 \subset I(f)$, it follows that, for $n \geq N_1$ say,

$$V_n \subset \mathbf{C} \setminus \widetilde{U_{m(n)}},$$

where $m(n) \rightarrow \infty$ as $n \rightarrow \infty$. Thus

$$f^n(\overline{V_0}) \subset \overline{V_n} \subset \mathbf{C} \setminus \widetilde{U_{m(n)}}, \text{ for } n \geq N_1,$$

and so $\partial V_0 \subset I(f) \cap J(f)$. Hence V_0 lies in the same component of $I(f)$ as does the connected set $A(f) \cup (I(f) \cap J(f))$. Part (b) now follows.

REMARKS

1. In [5] the proof was given that, for a transcendental entire function f , we have $J(f) \subset \overline{A(f)}$. It follows that $J(f) \subset \partial A(f)$, and it was stated in [5] that $J(f) = \partial A(f)$. We outline a proof of the inclusion $\partial A(f) \subset J(f)$, based on our Theorem 2. If this inclusion is not true, then there must be a closed disc Δ_0 in $F(f)$ which meets both $A(f)$, at z say, and the complement of $A(f)$, at z' say. It follows from Theorem 2 that Δ_0 cannot lie in a multiply connected component of $F(f)$. If Δ_0 lies in a simply connected component of $F(f)$, then it follows from [13, Theorem 3(b)], for example, that there exists C , $0 < C < 1$, such that

$$|f^n(z')| \geq C|f^n(z)|, \text{ for } n \in \mathbf{N}.$$

Thus, since $z \in A(f)$, there exist $R > 0$ and $L \in \mathbf{N}$ such that $J(f) \cap B(0, CR/8) \neq \emptyset$ and

$$|f^{n+L}(z')| \geq C|f^{n+L}(z)| \geq CM(R, f^n) \geq M(CR/8, f^n), \text{ for } n \in \mathbf{N},$$

by Lemma 2.3, together with the justification of (2.5). Hence $z' \in A(f)$, which is a contradiction.

2. We also observe that property (1.3) holds with $I(f)$ replaced by $A(f)$. It follows from Theorem 2 that $A(f) \cap J(f) \neq \emptyset$ if f has a Baker wandering domain. It is also true that $A(f) \cap J(f) \neq \emptyset$ if f has no Baker wandering domain. Indeed, Eremenko's proof of property (1.1) gives a point z_0 such that

$$(4.3) \quad |f^{n+1}(z_0)| \geq \frac{1}{2}M(|f^n(z_0)|, f) \text{ as } n \rightarrow \infty,$$

and also, assuming the absence of Baker wandering domains, such that $z_0 \in J(f)$ [7, Theorems 1 and 2]. Hence $z_0 \in A(f) \cap J(f)$, by [5, Lemma 2]. This result of Eremenko shows that if f has no Baker wandering domains, then $J(f)$ has a dense subset of points z_0 such that (4.3) holds.

3. We do not know an example of a transcendental entire function which has a Baker wandering domain and also a (bounded) simply connected wandering domain,

and it is natural to ask whether this can occur. It is possible for an entire function to have a bounded, simply connected, wandering domain. For example, the function

$$f(z) = 2 - \log 2 + 2z - e^z,$$

discussed by Bergweiler in [4], has a Fatou component containing the point $\log 2 + 2\pi i$, which is simply connected and wandering [4, Section 2]. This component can also be shown to be bounded (by finding a Jordan curve around $\log 2 + 2\pi i$, which is mapped outside itself by f ; see [10]).

4. The reasoning in Remark 1 shows that any simply connected component of $F(f)$ which meets $A(f)$ must lie entirely in $A(f)$. We do not know of a case where such a simply connected component of $F(f)$ exists.

REFERENCES

1. I.N. Baker. An entire function which has wandering domains. *J. Austral. Math. Soc. Ser. A* 22 (1976), 173-176. MR0419759 (54:7777)
2. I.N. Baker. Wandering domains in the iteration of entire functions. *Proc. London Math. Soc.* (3) 49 (1984), 563-576. MR0759304 (86d:58066)
3. W. Bergweiler. Iteration of meromorphic functions. *Bull. Amer. Math. Soc.*, 29 (1993), 151-188. MR1216719 (94c:30033)
4. W. Bergweiler. Invariant domains and singularities. *Math. Proc. Camb. Phil. Soc.*, 117 (1995), 525-532. MR1317494 (96b:30055)
5. W. Bergweiler and A. Hinkkanen. On semiconjugation of entire functions. *Math. Proc. Camb. Phil. Soc.*, 126 (1999), 565-574. MR1684251 (2000c:37057)
6. R.L. Devaney and F. Tangerman. Dynamics of entire functions near the essential singularity. *Ergod. Th. and Dynam. Sys.*, 6 (1986), 489-503. MR0873428 (88e:58057)
7. A.E. Eremenko. On the iteration of entire functions. *Dynamical systems and ergodic theory*, Banach Center Publ. 23 (Polish Scientific Publishers, Warsaw, 1989) 339-345. MR1102727 (92c:30027)
8. P. Fatou. Sur l'itération des fonctions transcendentes entières. *Acta Math.*, 47 (1926), 337-370.
9. W.K Hayman. *Meromorphic functions*. Clarendon Press, Oxford, 1964. MR0164038 (29:1337)
10. M. Kisaka, On the connectivity of Julia sets of transcendental entire functions. *Ergodic Theory Dynam. Systems*, 18 (1998), 189-205. MR1609471 (99a:30033)
11. M.H.A. Newman. *Elements of the topology of plane sets of points*, Cambridge University Press, 1961. MR0132534 (24A:2374)
12. D. Schleicher and J. Zimmer. Escaping points of exponential maps. *J. London Math. Soc.* (2), 67 (2003), 380-400. MR1956142 (2003k:37067)
13. P.J. Rippon and G.M. Stallard. On sets where iterates of a meromorphic function zip towards infinity. *Bull. London Math. Soc.*, 32 (2000), 528-536. MR1767705 (2001g:30019)

DEPARTMENT OF PURE MATHEMATICS, THE OPEN UNIVERSITY, WALTON HALL, MILTON KEYNES MK7 6AA, UNITED KINGDOM

E-mail address: p.j.rippon@open.ac.uk

DEPARTMENT OF PURE MATHEMATICS, THE OPEN UNIVERSITY, WALTON HALL, MILTON KEYNES MK7 6AA, UNITED KINGDOM

E-mail address: g.m.stallard@open.ac.uk