

Open Research Online

The Open University's repository of research publications and other research outputs

Ethiopian community health workers' beliefs and attitudes towards children with autism: Impact of a brief training intervention

Journal Item

How to cite:

Tilahun, Dejene; Fekadu, Abebaw; Tekola, Bethlehem; Araya, Mesfin; Roth, Ilona; Davey, Basiro; Hanlon, Charlotte and Hoekstra, Rosa A. (2019). Ethiopian community health workers' beliefs and attitudes towards children with autism: Impact of a brief training intervention. *Autism*, 23(1) pp. 39–49.

For guidance on citations see [FAQs](#).

© 2017 The Authors



<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Version: Accepted Manuscript

Link(s) to article on publisher's website:

<http://dx.doi.org/doi:10.1177/1362361317730298>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

Ethiopian community health workers' beliefs and attitudes towards children with autism: impact of a brief training intervention

Dejene Tilahun^{1,2}, Abebaw Fekadu^{1,3,4}, Bethlehem Tekola⁵, Mesfin Araya¹, Ilona Roth⁶, Basiro Davey⁶; Charlotte Hanlon^{1,7}, Rosa A. Hoekstra⁵

¹Addis Ababa University, Department of Psychiatry, School of Medicine, College of Health Sciences, Addis Ababa, Ethiopia; ²Jimma University, Department of Health Education and Behavioural Sciences, College of Public Health and Medical Science, Jimma, Ethiopia; ³Centre for Innovative Drug Development and Therapeutic Trials for Africa, College of Health Sciences, Addis Ababa University, Addis Ababa, Ethiopia; ⁴Department of Psychological Medicine, Centre for Affective Disorders, Institute of Psychiatry, Psychology and Neuroscience (IoPPN), King's College London, London, UK; ⁵Department of Psychology, IoPPN, King's College London, London, UK. ⁶The Open University, Department of Life, Health and Chemical Sciences, Milton Keynes, UK; ⁷Department of Health Services and Population Research, IoPPN, King's College London, London, UK.

Corresponding author:

Dr Rosa A. Hoekstra, Department of Psychology, Second Floor Addison House, room AH2.06, Guy's Campus, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London SE1 1UL, UK. Email: Rosa.Hoekstra@kcl.ac.uk; Tel: +44 (0)20 78488079.

Acknowledgements

The authors gratefully acknowledge Addis Ababa University and The Open University for technical support. We are also thankful to the study participants for their voluntary participation.

Funding

The study was funded by Autism Speaks (grant #7770).

This manuscript is in press as:

Tilahun, D., Fekadu, A., Tekola, B., Araya, M., Roth, I., Davey, B.; Hanlon, Hoekstra, R.A. Ethiopian community health workers' beliefs and attitudes towards children with autism: impact of a brief training intervention. *Autism*, in press.

Abstract

There is a severe shortage of services for children with autism in Ethiopia; access to services is further impeded by negative beliefs and stigmatising attitudes towards affected children and their families. To increase access to services, care provision is decentralised through task-shifted care by community health extension workers (HEWs). This study aimed to examine the impact of a brief training (Health Education and Training; HEAT) for Ethiopian rural HEWs and comprised three groups: i) HEWs who completed a basic mental health training module (HEAT group, N=104); ii) HEWs who received enhanced training, comprising basic HEAT as well as video-based training on developmental disorders and a mental health pocket guide (HEAT+ group, N=97); iii) HEWs untrained in mental health (N=108). All participants completed a questionnaire assessing beliefs and social distance towards children with autism. Both the HEAT and HEAT+ group showed fewer negative beliefs and decreased social distance towards children with autism compared to the untrained HEW group, with the HEAT+ group outperforming the HEAT group. However, HEAT+ trained HEWs were less likely to have positive expectations about children with autism than untrained HEWs. These findings have relevance for task-sharing and scale up of autism services in low-resource settings worldwide.

Keywords: Autism Spectrum Disorder; Developmental disabilities; Community Health Workers; Developing Countries; Ethiopia; Stigma; Attitude

Autism and other developmental disorders (DD) are increasingly recognised as conditions associated with long-term burden, disability and cost, requiring global action (Kieling et al., 2011; Patel et al., 2013; WHO, 2001). The global prevalence of autism is estimated to be around 0.6% (Elsabbagh et al., 2012). The prevalence of autism in sub-Saharan Africa is unknown (Elsabbagh et al., 2012; Franz et al., 2017), but prevalence studies of general mental health problems and intellectual disability in low and middle income countries (LMIC) suggest that these problems are at least as prevalent as in high-income countries (Maulik et al., 2011; WHO, 2008). In most LMIC mental health and developmental problems are aggravated by a severe shortage in mental health care provision, resulting in large treatment gaps (Saxena et al., 2007).

Ethiopia is a country located in the horn of Africa with a population of almost 100 million people, half of whom are children. Our recent situational analysis of autism services in Ethiopia (Tekola et al., 2016) showed that diagnostic and educational services for children with autism are largely confined to Ethiopia's capital, Addis Ababa, and therefore unavailable to the majority (85%) of families who live rurally. There are only two child psychiatrists working in Ethiopia; the two schools for children with autism in Addis Ababa are both highly oversubscribed with long waiting lists. The identification, care and

treatment of children with autism are further hindered by stigma, negative stereotypes and misconceptions about the causes of DD and child mental illness (Abera et al., 2015; Tilahun et al., 2016). Supernatural explanatory models, for example, attributing a child's problems to a curse or spirit possession, are common, both in caregivers of children with DD (Tilahun et al., 2016) and in parents of general population school children (Abera et al., 2015). Ethiopian caregivers of children with DD report high levels of stigma; caregivers who reported seeking help from traditional institutions and caregivers who gave a supernatural (as opposed to a biomedical) explanation for their child's condition reported higher experienced stigma (Tilahun et al., 2016). Supernatural explanatory models of mental illness and negative attitudes towards people with mental health problems are also common among nursing staff in Ethiopia (Deribew and Tesfaye, 2005; Abera et al., 2014). Examining beliefs regarding autism among health professionals, studies in Kenya (Gona et al., 2015) and Nigeria (Bakare et al., 2009) indicate that similar negative beliefs and misconceptions about the causes of autism are common in other sub-Saharan African countries. People with religious or supernatural beliefs about causation are likely to have less tolerant attitudes towards people with mental health or developmental problems (Gureje et al., 2006). These views and beliefs may, therefore, hamper identification and successful community rehabilitation of children with autism. Consequently it is likely that

the adult community also includes many individuals whose autism goes without recognition or support.

One of the most promising strategies to address the treatment gap for children with autism and other DD is decentralisation of mental health care provision through task-shifted or task-shared care by non-specialists (Saraceno et al., 2007; Patel et al., 2013). Community health workers, referred to as health extension workers (HEWs) in the Ethiopian health system, have a great potential to be involved in decentralised care provision. To improve the general health care facilities for people living in rural areas, Ethiopia launched a community-based health services extension programme in 2003. Since then 38,000 HEWs completed a one-year training course before starting work in rural Ethiopia. Every rural *kebele* (the smallest administrative unit in Ethiopia, comprising about 5000 inhabitants, half of whom are children) is assigned two HEWs. The HEWs deliver primary community healthcare services consisting of health promotion and prevention packages at their health post. In addition, HEWs are actively engaged in outreach services by visiting the houses of people in their community and training model families and community volunteers (Koblinsky et al., 2010). Evaluation of the initial Ethiopian health services extension programme indicated some notable successes, including a significantly higher proportion of

children vaccinated against communicable diseases and an increase in the use of bed nets to prevent malaria (Assefa et al., 2009). However, there was little impact of the programme in other health areas. More importantly in the context of autism, the initial programme did not include any training on mental health or DD.

To address these concerns, the Federal Ministry of Health of Ethiopia worked with The Open University (UK) to develop the Health Education and Training programme (HEAT; <http://www.open.ac.uk/africa/heat/>) to upgrade the training of existing HEWs. The HEAT curriculum covers topics in child and maternal health, family planning and sexual health, environmental hygiene, communicable and non-communicable diseases. Ten study sessions specifically concern mental health, though child mental health and developmental problems receive only limited attention, with virtually no content on autism spectrum disorders.

About 1,300 HEWs were trained using the HEAT modules in a pilot project across the country at the launch of HEAT in 2012; since then, over 12,700 have completed the upgrading training using HEAT study materials (Personal Communication, Federal Ministry of Health, 2016). Following their HEAT training, we examined the experiences and remaining training needs of 104 HEWs who had just completed the HEAT pilot

(Tilahun et al., 2017). The majority of surveyed HEWs were satisfied with the HEAT training materials on mental health and over two third of HEWs indicated they used these study materials at least once a month in their job. However, 74% of the HEWs also indicated they would benefit from additional training on child mental health and DD. Qualitative interviews with a subsample of HEWs indicated HEWs feel they lack knowledge and skills to adequately support children with DD in their community. In response to these findings, we developed an enhanced version of the HEAT mental health training, called HEAT+. This covered a range of adult and child mental health problems, and included training materials specifically focussing on autism and intellectual disability in children. In the HEAT+ research project we surveyed the experience, knowledge and attitudes of HEWs who were trained using HEAT or HEAT+, and compared their experience, knowledge and skills with HEWs not yet trained in mental health or DD. The aim of this study was therefore to evaluate the impact of the basic and enhanced Health Education and Training (HEAT and HEAT+) materials on the beliefs and attitudes towards children with autism in rural HEWs in Ethiopia.

Methods

Study setting

The study was conducted in the Southern Nations, Nationalities, and Peoples' Region (SNNPR), one of the eleven federal administrative regions of Ethiopia. From the Ethiopian 2007 census (Ethiopian Central Statistical Agency, 2012), SNNPR has an estimated population of around fifteen million people, comprising 45 ethnic groups. The region is predominantly rural; nearly 90% of inhabitants reside in rural areas. Upon the launch of the community-based health services extension programme, 7492 rural HEWs were trained in SNNPR between 2003 and 2009 (WHO, 2010). In 2011, the first cohort of 208 HEWs from SNNPR enrolled in the HEAT training, of which 204 successfully completed the training. In 2013 a new cohort were trained using the enhanced (HEAT+) mental health training materials.

Participants

This cross-sectional study comprised three groups: i) HEWs who completed the original HEAT mental health module (HEAT group; N=104); ii) HEWs trained using the enhanced HEAT mental health module (HEAT+ group; N=97); iii) practicing HEWs not yet enrolled in the upgrading programme, who had not yet received any mental health training (untrained group; N=108). The study participants in the HEAT and HEAT+ groups had all received their training in the study sites Hawassa or Hosanna, two of the four available

training facilities in SNNPR at the time of the study. The HEAT group had completed their training on average 16 months prior to the data collection and had since returned to their local community to work. All HEAT trained HEWs who studied in Hosanna and Hawassa (N=116) were invited to take part in this study; complete data are available for 104 HEWs (participation rate 89.7%). HEWs enrolled in the HEAT+ training had completed the HEAT+ mental health module on average 4 months prior, but were still completing other modules of their upgrading training and had thus not yet returned to their local community. Out of the 104 HEAT+ trained HEWs invited to take part, 97 participated (93.3%). Lastly, a total of 116 HEWs not yet trained in HEAT or HEAT+ were invited to take part, of which 108 participated (93.1%). All participating HEWs were female, following the policy of the Federal Ministry of Health of Ethiopia. Each HEW works alongside one HEW colleague in her local community. Working alongside a colleague who had already completed mental health training may affect a HEWs' knowledge and attitudes about autism; we therefore asked whether a HEW worked alongside a trained or untrained colleague. This information and further demographic data is provided in Table 1.

Training

Two types of educational interventions were provided as briefly described above: i) the basic mental health module (HEAT), and ii) the enhanced HEAT mental health module (HEAT+) of the upgrading programme for Ethiopia's HEWs. The basic HEAT mental health training materials comprise ten sessions, equivalent to two weeks of fulltime study, and include a focus on mental health management, assessment, and mental illness prevention strategies. One session focuses on child development and child mental health, including a discussion of developmental problems. Autism is described only very briefly, in two sentences. The HEAT training was provided through class teaching using printed module materials written in English, following the language policy of the Federal Ministry of Education; most students had access to their personal copy of the training materials (Tilahun et al., 2017).

The HEAT+ training comprised the basic HEAT materials, as well as a DVD and a mental health 'pocket guide'. The DVD includes five short video scenarios modelling an HEW interviewing mothers of children with autism or intellectual disability, demonstrating skills in early detection, supportive counselling and problem solving. The mental health pocket guide provides a brief introduction to mental health with a main focus on detection, mental health first aid and providing support to affected families. The pocket guide includes a

dedicated child section, with substantially more information on autism and intellectual disability than provided in the basic HEAT materials. It outlines the symptoms of intellectual disability and autism; provides guidance on referral; gives advice on how to approach parents and how to talk to a child with a DD; highlights the importance of checking for signs of child abuse and protecting the child from harm; gives advice on what information to provide caregivers of a child with a DD, and suggests strategies to help families support their child's development, managing limitations and challenging behaviours while building on areas of skill and promoting new skills. The pocket guide also advises on how to support integration of children with DD into the community, by giving guidance on how to organise community awareness and school-based awareness meetings. The pocket guide was prepared in English and subsequently translated to Ethiopia's official language Amharic. The scripts for the video scenarios were written in English, and then translated to and filmed in Amharic. In addition to the basic HEAT materials, HEWs in the HEAT+ group viewed the training videos as part of their in-class education, and received a personal copy of the pocket guide. The tutors teaching the HEAT+ training were given copies of all HEAT+ materials a month before they delivered the training to students, allowing them to prepare their lessons.

Both the basic HEAT materials and the HEAT+ materials are open educational resources, freely available online on the Open University's HEAT website (quick link: www.tinyurl.com/heatplus). All HEAT and HEAT+ resources were produced by a team of Ethiopian and UK-based experts in mental health and/or pedagogy, ensuring that the materials were culturally and contextually appropriate. The materials use clear learning outcomes directly linked to the key competencies expected of upgraded HEWs (Federal Democratic Republic of Ethiopia, Ministry of Education, 2016).

Measures

To evaluate the impact of the training, all HEWs consenting to take part in the research evaluation were asked to complete a structured questionnaire, including a written case vignette of a child with autism. The vignette described a mother concerned about her 8-year-old son who shows core symptoms of autism (lack of eye contact, limited speech and play with peers, repetitive activities and difficulty with change). The vignette also highlighted the parental love for their son to avoid any impression of family neglect. Following the presentation of the case vignette, the HEWs were asked to respond to various questions related to autism, including self-rating their general beliefs about autism and their preferred social distance (a measure of what level of intimacy and interaction they would

accept) from children with autism. These questions were adapted from the World Psychiatric Association's programme to reduce stigma and discrimination because of schizophrenia (Stuart and Arboleda-Florez, 2001). A similar questionnaire has been used to assess attitudes towards mental illness in Nigeria (Gureje et al., 2005). Adaptation was carried out through expert consensus meetings to make the questions directly relevant to autism and appropriate for the Ethiopian context. The general beliefs questionnaire comprised ten items including questions assessing positive beliefs and expectations (e.g. 'children with autism can make their parents proud') as well as negative beliefs (e.g. 'children with autism can bring bad luck on the community'). Items were assessed on a 4-point scale ranging from 0 = 'never' to 3 = 'nearly always'. The social distance questionnaire comprised six items, including e.g. 'would you feel ashamed to be seen out on the street taking care of a child with autism'. Item responses were recorded on a 4-point scale ranging from 0 = 'definitely not' to 3 = 'definitely', with higher scores indicating a greater preferred social distance from children with autism, suggesting stronger stigmatising attitudes. The survey was first piloted in five untrained HEWs in the Butajira area of SNNPR. Based on the pilot final revisions were implemented before the data collection began.

Data collection procedure

The questionnaires were administered to HEWs through a face-to-face interview in Amharic by trained and experienced data collectors. This data collection procedure is commonly used in mental health research in Ethiopia and is appropriate for the cultural setting. Data collectors were all women who completed secondary school (10th grade), and were of similar socio-economic status to the participants to limit social desirability bias in the HEWs' responses. Data collectors were trained over five days; the training focused on good interviewing skills, role plays and observed pilot interviews. The data collection was conducted at the training facilities in Hawassa and Hosanna. All study participants provided their written informed consent prior to participation. Ethical approval was obtained from the Institutional Review Board of Addis Ababa University's College of Health Sciences and the Open University's Human Research Ethics Committee.

Data management and analysis

Double data entry using Epidata version 3 (Lauritsen and Bruus, 2003) was employed to reduce the risk of data entry errors. The data were then exported to SPSS version 23 (IBM SPSS Statistics 23) and MPlus version 5.21 (Muthén and Muthén, 2007) for analysis. The factor structure of the adapted questionnaires was assessed in two steps, using exploratory

factor analyses (EFA) and confirmatory factor analyses (CFA) in MPlus. The total sample of 309 participants was randomly split in half. Half of the sample was used to conduct the EFA. The factor structure suggested by the EFA was subsequently tested employing CFA using the second half of the sample. Modification indices were used as guidelines to examine whether the fit of the resulting model could be further improved. The overall fit of the models was evaluated using the Comparative Fit Index (CFI) and the Root Mean Square Error of Approximation (RMSEA). A good model fit is indicated by CFI values above 0.90 and RMSEA below 0.08 (Schermelleh-Engel et al., 2003). Once the best fitting models were identified, further analyses were conducted in SPSS. The internal consistency of the resulting scales was estimated using Cronbach's alpha. Non-parametric Kruskal-Wallis test and subsequent Mann-Whitney U tests were used to examine group differences in a single item. Analyses of covariance were used to examine group differences in quantitative scales. The following four variables were included as covariates in these analyses: HEW's age; length of work experience practising as HEW; educational achievement in high school; and whether the HEW worked alongside a HEAT trained colleague.

Results

Participant characteristics

The sample characteristics for each of the three groups are reported in Table 1. The three groups had a similar distribution of religious affiliations; similar proportions in each group had completed high school to 10th or 12th grade level, and similar proportions worked alongside a HEW colleague who had already completed the upgrading programme. The HEAT group was significantly older and had a significantly longer experience working as an HEW compared to the HEAT+ and untrained groups ($p < .01$). Moreover, the untrained group had a slightly higher average grade in high school than the HEAT trained group ($p < .01$). No other group differences in background characteristics were observed.

Factor analyses

The ten items assessing positive and negative beliefs towards children with autism were examined using EFA. Initial EFA indicated that one item showed little covariance with the other nine items. Item content inspection ('Can improve their language skills with the right help') suggested this item mainly assesses a positive belief in efficacy of an intervention, rather than a positive belief towards children with autism directly. The EFA was re-run including the nine remaining items only. The results indicated a 2-factor structure: one

factor comprising four items assessing positive beliefs and expectations about children with autism, and one factor comprising five negative belief items. This 2-factor model was subsequently fitted to the second half of the data. The modification indices suggested that the model fit could be improved if we would allow a small tweak to our specified model: a cross-loading of one of the positive belief items on the negative belief factor (allowing for some covariation of this item with the negative belief factor). Implementing this modification resulted in a good model fit (CFI= 0.927; RMSEA = 0.064). Next, the six items assessing social distance were subjected to EFA. The EFA suggested a 1-factor structure; testing this structure using CFA in the second half of the data indicated a very good model fit (CFI= 0.987; RMSEA = 0.063); there were no modification indices above the minimum value. The 4-item positive beliefs scale had moderate internal consistency (Cronbach's $\alpha = 0.54$), acceptable when taking into account the low number of items included. The internal consistency of the 5-item negative beliefs scale ($\alpha = 0.67$) and 6-item social distance scale ($\alpha = 0.72$) were adequate.

Impact of training on beliefs and attitudes

Tables 2 and 3 show the item-by-item responses of the participants in relation to positive (Table 2) and negative (Table 3) beliefs and stereotypes towards children with autism, while Table 4 presents the responses related to preferred social distance.

The groups differed significantly in their belief that a child with autism can improve their language skills with the right help ($H(2) = 52.371, p < 0.001$). Subsequent Mann-Whitney U tests indicated that HEAT+ trained ($Z = -6.24, p < 0.001, r = -.44$) and basic HEAT trained ($Z = -6.14, p < 0.001, r = -.42$) HEWs were more likely to believe that children with autism can improve their language skills compared to untrained HEWs (Table 2). The HEAT and HEAT+ groups did not differ from each other ($Z = -.46, p > .05$).

Figure 1 shows the mean scores and distributions of the positive beliefs (Figure 1a), negative beliefs (Figure 1b) and social distance (Figure 1c) scales in each of the three groups. There was a significant group effect on the positive beliefs scale ($F(2, 301) = 4.57, p = .01, \text{partial } \eta^2 = .03$). Planned contrasts using Bonferroni correction for multiple comparisons revealed that the HEAT+ group had significantly lower positive beliefs scores than the untrained group ($p = .01$); none of the other group comparisons were significant. None of the included covariates had a significant effect on positive beliefs (all $p > .05$).

Significant group differences were observed for the negative beliefs scores ($F(2, 301) = 45.91, p < .001, \text{partial } \eta^2 = .23$). This comparison took into account the effect of whether the surveyed HEW had a co-worker trained in mental health ($F(1, 301) = 4.08, p = .04, \text{partial } \eta^2 = .01$); none of the other three covariates was significant (all $p > .05$). Considering all three groups together, HEWs with a HEAT-trained co-worker ($N=19$) reported fewer negative beliefs. Both the HEAT ($p = .004$) and HEAT+ group ($p < .001$) showed fewer negative beliefs towards children with autism than the untrained group. The HEAT+ group in turn displayed fewer negative beliefs than the HEAT group ($p < .001$).

Significant differences were also observed between the groups' social distance scores ($F(2, 301) = 103.14, p < .001, \text{partial } \eta^2 = .41$), taking into account the effect of having a trained or untrained HEW co-worker ($F(1, 301) = 4.99, p = .03, \text{partial } \eta^2 = .02$). None of the other covariates was significant (all $p > .05$). Overall, HEWs with a trained co-worker reported greater social distance. Considering this covariate separately for each group revealed that this effect was only significant in the group of HEWs untrained in mental health ($p = .033$). Taking the effect of this covariate into account, both the HEAT ($p < .001$) and the HEAT+ group ($p < .001$) showed decreased social distance towards children with autism compared to

the untrained group; the HEAT+ group displayed a smaller preferred social distance than the HEAT group ($p=.017$).

Discussion

To our knowledge this is the first study in Africa that provides evidence of the impact of both a basic and enhanced educational intervention on the beliefs and attitudes of community health workers towards children with autism. We find that both educational approaches have a significant impact in improving attitudes and negative beliefs, with more mixed findings regarding positive beliefs about children with autism.

Impact of training on positive beliefs and expectations

Compared to untrained HEWs, both the HEAT+ and the HEAT trained groups of HEWs were more likely to think that children with autism can improve their language skills with the right help. While two thirds of the HEWs not yet trained in mental health and DD thought children with autism will never or rarely improve their language with the right help, only a minority of HEAT (23%) and HEAT+ (16%) trained HEWs had such low expectations of the efficacy of a language intervention.

HEAT+ trained HEWs were less likely to have positive expectations about children with autism, e.g. that they can make their parents proud, can attend school, can get married when they grow up, and can play normally with other children. This unexpected finding may perhaps be explained by the exposure of HEWs in rural Ethiopia to predominantly severe cases of autism, as well as the direction of our training materials towards cases with severe symptoms of developmental disorders. The training materials were developed based on our previous research and clinical experience in Ethiopia. Identified children with autism in Ethiopia nearly always have comorbid intellectual disability (Tilahun et al., 2016), and tend to have a classic form of autism, with no or limited speech, and frequent additional challenging behaviours such as aggression (Hoekstra et al., 2017). To make the training directly relevant to rural Ethiopian context the HEAT+ materials primarily characterised the difficulties of children with classic autism. Though areas of skill were mentioned, the predominant emphasis on problems may have somewhat limited the scope for HEAT+ trained HEWs to formulate more positive expectations of outcomes, especially given the context in which the HEWs work, in rural settings where intervention services are unavailable (Tekola et al., 2016). The limited positive expectations of HEAT+ trained HEWs' may reflect their improved knowledge about autism (compared to the other two groups who received little or no autism-specific training), but also the practical reality that

severely affected children are unlikely to get married later, and unlikely to fit in mainstream education, in the absence of specialised autism education or behavioural intervention services (Tekola et al., 2016). Our finding of no difference between the HEAT and untrained group is consistent with a number of previous mental health education studies that observed no significant change in positive attitudes or beliefs in a positive outcome for mental health patients in community health workers (Armstrong et al., 2011; Makanjuola et al., 2012) and occupational health students (Penny et al., 2001) after training.

Impact of training on negative beliefs

Compared to HEWs untrained in mental health, both HEAT and HEAT+ trained HEWs were less likely to endorse negative beliefs relating to children with autism. The HEAT+ trained group in turn endorsed fewer negative beliefs than the HEAT trained group. HEWs who had a HEAT-trained co-worker (irrespective of group) were also less likely to report negative beliefs. Importantly, HEAT or HEAT+ trained HEWs were unlikely to believe that children with autism ‘need to be chained up at home’, while 18% of untrained HEWs thought this would ‘often’ or ‘nearly always’ be needed. Chaining and beating children with DD is still common in Ethiopia (Tilahun et al., 2016), it is thus encouraging to see that a brief training appears to reduce the belief in this practice.

The effectiveness of the HEAT and HEAT+ programme to change negative beliefs is in keeping with previous mental health education intervention studies in health workers (Li et al., 2014; Li et al., 2015; Liu et al., 2016; Mansouri et al., 2009) occupational health students (Penny et al., 2000), police officers (Pinfold et al., 2003) and young people (Yamaguchi et al., 2011). One explanation of the greater impact of HEAT+ compared to basic HEAT may be the use of videos in the training. Previous mental health intervention education studies suggest that video-based education is more effective in changing attitudes than lecture-only education (Clement et al., 2013; Yamaguchi et al., 2013). An alternative explanation may be that the effect of basic HEAT training is reduced in this study as the training was provided sixteen months prior to data collection (in contrast of HEAT+, which was provided four months prior). Previous studies suggest that positive effects of mental health training may decline over time (Baxter et al., 2001; Yamaguchi et al., 2011).

Impact of training on social distance

Both the HEAT and the HEAT+ group showed reduced preferred social distance towards children with autism compared to the untrained group, suggesting decreased stigmatising attitudes. The HEAT+ group displayed a lower preferred social distance than the HEAT

group. Fifty percent of the HEAT+ HEWs obtained a score of zero on the social distance scale, suggesting they have no need for social distance towards children with autism. The effect of the intervention is in keeping with previous reports that examined changes in the social distance towards people with mental health problems in community health workers (Li et al., 2014; Li et al., 2015) , police officers (Hansson and Markström, 2014) and young people (Yamaguchi et al., 2011). The result are also consistent with previous reports that video-based training is more effective in reducing social distance than lecture-only methods (Clement et al., 2013; Yamaguchi et al., 2013).

Somewhat counterintuitively, HEWs who had a HEAT-trained co-worker reported *greater* preferred social distance. Further analyses exploring the effect of this covariate per group showed that the effect was only significant in the untrained group: HEWs untrained in mental health who had a HEAT-trained co-worker indicated a preference for greater social distance towards children with autism. This finding may suggest that gaining some limited mental health insights through working with a trained colleague, in the absence of personal education on this topic, may increase awareness of some mental health-related issues (for example the importance of personal safety) without providing a full picture of mental health and developmental disorders. It also illustrates the complex trajectory of learning

that HEWs embark upon when they start to learn about autism. There is not a straightforward linear acquisition of knowledge and understanding, but a ‘bumpy’ trajectory in which some attitudes go backwards before they can go forward. It is important to note that only a small subsample of HEWs had a HEAT-trained colleague, therefore these covariate results should be interpreted with care.

Limitations

Our study has some limitations. First, the study only included HEWs from SNNPR; views expressed by our participants may not reflect views of HEWs working in other regions. However, stigma and misconceptions about autism are widespread in sub-Saharan Africa (Bakare et al., 2009; Gona et al., 2016; Ruparelia et al., 2016), suggesting that our findings may apply more widely across Ethiopia and be relevant to settings elsewhere in Africa. Second, although we tried to limit social desirability bias by employing independent data collectors of similar socio-economic status to the participants, it is conceivable that socially desirable responding may have played a role. Against this interpretation, the HEWs who took part in our initial evaluation of HEAT (Tilahun et al., 2017) appeared open to share their opinion on what they did not like about the programme (for example, dissatisfaction that the original HEAT materials were only available in English, rather than Amharic). A

third limitation is that our groups were not perfectly matched: The HEAT group was significantly older and had longer work experience as HEW compared to the other two groups, and the untrained group had a slightly higher mark in high school compared to the HEAT group. Moreover, the time lag in between the survey and the HEAT and HEAT+ training was different: the HEAT group had completed their training sixteen months prior, while the HEAT+ group had completed their training only four months prior. Differences between the HEAT and HEAT+ group may thus also be explained by this time lag difference. Fourth, the HEAT+ materials included both a DVD and a mental health pocket guide. From our current study we cannot deduce whether it was the exposure to video materials or access to the pocket guide or both that generated the impact in the HEAT+ group. Fifth, this study did not include a comparison in beliefs and attitudes pre- and post-training. Sixth, this study did not consider whether the training had a direct impact on children with autism and their families. Our study is limited to self-reported beliefs and attitudes; future studies are warranted to examine whether these self-reported changes in beliefs and attitudes translate into tangible differences for families with children with autism, for example in terms of increased identification of children with autism, and decreased stigma experienced by families. Moreover, it would be useful for future studies to explore in more detail how the training tools were used in practice and which aspects of

the training were most important in affecting change. Lastly, the training evaluated in this study was brief and primarily focused on identification of children with DD, raising awareness in the community, decreasing stigma and promoting community rehabilitation. The two-week long training did not allow for inclusion of comprehensive guidance on the use of social communication intervention strategies to support the child's development. In an ongoing project we are piloting the feasibility and acceptability of a parent skills training programme for caregivers of children with DD; if evaluated positively, this training could help address the lack of available intervention services for children with DD in rural Ethiopia.

Implications

Our finding that a brief training on mental health and DD is associated with decreased negative beliefs and an increased willingness to have contact with children with autism and their parents bodes well for the role community health workers could play in decreasing stigma in the community and in community rehabilitation for children with DD. Apart from health prevention and treatment activities, a main responsibility of Ethiopian HEWs concerns outreach services, including home visits and raising health awareness in their local community. Decreased negative beliefs amongst HEWs may contribute to decreased

stigma and negative beliefs in the community as a whole; decreased social distance from HEWs to families with a child with autism is likely to facilitate community rehabilitation, for example through the use of non-specialist workers to administer psychosocial interventions in the community (Patel et al., 2013).

The mixed effects of the HEAT+ training on positive expectations about children with autism highlights the need for carefully designed educational interventions, taking into account the local practical and sociocultural context. The finding of a stronger effect of HEAT+ compared to HEAT training (with the caveat of the greater time lag in the HEAT group) echoes the emerging finding in mental health education research that different forms of training do not all create the same effect, and video presentations may be an especially powerful training method to reduce stigma, especially in training situations where direct social contact is not feasible (Clement et al., 2012; Yamaguchi et al., 2013).

The HEAT and HEAT+ study materials are open educational resources that are free to be adapted and used elsewhere. Given that similar challenges related to autism and mental health are reported elsewhere, the lessons learned in this study are likely to have relevance for training community health workers in other low-resource settings. A recent review

highlighted the lack of evidence for interventions to reduce mental health-related stigma conducted in LMIC (Thornicroft et al., 2016). Our findings are thus likely to be relevant to addressing stigma relating to a broader range of mental health and developmental conditions too.

Conclusion

This study suggests that a two-week training course on mental health and DD has a significant impact in decreasing community health workers' negative beliefs and social distance towards children with autism and their parents. Training delivered in a widely used Ethiopian language, including video materials and a pocket guide, proved especially effective in decreasing negative attitudes and social distance, though was unsuccessful in increasing positive attitudes. Future research needs to investigate how beliefs in a positive outcome can be preserved while developing an increasing awareness about autism in challenging low-resource settings. The findings presented here are likely to be relevant for task-sharing and scale up of services for children with DD in low-resource settings worldwide.

References

- Abera M, Robbins JM and Tesfaye M (2015) Parents' perception of child and adolescent mental health problems and their choice of treatment option in southwest Ethiopia. *Child and Adolescent Psychiatry and Mental Health* 22(9):40.
- Abera M, Tesfaye M, Belachew T, Hanlon C (2014) Perceived challenges and opportunities arising from integration of mental health into primary care: a cross-sectional survey of primary health care workers in south-west Ethiopia. *BMC Health Services Research* 14:113.
- Assefa A, Degnet A and Andinet D (2009) Impact evaluation of the Ethiopian Health Services Extension Programme. *Journal of Development Effectiveness* 1(4):430-449.
- Armstrong G, Kermode M, Raja S, Suja S, Chandra P and Jorm AF (2011) A mental health training program for community health workers in India: impact on knowledge and attitudes. *International Journal of Mental Health Systems* 5:17.
- Bakare MO, Agomoh AO, Ebigbo PO, Eaton J, Okonkwo KO, Onwukwe JU and Onyeama GM (2009) Etiological explanation, treatability and preventability of childhood autism: a survey of Nigerian healthcare workers' opinion. *Annals of General Psychiatry* 8:6.
- Baxter H, Singh SP, Standen P and Duggan C (2001) The attitudes of 'tomorrow's doctors' towards mental illness and psychiatry: changes during the final undergraduate year. *Medical education* 35:381-383.
- Clement S, Nieuwenhuizen AV, Kassam A, Flach C, Lazarus A, Castro Mde, McCrone P, Norman I and Thornicroft G (2012) Filmed v. live social contact interventions to

- reduce stigma: randomised controlled trial. *The British Journal of Psychiatry* 201:57-64.
- Deribew A and Tesfaye M (2005) Assessment of knowledge, attitude, and practice of nursing staff toward mental health problems in Jimma zone, South West Ethiopia. *Ethiopian Journal of Health Sciences* 15(2):199-206.
- Elsabbagh M, Divan G, Koh YJ KY, Kauchali S, Marcín C, Patel V, Paula C, Wang C, Yasamy M and Fombonne E (2012) Global prevalence of autism and other pervasive developmental disorders. *Autism Research* 5:160-179.
- Ethiopian Central Statistical Agency (2012) 2007 Ethiopian population and housing Census, administrative report, Addis Ababa, Ethiopia. *Ethiopian Central Statistical Authority*.
- Federal Democratic Republic of Ethiopia, Ministry of Education (2016) Occupational standard health extension service, NTFQ Level IV. Addis Ababa: FDRE.
- Franz L, Chambers N, von Isenburg M, de Vries PJ (2017) Autism Spectrum Disorder in sub-Saharan Africa: A comprehensive scoping review. *Autism Research* 10(5):723-749.
- Gona JK, Newton CR, Rimba K, Mapenzi R, Kihara M, Van de Vijver FJR, Abubakar A (2015) Parents' and professionals' perceptions on causes and treatment options for autism spectrum disorders (ASD) in a multicultural context on the Kenyan coast. *PLoS ONE* 10(8): e0132729.
- Gureje O, Lasebikan VO, Ephraim-Oluwanuga O, Olley BO and Kola L (2005) Community study of knowledge of and attitude to mental illness in Nigeria. *British Journal of Psychiatry* 186:436-441.

- Gureje O, Olley BO, Ephraim-Oluwanuga O and Kola L (2006) Do beliefs about causation influence attitudes to mental illness. *World Psychiatry* 5:2.
- Hansson L and Markström U (2014) The effectiveness of an anti-stigma intervention in a basic police officer training programme: a controlled study. *BMC Psychiatry* 14:55.
- Hoekstra, R.A., Girma Bayouh, F., Mihretu, A., Adamu, W., Klasen, H., Hanlon, C. The expression, recognition and reporting of autism symptoms in the Ethiopian context. *16th International Meeting For Autism Research, San Francisco, 10-13 May 2017.*
- IBM Corp. Released 2015. IBM SPSS Statistics for Windows, Version 23.0. Armonk, NY: IBM Corp.
- Kieling C, Baker-Henningham H, Belfer M, Conti G, Ertem I, Omigbodun O, Rohde L, Srinath S, Ulkuer N and Rahman A (2011) Global Mental Health 2: Child and adolescent mental health worldwide: evidence for action. *Lancet* 378(9801):1515-1525
- Koblinsky M, Tain F, Gaym A, Karim A, Carnell M and Tesfaye S (2010) Responding to the challenge-the Ethiopian Health Extension Programme and back up support for maternal health care. *Ethiopian Journal of Health Development* 24(1):105-109.
- Lauritsen JM and Bruus M (2003) Epidata (Version 3). A Comprehensive Tool for Validated Entry and Documentation of Data. *The Epidata Association.*
- Li J, Li J, Huang Y and Thornicroft G (2014) Mental health training program for community mental health staff in Guangzhou, China: effects on knowledge of mental illness and stigma. *International Journal of Mental Health Systems* 8:49.
- Li J, Li J, Thornicroft G, Yang H, Chen W and Huang Y (2015) Training community mental health staff in Guangzhou, China: evaluation of the effect of a new training model. *BMC Psychiatry* 15:263.

- Liu G, Jack H, Piette A, Mangezi W, Machando D, Rwafa C, Goldenberg M, Abas M (2016) Mental health training for health workers in Africa: a systematic review. *Lancet Psychiatry* 3(1):65-76
- Makanjuola V, Doku V, Jenkins R and Gureje O (2012) Impact of a one-week intensive ‘training of trainers’ workshop for community health workers in south-west Nigeria. *Mental Health in Family Medicine* 9:33-38.
- Mansouri N GB, Shariat SV, Bolhari J, Nooraie RY, Rahimi-Movaghar A and Alirezaie N (2009) The change in attitude and knowledge of health care personnel and general population following trainings provided during integration of mental health in Primary Health Care in Iran: a systematic review. *International Journal of Mental Health Systems* 3:15.
- Maulik PK, Mascarenhas MN, Mathers CD, Dua T, Saxena S. (2011) Prevalence of intellectual disability: a meta analysis of population based studies. *Research in Developmental Disability* 32:419–36.
- Muthén, L. K., and Muthén, B. O, (2007) Mplus user’s guide (5th ed.). Los Angeles, CA: Muthén & Muthén.
- Patel V, Kieling C, Maulik P and Divan G (2013) Improving access to care for children with mental disorders: a global perspective. *Archives of Disease in Childhood* 98(5): 323–327.
- Penny NH, Kasar J and Sinay T (2001) Student attitudes toward persons with mental illness: The influence of course work and level I fieldwork. *The American Journal of Occupational Therapy* 55:217-220.
- Ruparelia K, Abubakar A, Badoe E, Bakare M, Visser K, Chugani DC, Chugani HT, Donald KA, Wilmschurst JM and Shih A *et al* (2016) Autism spectrum disorders in

- Africa: current challenges in identification, assessment, and treatment: A report on the international child neurology association meeting on ASD in Africa, Ghana, April 3-5, 2014. *Journal of Child Neurology* 31(8):1018-1026.
- Saraceno B, Ommeren MV, Batniji R, Cohen A, Gureje O, Mahoney J, Sridhar D and Underhill C (2007) Barriers to improvement of mental health services in low-income and middle-income countries. *Lancet* 370(9593):1164-1174.
- Saxena S, Thornicroft G, Knapp M and Whiteford H (2007) Global mental health 2. Resources for mental health: scarcity, inequity, and inefficiency. *Lancet* 370(9590):878-889.
- Schermelleh-Engel K, Moosbrugger H and Muller H (2003) Evaluating the fit of structural equation models: Tests of significance and descriptive goodness-of-fit measures. *Methods of Psychological Research Online* 8:23-74.
- Stuart H and Arboleda-Florez J (2001) Community attitudes toward people with schizophrenia. *The Canadian Journal of Psychiatry* 46:245-252.
- Tekola B, Baheretibeb Y, Roth I, Tilahun D, Fekadu A, Hanlon C and Hoekstra RA (2016) Challenges and opportunities to improve autism services in low-income countries: lessons from a situational analysis in Ethiopia: Policy and system review. *Global Mental Health* 3(e21):1-11.
- Thornicroft G, Mehta N, Clement S, Evans-Lacko S, Doherty M, Rose D, Koschorke M, Shidhaye R, O'Reilly C and Henderson C (2016) Evidence for effective interventions to reduce mental-health-related stigma and discrimination. *Lancet* 387(10023):1123-1132.
- Tilahun D, Hanlon C, Araya M, Davey B, Hoekstra RA and Fekadu A (2017) Training needs and perspectives of community health workers in relation to integrating child

- mental health care into primary health care in a rural setting in sub-Saharan Africa: a mixed methods study. *International Journal of Mental Health Systems* 11:15.
- Tilahun D, Hanlon C, Fekadu A, Tekola B, Baheretibeb Y and Hoekstra RA (2016) Stigma, explanatory models and unmet needs of caregivers of children with developmental disorders in a low-income African country: a cross-sectional facility based survey. *BMC Health Services Research* 16:152.
- Yamaguchi S, Mino Y and Uddin S (2011) Strategies and future attempts to reduce stigmatization and increase awareness of mental health problems among young people: A narrative review of educational interventions. *Psychiatry and Clinical Neurosciences* 65:405-415.
- Yamaguchi S, Wu SI, Biswas M, Yate M, Aoki Y, Barley EA and Thornicroft G (2013) Effects of short-term interventions to reduce mental health-related stigma in university or college students: a systematic review. *The Journal of Nervous and Mental Disease* 201(6):490-503.
- WHO (2001) Mental health: new understanding, new hope. Geneva: World Health Organization.
- WHO (2008) Mental Health Gap Action Program (mhGAP): Scaling up care for mental, neurological and substance use disorders. Geneva: World Health Organization.
- WHO (2010) Human resources for health country profile: Ethiopia, Africa Health Workforce Observatory. Geneva: World Health Organization.

Table 1: Participants' socio demographic characteristics in all three study groups

Characteristic	Untrained	HEAT trained	HEAT+ trained
Time since completing training	N/A	16 months	4 months
Mean age (years)	25.6 ±3.3	27.0±3.5*	25.4 ±3.5
Mean work experience (months)	78.1 ± 19.2	86.9±19.4*	78.7±17.7
Religious affiliation	N (%)	N (%)	N (%)
Muslim	11 (10.2)	13 (12.5)	6 (6.2)
Orthodox Christian	11 (10.2)	19 (18.3)	24 (24.7)
Catholic	3 (2.8)	2 (1.9)	1 (1.0)
Protestant	83 (76.9)	69 (66.3)	65 (67.0)
Other	0 (0)	1 (1.0)	1 (1.0)
HEAT-trained colleague employed in same community	N (%)	N (%)	N (%)
No	102 (94.4)	96 (92.3)	92 (94.8)
Yes	6 (5.6)	8 (7.7)	5 (5.2)
Educational achievement in high school (grade)	2.25 ±0.29	2.13±0.26 ^{\$}	2.21±0.26

Grade they completed at school	N (%)	N (%)	N (%)
Grade 10	73 (67.6)	63 (60.0)	71 (73.2)
Grade 12	35 (32.4)	42 (40.0)	26 (26.8)

* significantly different from other two groups, $p < 0.01$

§ significantly different from untrained group only, $p < 0.01$

Table 2: Positive beliefs and expectations about children with autism in untrained, HEAT trained and HEAT+ trained HEWs

Items	Untrained	HEAT	HEAT+
	N (%)	trained N (%)	trained N (%)
<i>Single item: positive belief in effect language intervention</i>			
Can improve their language skills with the right help			
Never	24 (22.2)	6 (5.7)	4 (4.1)
Rarely	45 (41.7)	18 (17.1)	12 (12.4)
Often	19 (17.6)	26 (24.8)	39 (40.2)
Nearly always	20 (18.5)	54 (51.4)	42 (43.3)
 <i>Positive beliefs scale</i>			
Can make their parents proud			
Never	60 (55.5)	77 (74.1)	84 (86.6)
Rarely	14 (13.0)	20 (19.2)	5 (5.2)
Often	20 (18.5)	5 (4.8)	5 (5.2)
Nearly always	14 (13.0)	2 (1.9)	3 (3.0)

Can attend school

Never	54 (50.0)	39 (37.5)	51 (52.6)
Rarely	38 (35.2)	46 (44.2)	41 (42.3)
Often	8 (7.4)	14 (13.5)	3 (3.1)
Nearly always	8 (7.4)	5 (4.8)	2 (2.0)

Can get married when they grow up

Never	59 (54.6)	37 (35.6)	46 (47.4)
Rarely	38 (35.2)	49 (47.1)	35 (36.1)
Often	7 (6.5)	14 (13.5)	11 (11.3)
Nearly always	4 (3.7)	4 (3.8)	5 (5.2)

Can play normally with other children

Never	65 (60.2)	65 (62.5)	68 (70.1)
Rarely	28 (25.9)	26 (25.0)	25 (25.7)
Often	9 (8.3)	6 (5.8)	2 (2.1)
Nearly always	6 (5.6)	7 (6.7)	2 (2.1)

Table 3: Negative beliefs towards children with autism in untrained, HEAT trained and HEAT+ trained HEWs

Items		Untrained	HEAT	HEAT+
		N (%)	trained N (%)	trained N (%)
Are a public nuisance due to poor hygiene or odd behaviour				
	Never	12 (11.1)	7 (6.7)	37 (38.1)
	Rarely	45 (41.7)	39 (37.5)	29 (29.9)
	Often	26 (24.1)	32 (30.8)	22 (22.7)
	Nearly always	25 (23.1)	26 (25.0)	9 (9.3)
Can bring bad luck on the community				
	Never	24 (22.2)	47 (45.2)	65 (67.0)
	Rarely	39 (36.1)	36 (34.6)	21 (21.6)
	Often	28 (25.9)	4 (3.9)	5 (5.2)
	Nearly always	17 (15.8)	17 (16.3)	6 (6.2)
Can be seen talking to themselves				
	Never	10 (9.3)	7 (6.7)	34 (35.1)
	Rarely	19 (17.6)	45 (43.3)	37 (38.1)

Often	43 (39.8)	29 (27.9)	16 (16.5)
Nearly always	36 (33.3)	23 (22.1)	10 (10.3)
Are dangerous to the public because of violent behaviour			
Never	20 (18.5)	20 (19.2)	46 (47.4)
Rarely	32 (29.6)	40 (38.5)	36 (37.1)
Often	38 (35.2)	23 (22.1)	7 (7.2)
Nearly always	18 (16.7)	21 (20.2)	8 (8.3)
Need to be chained up in the home			
Never	77 (71.3)	102 (98.1)	96 (99.0)
Rarely	11 (10.2)	2 (1.9)	1 (1.0)
Often	9 (8.3)	0 (0)	0 (0)
Nearly always	11 (10.2)	0 (0)	0 (0)

Table 4: Social distance towards children with autism in untrained, HEAT trained and HEAT+ trained HEWs

	Untrained	HEAT trained	HEAT+ trained
Items	N (%)	N (%)	N (%)
Feel afraid to have a conversation with a child with autism			
Definitely not	42 (38.9)	75 (72.1)	87 (89.7)
Probably not	8 (7.4)	7 (6.7)	3 (3.1)
Probably	26 (24.1)	9 (8.7)	4 (4.1)
Definitely	32 (29.6)	13 (12.5)	3 (3.1)
Upset about working with a colleague who is a parent of a child with autism			
Definitely not	55 (50.9)	85 (81.7)	84 (86.6)
Probably not	14 (13.0)	5 (4.8)	7 (7.2)
Probably	18 (16.7)	11 (10.6)	3 (3.1)
Definitely	21 (19.4)	3 (2.9)	3 (3.1)
Able to maintain a friendship with a parent of a child with autism*			
Definitely not	50 (46.3)	19 (18.2)	10 (10.3)
Probably not	13 (12.0)	6 (5.8)	3 (3.1)
Probably	7 (6.5)	11 (10.6)	1 (1.0)

Definitely	38 (35.2)	68 (65.4)	83 (85.6)
------------	-----------	-----------	-----------

Feel upset or disturbed about being alone in a room with a child with autism

Definitely not	39 (36.1)	61 (58.6)	68 (70.1)
----------------	-----------	-----------	-----------

Probably not	10 (9.3)	10 (9.6)	14 (14.4)
--------------	----------	----------	-----------

Probably	25 (23.1)	19 (18.3)	9 (9.3)
----------	-----------	-----------	---------

Definitely	34 (31.5)	14 (13.5)	6 (6.2)
------------	-----------	-----------	---------

Feel ashamed if people knew a child in family has been diagnosed with autism

Definitely not	55 (50.9)	88 (84.6)	87 (89.7)
----------------	-----------	-----------	-----------

Probably not	12 (11.1)	7 (6.7)	4 (4.1)
--------------	-----------	---------	---------

Probably	15 (13.9)	6 (5.8)	3 (3.1)
----------	-----------	---------	---------

Definitely	26 (24.1)	3 (2.9)	3 (3.1)
------------	-----------	---------	---------

Feel ashamed to be seen out on the street taking care of a child with autism

Definitely not	62 (57.4)	96 (92.3)	94 (96.9)
----------------	-----------	-----------	-----------

Probably not	10 (9.3)	8 (7.7)	2 (2.1)
--------------	----------	---------	---------

Probably	8 (7.4)	0 (0)	1 (1.0)
----------	---------	-------	---------

Definitely	28 (25.9)	0 (0)	0 (0)
------------	-----------	-------	-------

Note: * denotes a reverse scored item in the social distance sum score

Figure 1 Positive and negative beliefs and social distance towards children with autism in untrained, HEAT trained, and HEAT+ trained HEWs




Figure 1a




Figure 1b




Figure 1c

Note: a higher score on the positive beliefs (Figure 1a) and negative beliefs (Figure 1b) scales indicates stronger beliefs; a lower social distance score (Figure 1c) indicates less desire to keep a distance from the child with autism suggesting decreased stigmatising attitudes