

# Open Research Online

---

The Open University's repository of research publications and other research outputs

## Role of Asian summer monsoon subsystems in the inter-hemispheric progression of deglaciation

### Journal Item

#### How to cite:

Nilsson-Kerr, K.; Anand, P.; Sexton, P. F.; Leng, M. J.; Misra, S.; Clemens, S. C. and Hammond, S. J. (2019). Role of Asian summer monsoon subsystems in the inter-hemispheric progression of deglaciation. *Nature Geoscience*, 12 pp. 290–295.

For guidance on citations see [FAQs](#).

© 2019 The Authors, under exclusive licence to Springer Nature Limited



<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Version: Accepted Manuscript

Link(s) to article on publisher's website:  
<http://dx.doi.org/doi:10.1038/s41561-019-0319-5>

---

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

---

[oro.open.ac.uk](http://oro.open.ac.uk)

1 **Role of Asian summer monsoon subsystems in the inter-hemispheric progression of**  
2 **deglaciation**

3 K. Nilsson-Kerr<sup>1\*</sup>, P. Anand<sup>1</sup>, P. F. Sexton<sup>1</sup>, M. J. Leng<sup>2</sup>, S. Misra<sup>3</sup>, S. C. Clemens<sup>4</sup>, S. J.  
4 Hammond<sup>1</sup>

5 <sup>1</sup>School of Environment, Earth and Ecosystem Sciences, Faculty of Science, Technology,  
6 Engineering and Mathematics, The Open University, Milton Keynes, MK76AA, UK

7 <sup>2</sup>NERC Isotope Geoscience Facilities, British Geological Survey, Nottingham, NH125GG, UK  
8 and Centre for Environmental Geochemistry, School of Biosciences, Sutton Bonington  
9 Campus, University of Nottingham, Loughborough, LE12 5RD, UK

10 <sup>3</sup>Centre for Earth Sciences, Indian Institute of Science, Bangalore, 560012, India

11 <sup>4</sup>Earth, Environmental, and Planetary Sciences, Brown University, Providence, 02912, USA

12 (\*katrina.kerr@open.ac.uk)

13 **The response of Asian Monsoon subsystems to both hemispheric climate forcing and**  
14 **external orbital forcing are currently issues of vigorous debate. The Indian Summer**  
15 **Monsoon is the dominant monsoon subsystem in terms of energy flux, constituting one of**  
16 **Earth's most dynamic expressions of ocean-atmosphere interactions. Yet the Indian**  
17 **Summer Monsoon is grossly under-represented in Asian Monsoon palaeoclimate records.**  
18 **Here we present high-resolution records of Indian Summer Monsoon induced rainfall**  
19 **and fluvial runoff recovered in a sediment core from the Bay of Bengal across**  
20 **Termination II, 139 to 127 thousand years ago, including coupled measurements of the**  
21 **oxygen isotopic composition and Mg/Ca, Mn/Ca, Nd/Ca and U/Ca ratios in surface-ocean**  
22 **dwelling foraminifera. Our data reveal a millennial-scale transient strengthening of the**  
23 **Asian Monsoon that punctuates Termination II associated with an oscillation of the**

24 **bipolar seesaw. The progression of deglacial warming across Termination II emerges first**  
25 **in the southern hemisphere then the tropics in tandem with Indian Summer Monsoon**  
26 **strengthening and finally the northern hemisphere. We therefore suggest that the Indian**  
27 **Summer Monsoon was a conduit for conveying southern hemisphere latent heat**  
28 **northwards, thereby promoting subsequent northern hemisphere deglaciation.**

29 Early modelling studies that attempted to evaluate the response of the boreal summer monsoon  
30 to orbital forcing identified Northern Hemisphere (NH) solar insolation (during precession<sub>min</sub>)  
31 as a primary driver, via its influence on land-ocean thermal contrasts<sup>1</sup>. Palaeoclimate records  
32 of the ISM support this view but also commonly invoke NH climate controls<sup>2</sup> owing to the  
33 coincidence of weak Indian Summer Monsoon (ISM) intervals with North Atlantic Heinrich  
34 Events<sup>2, 3</sup>. These millennial scale cooling events originating in the high latitudes of the NH  
35 have been linked to the ISM via atmospheric<sup>3</sup> and oceanic<sup>4</sup> teleconnections. Similarly, East  
36 Asian Summer Monsoon (EASM) speleothem oxygen isotope ( $\delta^{18}\text{O}$ ) records, inferred to reflect  
37 both upstream depletion of  $\delta^{18}\text{O}$  from tropical moisture sources and regional precipitation  
38 amount<sup>5</sup>, have been linked to both NH solar insolation and North Atlantic forcing<sup>6</sup>, although  
39 this interpretation has been recently questioned in light of new EASM rainfall records<sup>7, 8</sup>.  
40 Despite this prevailing view of NH forcing of the Asian Monsoon on millennial to orbital  
41 timescales, some observations from ISM records have pointed to additional mechanisms  
42 influencing ISM behaviour<sup>9-11</sup>. The nature of variance in the obliquity band and lag of ISM  
43 maxima with precession<sub>min</sub> suggests a component of Southern Hemisphere (SH) forcing  
44 through latent heat export<sup>9, 10</sup>. Understanding of the ISM at timescales beyond the last glacial  
45 period mainly derives from orbital-scale records from the Arabian Sea and southern Bay of  
46 Bengal (BoB) (Fig. 1a). Records from these locations have applied proxies that have been  
47 assumed to be representative of upwelling and changes in water column stratification driven  
48 by ISM winds. However, the extent to which the ISM exclusively controls these proxies

49 remains unclear. Thus, what is urgently required to enhance our understanding of the ISM are  
50 records of rainfall and runoff from the ISM's core convective region, the northern BoB in order  
51 to isolate a primary and direct signal of ISM strength.

52 Here we report new geochemical records from well preserved planktic foraminifera at a sub-  
53 millennial scale resolution (~250-500 years) spanning Termination II (TII, 139 to 127 thousand  
54 years ago (ka)) from IODP 353, Site U1446 in the northern BoB. Site U1446 is situated in the  
55 core convective region of the ISM, under the direct influence of ISM-induced rainfall and  
56 fluvial runoff received from one of the world's largest river systems (Ganges-Brahmaputra).  
57 Figure 1(b-e) shows the southward propagation of the ISM induced freshwater plume derived  
58 from the Ganges-Brahmaputra systems, engulfing Site U1446 during the peak summer  
59 monsoon season. This site is thus ideally situated to capture the signal of ISM derived rainfall,  
60 fluvial runoff and sediment delivery from the Indian subcontinent. We have produced a detailed  
61 stratigraphy for Site U1446 that is tied to the Antarctic Ice Core (AICC2012) chronology<sup>12</sup>  
62 (Methods, Supplementary Fig. 1). To evaluate changes in the surface ocean salinity response  
63 to rainfall and runoff, we combine oxygen isotope ( $\delta^{18}\text{O}_c$ ) and Mg/Ca-derived SSTs from the  
64 planktic foraminifera *Globigerinoides ruber (sensu-stricto)* to reconstruct  $\delta^{18}\text{O}$  of seawater  
65 ( $\delta^{18}\text{O}_{\text{sw}}$ ) (Methods) (Fig. 2B n).

66 We also present Mn/Ca, Nd/Ca and U/Ca ratios (Supplementary Fig. 8) of *G. ruber ss* calcite  
67 in a novel application to reconstruct fluvial runoff, where high concentrations of Mn, Nd and  
68 U are delivered from the continental hinterland by the ISM's vigorous hydrological and  
69 weathering regime<sup>13-15</sup>. This regime exerts a strong seasonal bias on the vertical and lateral  
70 distribution of dissolved 'lithogenic' elements within the BoB<sup>14</sup>, with a strong lithogenic signal  
71 existing in the upper 100m of the northern BoB as a result of high terrigenous fluxes<sup>15</sup>. The  
72 origin of Nd in planktic foraminiferal calcite remains controversial with the Nd being attributed  
73 to either reflect in-situ seawater Nd signal<sup>16</sup>, a mixed signal from sediments and bottom

74 waters<sup>17</sup> or to arise from intra-test organic matter<sup>18</sup>. We interpret our foraminiferal Mn/Ca,  
75 Nd/Ca and U/Ca data to reflect a primary signal of upper ocean chemistry modulated by high  
76 fluxes of lithogenic elements from high fluvial runoff for several reasons. First, the  
77 foraminifera cleaning method we applied included a reductive cleaning step that ensures  
78 removal of Fe-Mn coatings added on the foraminifera test at the sediment-water interface<sup>19</sup>.  
79 Second, Mn/Ca correlates with Nd/Ca and U/Ca (Supplementary Fig. 7), suggesting that the  
80 concentrations of these elements are all derived from the same dominant process (i.e. in this  
81 hydrographic setting, fluvial runoff). Third, the concentrations of lithogenic elements in  
82 modern seawater in the northern BoB are much higher than for global average seawater<sup>15</sup>  
83 (owing to high dissolved elemental fluxes from the continent, driven by the ISM). Fourth, the  
84 observed concentrations of these elements are beyond what is typically found in planktic  
85 foraminifera<sup>20</sup>. We normalised Mn/Ca, Nd/Ca and U/Ca to unit variance<sup>21</sup> to produce a stack  
86 of *G. ruber ss* geochemical tracers of fluvial runoff (Fig. 2B m) (Methods). The range of values  
87 exhibited by this runoff tracers record overlaps with the range of these same elements in  
88 modern *G. ruber ss* as measured from a 2005 sediment trap in the northern BoB (red vertical  
89 bar in Fig. 2B m). This underscores that our *G. ruber ss*-based stacked record of Mn, Nd and  
90 U concentrations is recording high concentrations of these elements in local seawater (derived  
91 from high runoff fluxes), rather than being a post-depositional phenomenon via diagenetic  
92 alteration of the foraminiferal calcite. Therefore, comparing *G. ruber ss*  $\delta^{18}\text{O}_{\text{sw}}$  and *G. ruber ss*  
93 runoff tracers together provides a novel opportunity to reconstruct changes in both salinity and  
94 fluvial runoff sourced directly from the ISM. Application of these runoff tracers in *G. ruber ss*  
95 as representing ISM river fluxes is supported by elemental signatures of continental origin from  
96 discrete portable X-Ray Fluorescence (pXRF) measurements on bulk sediment samples that  
97 are purely diagnostic of continental detrital input from runoff (Al, Ti, K, Rb) (Fig. 2B l)  
98 (Methods, Supplementary Fig. 9).

99 Our high-resolution time series of  $\delta^{18}\text{O}_{\text{sw}}$ , *G. ruber ss* runoff tracers and pXRF element stack  
100 show a similar pattern of ISM behaviour across TII, accounting for the differing intensity in  
101 the response and thresholds between surface freshening and riverine sediment fluxes<sup>22</sup>. The  
102 data reveal a brief intensification of the ISM from ~134 to 133 ka, reflected as a decrease in  
103  $\delta^{18}\text{O}_{\text{sw}}$  (Fig. 2B n), an increase in *G. ruber ss* runoff tracers (Fig. 2B m), and pXRF element  
104 stack (Fig. 2B l) late in Marine Isotope Stage (MIS) 6, prior to TII onset. This was immediately  
105 preceded by a ~1 kyr duration SST warming in the BoB (Fig. 2B o), suggesting advection of  
106 SH heat across the equator provided a crucial precondition<sup>23</sup> for the subsequent transient  
107 strengthening of monsoonal circulation at 134 ka. Our data show that the ISM then undergoes  
108 two phases of deglacial strengthening; first at ~131 to 130 ka, followed by a further  
109 strengthening at ~129 ka, with the final attainment of a vigorous interglacial ISM coeval with  
110 the development of full deglaciation into the Last Interglacial (MIS 5e) (Fig. 2B).

### 111 **Interstadial within Termination II**

112 The structure of the last two terminations, TI and TII, is fundamentally different (Fig. 2,  
113 Methods). TI is punctuated by several millennial scale events, manifested in the Bølling-  
114 Allerød and Younger Dryas, associated with fluctuations in Atlantic Meridional Overturning  
115 Circulation (AMOC)<sup>24</sup> (Fig. 2A). Such millennial scale events have remained largely  
116 unidentified in reconstructions of TII. However, we identify a climatic event punctuating TII,  
117 evident in ISM rainfall and runoff (Fig. 2B l, m and n) at ~134 to 133 ka, prior to the timing of  
118 TII deglaciation in the NH<sup>25</sup>. We refer to this event as the Termination II Interstadial (TII IS).  
119 ISM strengthening during the TII IS was preceded by a 1°C warming in *G. ruber ss* derived  
120 SSTs at ~135 ka (Fig. 2B o). This warming coincides with early deglaciation in the SH (Fig.  
121 2B i, k) but with the establishment of cool condition in the North Atlantic associated with  
122 Heinrich Stadial 11 (HS11) onset<sup>26</sup>. We infer that this SST warming in the BoB reflects cross-  
123 equatorial heat transport in response to contemporaneous warming in the SH. These SH-

124 derived energy fluxes, advecting northwards, leads to the transient strengthening of the ISM  
125 that marks the TII IS (Fig. 2B l, m and n). We thus attribute the TII IS to a transient oscillation  
126 of the bipolar seesaw, akin to mechanisms proposed for TI<sup>24, 27</sup>. The TII IS is also depicted in  
127 other NH records, a western Mediterranean Sea SST record<sup>26, 28</sup> (Fig. 2B f) and the EASM  
128 speleothem  $\delta^{18}\text{O}$  record<sup>6, 29</sup> (Fig. 2B e). Further support for a cross-equatorial northward flux  
129 of SH-derived heat through a bipolar seesaw mechanism is provided by a cooling in the South-  
130 East Atlantic coeval with the TII IS, which has been attributed to a reduction in Agulhas  
131 Leakage associated with a northward shift of the atmospheric belts towards the warmer  
132 (northern) hemisphere<sup>30</sup>. The timing of TII IS is within error of Meltwater Pulse 2B (MWP 2B,  
133  $133\pm 1$  ka)<sup>26</sup>. Thus, it appears that TII IS may have contributed to rapid retreat of NH ice sheets  
134 and the resulting MWP 2B owing to heat import into the NH. The resulting enhanced  
135 freshwater fluxes into the North Atlantic<sup>31</sup> causes an intensification of HS11 (Fig. 2B c, d),  
136 cooling of the NH and the ending of TII IS associated with a southward shift of the Inter-  
137 Tropical Convergence Zone (ITCZ)<sup>32</sup>. Recent work has argued for a robust North Atlantic  
138 control on the EASM<sup>6, 29</sup>. Yet our findings for a SH origin for the transient EASM strengthening  
139 during TII IS, perhaps via the ISM, reveal that the nature of these inter-hemispheric controls  
140 on a given monsoonal subsystem is not fixed, but dynamic across different timescales.

### 141 **Inter-hemispheric progression of deglaciation**

142 The nature of deglaciation during TII is thought to be a result of orbital preconditioning; an  
143 earlier maximum in SH solar insolation 10 kyr prior to NH solar insolation maxima promoting  
144 earlier Antarctic warming<sup>25</sup> (Fig 2B b). Furthermore, obliquity<sub>max</sub> (Fig. 2B a) was reached prior  
145 to precession<sub>min</sub><sup>33</sup> (Fig. 2B b), triggering an increased inter-hemispheric temperature contrast  
146 and strengthening of the Hadley Cell in the warmer (southern) hemisphere. The colder  
147 (northern) hemisphere is compensated by increased cross-equatorial heat transport<sup>34</sup>. Figure 3  
148 shows the statistically determined timings<sup>35</sup> of regional deglaciation throughout TII. The

149 combination of obliquity<sub>max</sub> and early deglacial SH warming (Fig. 3h) dictates that heat and  
150 moisture transported to the ISM would have been across the equator from the southern Indian  
151 Ocean. We thus conclude that SH sourced energy fluxes (Fig. 3h) were responsible for early  
152 deglacial strengthening of the ISM at ~131 to 130 ka (Fig. 3e-g). A contemporaneous early  
153 deglacial warming occurs in the western Mediterranean<sup>26, 28</sup> (Fig. 3d) that we infer reflects  
154 adiabatic descent from the descending limb of the Hadley Cell<sup>36</sup>, propagating SH sourced  
155 energy fluxes northward. This northward propagation of SH heat and moisture into higher NH  
156 latitudes was slowed by the persistence of a cold North Atlantic with HS11 (Fig. 3a, b).  
157 Subsequently, the inter-hemispheric progression of deglacial warming is propagated into the  
158 higher latitudes of the North Atlantic (Fig. 3a, b) with associated EASM strengthening (Fig.  
159 3c). Our ISM records (Fig. 2B m, n) show strong covariance with the Antarctic CH<sub>4</sub> record  
160 (Fig. 2B j) during both the TII IS and broader deglaciation. This finding supports hypotheses  
161 that call for tropical wetlands as being an important global methane source during glacial-  
162 interglacial transitions and that the tropical monsoonal system plays a fundamental role in  
163 regulating concentrations of this greenhouse gas<sup>37</sup>.

#### 164 **Millennial-scale phasing of Asian Monsoon subsystems**

165 Our ISM records across TII provide insights into the relationship between the two main Asian  
166 Monsoon subsystems at the millennial scale. Deglacial ISM strengthening is temporally  
167 decoupled from EASM strengthening by ~1 to 2 kyr (Fig. 3). We infer that this lag is not  
168 associated with respective age-models and instead ultimately reflects the time transgressive  
169 nature of deglacial strengthening in the Asian Monsoon subsystems and influence of differing  
170 forcing mechanisms triggering this strengthening. The makeup of these two monsoonal  
171 subsystems is quite different; differing land-ocean configurations, atmospheric and ocean  
172 dynamics<sup>38</sup> thus, it is likely that during major changes in background climate state the ISM and  
173 EASM exhibit such time-transgressive responses.



174 Our findings thus allow us to reject the hypothesis of a singular common (NH) forcing  
175 mechanism of the Asian Monsoon<sup>6</sup>. Therefore, despite the iconic nature of the EASM  
176 speleothem records<sup>6</sup>, our high-resolution ISM rainfall and runoff data suggest that the  
177 assumption that they are representative of the Asian Monsoon as a whole needs to be  
178 reconsidered, at least on millennial timescales. This decoupling of the ISM and EASM across  
179 TII may owe its origins to the complexities and large-scale variation in the moisture supply  
180 amalgamated in the speleothem  $\delta^{18}\text{O}$  signal<sup>8, 39</sup>. Our new records point to a greater dynamism  
181 in the mechanisms regulating Asian Monsoon rainfall beyond just teleconnections to the North  
182 Atlantic<sup>6</sup>. This emphasises the need for more high-resolution palaeoclimate time series that are  
183 directly influenced by monsoonal rainfall, for both the EASM and ISM, in order to shed further  
184 light on the mechanism and feedbacks regulating monsoonal subsystems.

185 Our findings from TII indicate that the ISM is a key inter-hemispheric link in the transfer of  
186 heat and moisture between the warm SH into the colder NH (Fig. 3). Our sub-millennial scale  
187 records provide support for hypotheses that argue for an important role of the tropics<sup>40</sup> in  
188 conveying SH latent heat northwards into the NH, thereby promoting NH deglaciation.  
189 However, the evolution of the ISM captured in our data suggests that a fully strengthened  
190 ‘interglacial’ mode of the ISM cannot be attained until the NH experiences full deglacial  
191 climatic amelioration (Fig. 3). Our results highlight the need for explicit differentiation  
192 between the ISM and EASM owing to their respective sensitivities to fundamentally different  
193 components of the Earth system during global climate change. Our data also reveal that inter-  
194 hemispheric climatic controls on the two primary monsoonal subsystems are dynamic across  
195 different timescales and that, during a glacial transition, these two monsoonal subsystems can  
196 be governed by different inter-hemispheric controls.

## 197 **References**

- 198 1. Kutzbach, J. E. Monsoon climate of the early Holocene: climate experiment with the  
199 Earth's orbital parameters for 9000 years ago. *Science*. **214**, 59-61 (1981).
- 200 2. Kathayat, G. *et al.* Indian monsoon variability on millennial-orbital timescales. *Sci.*  
201 *Rep.* **6**, 24374 (2016).
- 202 3. Deplazes, G. *et al.* Weakening and strengthening of the Indian monsoon during  
203 Heinrich events and Dansgaard-Oeschger oscillations. *Paleoceanography*. **29**,  
204 2013PA002509 (2014).
- 205 4. Tierney, J. E., Pausata, F, S, R. & deMenocal, P. Deglacial Indian monsoon failure and  
206 North Atlantic stadials linked by Indian Ocean surface cooling. *Nat. Geosci.* **9**, 46-50  
207 (2015).
- 208 5. Orland, I. J. *et al.* Direct measurements of deglacial monsoon strength in a Chinese  
209 stalagmite. *Geology*. **43**, 555-558 (2015).
- 210 6. Cheng, H. *et al.* The Asian monsoon over the past 640,000 years and ice age  
211 terminations. *Nature*. **534**, 640-646 (2016).
- 212 7. Beck, J. W. *et al.* A 550,000-year record of East Asian monsoon rainfall from <sup>10</sup>Be in  
213 Loess. *Science*. **360**, 877-881 (2018).
- 214 8. Clemens, S.C. *et al.* Precession-band variance missing from East Asian monsoon  
215 runoff. *Nat. Commun.* **9**, 3364 (2018).
- 216 9. Clemens, S. C. & Prell, W. L. 350,000 year summer-monsoon multi-proxy from the  
217 Owen Ridge, Northern Arabian Sea. *Mar. Geol.* **201**, 35-51 (2003).
- 218 10. Caley, T. *et al.* New Arabian Sea records help decipher orbital timing of Indo-Asian  
219 monsoon. *Earth Planet. Sci. Lett.* **308**, 433-444 (2011).
- 220 11. Zhisheng, A. *et al.* Glacial-Interglacial Indian Summer Monsoon Dynamics. *Science*.  
221 **333**, 719-723 (2011).

- 222 12. Bazin, L. *et al.* An optimized multi-proxy, multi-site Antarctic ice and gas orbital  
223 chronology (AICC2012): 120-800 ka. *Clim. Past.* **9**, 1715-1731 (2013).
- 224 13. Sarin, M. M., Krishnaswami, S., Somayajulu, B. L. K. & Moore, W. S. Chemistry of  
225 uranium, thorium, and radium isotopes in the Ganga-Brahmaputra river system:  
226 Weathering processes and fluxes into the Bay of Bengal. *Geochim. Cosmochim. Acta.*  
227 **54**, 1387-1396 (1990).
- 228 14. Singh, S. P. *et al.* Spatial distribution of dissolved neodymium and  $\epsilon_{Nd}$  in the Bay of  
229 Bengal: Role of particulate matter and mixing of water mass. *Geochim. Cosmochim.*  
230 *Acta.* **94**, 38-56 (2012).
- 231 15. Yu, Z. *et al.* Seasonal variations in dissolved neodymium isotope composition in the  
232 Bay of Bengal. *Earth Planet. Sci. Lett.* **479**, 310-321 (2017).
- 233 16. Vance, D. & Burton, K. Neodymium isotopes in planktonic foraminifera: a record of the  
234 response of continental weathering and ocean circulation rates to climate change. *Earth*  
235 *Planet. Sci. Lett.* **173**, 365-379 (1999).
- 236 17. Pomiès, C., Davies, G. R. & Conan, S. M. –H. Neodymium in modern foraminifera from  
237 the Indian Ocean: implications for the use of foraminiferal Nd isotope compositions in  
238 paleo-oceanography. *Earth Planet. Sci. Lett.* **203**, 1031-1045 (2002).
- 239 18. Martínez-Botí, M. A., Vance, D. & Mortyn, P. G. Nd/Ca ratios in plankton-towed and  
240 core top foraminifera: Confirmation of the water column acquisition of Nd. *Geochem.*  
241 *Geophys. Geosyst.* **10**, Q08018 (2009).
- 242 19. Boyle, E. A. & Keigwin, L. D. Comparison of Atlantic and Pacific paleochemical  
243 records for the last 215,000 years: changes in deep ocean circulation and chemical  
244 inventories. *Earth Planet. Sci. Lett.* **76**, 135-150 (1985).

- 245 20. Russell, A. N., Emerson, S., Nelson, B. K., Erez, J. & Lea, D. W. Uranium in  
246 foraminiferal calcite as recorder of seawater uranium concentrations. *Geochim.*  
247 *Cosmochim. Acta.* **58**, 671-681 (1994).
- 248 21. Abdi, H. *Normalizing data: In Salkind, N. (Ed.), Encyclopoedia of Research Design*  
249 (Sage, Thousand Oaks, CA, 2010).
- 250 22. Dearing, J. A. & Jones, R. T. Coupling temporal and spatial dimensions of global  
251 sediment flux through lake and marine sediment records. *Glob Planet Change.* **39**, 147-  
252 168 (2003).
- 253 23. McCreary, J. P., Kundu, P. K. & Molinari, R. L. A numerical investigation of dynamics,  
254 thermodynamics and mixed-layer processes in the Indian Ocean. *Prog. Oceanog.* **31**,  
255 181-244 (1993).
- 256 24. Barker, S. *et al.* Interhemispheric Atlantic seesaw response during the last deglaciation.  
257 *Nature.* **457**, 1097-1101 (2009).
- 258 25. Broecker, W. S. & Henderson, G. M. The sequence of events surrounding Termination  
259 II and their implications for the cause of glacial-interglacial CO<sub>2</sub> changes.  
260 *Paleoceanography.* **13**, 352-364 (1998).
- 261 26. Marino, G. *et al.* Bipolar seesaw control on last interglacial sea level. *Nature.* **522**, 197-  
262 201 (2015).
- 263 27. Knorr, G. & Lohmann, G. Rapid transitions in the Atlantic thermohaline circulation  
264 triggered by global warming and meltwater during the last deglaciation. *Geochem.*  
265 *Geophys. Geosyst.* **8**, Q12006 (2007).
- 266 28. Martrat, B., Jimenez-Amat, P., Zahn, R. & Grimalt, J. O. Similarities and dissimilarities  
267 between the last two deglaciations and interglaciations in the North Atlantic region.  
268 *Quat. Sci. Rev.* **99**, 122-134 (2014).
- 269 29. Cheng, H. *et al.* Ice Age Terminations. *Science.* **326**, 248-252 (2009).

- 270 30. Scussolini, P., Marino, G., Brummer, G. –J. & Peeters, F. J. C. Saline Indian Ocean  
271 waters invaded the South Atlantic thermocline during glacial termination II. *Geology*.  
272 **43**, 139-142 (2015).
- 273 31. Carlson, A. E. & Winsor, K. Northern Hemisphere ice-sheet responses to past climate  
274 warming. *Nat. Geosci.* **5**, 607-613 (2012).
- 275 32. Broccoli, A. J., Dahl, K. A. & Stouffer, R. J. Response of the ITCZ to Northern  
276 Hemisphere cooling. *Geophys. Res. Lett.* **33**, L01702 (2006).
- 277 33. Laskar, J. *et al.* A long-term numerical solution for the insolation quantities of the Earth.  
278 *Astron. Astrophys.* **428**, 261-285 (2004).
- 279 34. Mantis, D. F. *et al.* The Response of Large-Scale Circulation to Obliquity-Induced  
280 Changes in Meridional Heating Gradients. *J. Clim.* **27**, 5504-5516 (2014).
- 281 35. Mudelsee, M. Ramp function regression: a tool for quantifying climate transitions.  
282 *Comput. Geosci.* **26**, 293-307 (2000).
- 283 36. Rodwell, M. J. & Hoskins, B. J. Subtropical Anticyclones and Summer Monsoons. *J.*  
284 *Clim.* **14**, 3192-3211 (2001).
- 285 37. Louergue, L. *et al.* Orbital and millennial-scale features of atmospheric CH<sub>4</sub> over the  
286 past 800,000 years. *Nature.* **453**, 383-386 (2008).
- 287 38. Wang, B., Clemens, S. C. & Liu, P. Contrasting the Indian and East Asian monsoons:  
288 implications on geologic timescales. *Mar. Geol.* **201**, 5-21 (2003).
- 289 39. Caley, T., Roche, D. M. and Renssen, H. Orbital Asian summer monsoon dynamics  
290 revealed using an isotope-enabled global climate model. *Nat. Commun.* **5**, 5371 (2014).
- 291 40. Rodgers, K. B. *et al.* A tropical mechanism for Northern Hemisphere deglaciation.  
292 *Geochem. Geophys. Geosyst.* **4**, 1046 (2003).

- 293 41. Bolton, C. T. *et al.* A 500,000 year record of Indian summer monsoon dynamics  
294 recorded by eastern equatorial Indian Ocean upper water-column structure. *Quat. Sci.*  
295 *Rev.* **77**, 167-180 (2013).
- 296 42. Budziak, D. *et al.* Late Quaternary insolation forcing on total organic carbon and C<sub>37</sub>  
297 alkenone variations in the Arabian Sea. *Paleoceanography*. **15**, 307-321 (2000).
- 298 43. Ziegler, M. *et al.* Precession phasing offset between Indian summer monsoon and  
299 Arabian Sea productivity linked to changes in Atlantic overturning circulation.  
300 *Paleoceanography*. **25**, PA3213 (2010).
- 301 44. Reichert, G. -J., Lourens, L. J. & Zachariasse, W. J. Temporal variability in the northern  
302 Arabian Sea Oxygen Minimum Zone (OMZ) during the last 225,000 years.  
303 *Paleoceanography*. **13**, 607-621 (1998).
- 304 45. Meissner, T. & Wentz, F. J. Remote Sensing Systems SMAP Ocean Surface Salinities  
305 [Level 3 Monthly], Version 2.0 validated release. Remote Sensing Systems, Santa Rosa,  
306 Ca, USA. 2016. Available online at [www.remss.com/missions/smap](http://www.remss.com/missions/smap).
- 307 46. Boyer, T. P. *et al.* *World Ocean Database 2013*. (NEDSIS, Silver Spring, 2013).
- 308 47. Barker, S. *et al.* Icebergs not the trigger for North Atlantic cold events. *Nature*. **520**,  
309 333-336 (2015).
- 310 48. Kudrass, H. R., Hofmann, A., Dooze, H., Emeis, K. & Erlenkeuser, H. Modulation and  
311 amplification of climatic changes in the Northern Hemisphere by the Indian summer  
312 monsoon during the past 80 k.y. *Geology*. **29**, 63-66 (2001).
- 313 49. Rashid, H., Flower, B. P., Poore, R. Z. & Quinn, T. M. A ~25 ka Indian Ocean monsoon  
314 variability record from the Andaman Sea. *Quat. Sci. Rev.* **26**, 2586-2597 (2007).
- 315 50. Saraswat, R., Lea, D. W., Nigam, R., Mackensen, A. & Naik, D. K. Deglaciation in the  
316 tropical Indian Ocean driven by interplay between the regional monsoon and global  
317 teleconnections. *Earth Planet. Sci. Lett.* **375**, 166-175 (2013).

318 51. Pahnke, K. & Sachs, J. P. Sea surface temperatures of southern midlatitudes 0-160 kyr  
319 B. P. *Paleoceanography*. **21**, PA2003 (2006).

320 52. Jouzel, J. *et al.* Orbital and Millennial Antarctic Climate Variability over the Past  
321 800,000 Years. *Science*. **317**, 793-795 (2007).

322 53. Deaney, E. L., Barker, S. & van de Flierdt. T. Timing and nature of AMOC recovery  
323 across Termination 2 and magnitude of deglacial CO<sub>2</sub> change. *Nat. Commun.* **8**, 14595  
324 (2017).

### 325 **Acknowledgements**

326 We thank Peter Webb for his help with setting up pXRF analysis, Hilary Sloane for her help  
327 with stable isotope analysis, P. Divakar Naidu for providing 2005 NBBT sediment traps. We  
328 thank three anonymous reviewers for their insightful comments and suggestions that improved  
329 this manuscript. P. A. and K.N-K. acknowledge funding through a NERC PhD grant  
330 (NE/L002493/1) associated with the CENTA Doctoral Training Partnership. Samples were  
331 provided by the International Ocean Discovery Program (IODP). Stable isotope analysis of  
332 planktic foraminifera was funded by NIGFSC grant IP-1649-1116 to P.A.

### 333 **Author contributions**

334 P.A. conceived the research idea and further developed it with K.N-K. K.N-K processed  
335 samples, picked foraminifera, and conducted foraminifera cleaning and trace element analysis  
336 under guidance from P.A. and S.M. M.J.L. over saw the stable isotope analysis and S.J.H.  
337 helped with trace element analysis. S.C.C. produced benthic oxygen isotope data for age model  
338 development. K.N-K., P.A. and P.F.S. discussed the data interpretation and wrote the  
339 manuscript, and all authors contributed to the final text.

### 340 **Competing interests**

341 The authors declare no competing interests.

342 **Correspondence** should be addressed to [katrina.kerr@open.ac.uk](mailto:katrina.kerr@open.ac.uk) and **requests for materials**  
343 to [pallavi.anand@open.ac.uk](mailto:pallavi.anand@open.ac.uk)

344

345 **Figure 1. ISM induced freshening in the Bay of Bengal** a) Map depicting ISM inferred wind-  
346 driven upwelling and stratification records (as circles: pink<sup>41</sup>, purple<sup>42</sup>, blue<sup>9</sup>, orange<sup>44</sup>, black<sup>45</sup>  
347 and green<sup>10</sup>) that extend across TII. Yellow circle indicates Bittoo cave<sup>2</sup>. b-e) Average monthly  
348 sea surface salinity during 2017 ISM months<sup>45</sup> exhibiting proliferation of fluvial input. Site  
349 U1446 is indicated by red star. f) Winter (black) and summer (red) monsoon season  
350 temperature (solid) and salinity (dashed) depth profiles<sup>46</sup> above Site U1446. Shaded bar  
351 indicates inferred depth range of *G. ruber ss*. Figure created using Ocean Data View software  
352 (<http://odv.awi.de/>).

353

354 **Figure 2. Sequence of global events across TI (A) and TII (B)** a) Obliquity<sup>33</sup> b) June 21<sup>st</sup>  
355 and December 21<sup>st</sup> insolation<sup>33</sup> c) ODP 983, North Atlantic, IRD<sup>47</sup> d) ODP 983 % NPS<sup>47</sup> e)  
356 EASM speleothem  $\delta^{18}\text{O}^6$  f) ODP 976, western Mediterranean, SST<sup>26, 28</sup> g) Bittoo Cave  
357 speleothem  $\delta^{18}\text{O}^2$  h) ISM  $\delta^{18}\text{O}_{\text{sw-IVC}}$  stack<sup>48-50</sup> i) MD97-2120, southwest Pacific, SST<sup>51</sup> j) EDC  
358  $\text{CH}_4$ <sup>37, 12</sup> k) EDC  $\delta\text{D}^{52, 12}$  l) U1446 pXRF stack m) U1446 *G. ruber ss* (red) and *N. dutertrei*  
359 (brown) runoff tracers. Red bar shows modern sediment trap data range n) U1446 *G. ruber ss*  
360 (green) and *N. dutertrei* (grey)  $\delta^{18}\text{O}_{\text{sw-IVC}}$  and o) U1446 SST. Star represents modern day mean  
361 annual SST at study site<sup>46</sup>. Shaded envelopes represent  $1\sigma$  (Methods). Red triangles represent  
362 age control points contained within interval shown and associated AICC2012 chronology  
363 errors<sup>12</sup> (Methods).

364



365 **Figure 3. TII onset and duration** a) ODP 983, North Atlantic, % NPS<sup>47</sup> on AICC2012  
366 chronology<sup>12</sup> b) ODP 1063, Atlantic Ocean, % Warm species<sup>53</sup> on AICC2012 chronology<sup>12</sup> c)  
367 EASM speleothem  $\delta^{18}\text{O}$ <sup>6</sup> d) ODP 976, western Mediterranean Sea, SST<sup>28</sup> on Corchia Cave  
368 radiometrically constrained chronology<sup>26</sup> e) U1446 *G. ruber ss* runoff tracers (this study) f)  
369 U1446 *G. ruber ss*  $\delta^{18}\text{O}_{\text{sw-IVC}}$  (this study) g) U1446 pXRF stack h) EDC  $\delta\text{D}$ <sup>52</sup> on AICC2012  
370 chronology<sup>12</sup>. Pink shaded area denotes  $t_2$  (deglaciation onset) and  $t_1$  (attainment of  
371 interglacial) as modelled using RAMPFIT<sup>35</sup> (Methods).

## 372 **Methods**

373 Site U1446 (19°5.02'N, 85°44'E) was drilled during IODP Expedition 353 and located at a  
374 depth of 1430 meters below sea level in the Mahanadi Basin<sup>54</sup>. The BoB represents the core  
375 convective region of the ISM due to the thermodynamic structure of the water column resulting  
376 in positive ocean-atmosphere feedbacks favouring high SSTs (>28°C) allowing convection to  
377 be sustained during the summer monsoon months of June through to September<sup>55</sup>. The ISM  
378 exerts a strong seasonal signature of surface water freshening and stratification within the BoB  
379 due to a net surface water exchange of  $184 \times 10^{10} \text{ m}^3$  during the ISM months<sup>56</sup>. ISM induced  
380 river runoff generates a north-south salinity gradient; the northern BoB undergoes a reduction  
381 in salinity of 9‰ during this period<sup>57</sup>.

## 382 **Age Model**

383 The much expanded nature of the sediment sequence at Site U1446 (~25 cm/ka), and  
384 consequent high fidelity of our palaeoclimatic records, significantly reduces the error of the  
385 duration of events and the rates of change inferred from our records<sup>58</sup>. Using Analyseries<sup>59</sup> we  
386 graphically correlated benthic foraminifera (*Uvigerina spp.* and *Cibicidoides wuellerstorfi*)  
387  $\delta^{18}\text{O}$  (Clemens, S. C., unpublished data) to benthic  $\delta^{18}\text{O}$  from south Pacific core PS75/059-2<sup>60</sup>  
388 (Supplementary Fig. 1a). This itself is tied to the AICC2012 chronology<sup>12</sup> by exploiting the

389 age-depth relationship from PS75/059-2 Fe dust flux record<sup>61</sup> which has been tuned to the EDC  
390 Antarctic ice core<sup>61, 62</sup> (Supplementary Fig. 2). Tuning to AICC2012 was chosen rather than  
391 the absolute dated EASM speleothem record to allow for independent assessment of the  
392 lead/lag relationship between the ISM and the EASM. We infer that our records are not biased  
393 to the high latitudes of the southern hemisphere by our tuning strategy due to synchronicity  
394 existing between the Chinese Loess magnetic susceptibility record with EDC Antarctic ice core  
395 dust fluxes<sup>62</sup>. To ascertain our confidence in our age model, we further tied U1446 benthic  $\delta^{18}\text{O}$   
396 to ODP Leg. 117, Site 1146 benthic  $\delta^{18}\text{O}$  which has been transferred to the speleothem  
397 chronology<sup>63</sup>. We present site U1446 benthic  $\delta^{18}\text{O}$  on three different age models (AICC2012<sup>12</sup>,  
398 RC2011<sup>63</sup> and LR04<sup>64</sup>) (Supplementary Fig. 1c) in order to confirm the lead of U1446 ISM  
399 records over the EASM across TII regardless of chronology (Supplementary Fig. 3).

400 We used Bchron<sup>65</sup>, a Bayesian probability model, to model the 95% uncertainty envelope  
401 between tie points with the AICC2012 chronology error (modelled as Gaussian distribution)  
402 of EPICA Dome C at those points<sup>12</sup> (Supplementary Fig. 1b).

403 All datasets used to assess relative lead and lag relationships are on a consistent age-model;  
404 that of AICC2012<sup>12</sup> or absolute radiometrically constrained chronology<sup>2, 6, 26</sup> (see original  
405 references for detail).

#### 406 **Foraminiferal stable isotope and trace metal analysis**

407 The planktic foraminifera *Globigerinoides ruber sensu-stricto* (*ss*), was identified using the  
408 taxonomic description in ref. 66. Between 6 to 30 individuals were picked from the 250-355 $\mu\text{m}$   
409 size-fraction and gently crushed prior to analysis. Oxygen isotope analyses were performed at  
410 the British Geological Survey, NERC Isotope Geoscience Facilities, Keyworth using an  
411 Isoprime dual inlet mass spectrometer with Multiprep device. The reproducibility of oxygen  
412 isotope measurements is  $\pm 0.05\text{‰}$  ( $1\sigma$ ) based on replicate measurements of carbonate standards.

413 All data are reported in the usual delta notation ( $\delta^{18}\text{O}$ ) in ‰ on the VPDB scale.

414 For trace metal analysis, samples were cleaned using a modification of the method described  
415 in ref. 19 and reversal of the oxidative and reductive steps<sup>67</sup>. Due to the proximal setting of Site  
416 U1446 an extended clay removal step was essential in order to ensure removal of any fine clays  
417 that may bias Mg content in carbonate samples. Samples were initially rinsed with repeated  
418 MQ and methanol rinses with ultrasonification of 40 seconds between each rinse. Samples were  
419 then inspected under a microscope and any discoloured fragments, fragments with pyrite or  
420 silicate particles were removed. Subsequently samples were subjected to a reductive and 10%  
421 oxidative step to ensure removal of any coatings and organics. Samples were then polished  
422 using a weak (0.001M) HNO<sub>3</sub> leaching step and dissolved (0.075M HNO<sub>3</sub>) on the day of  
423 analysis. Samples were analysed at the Open University using an Agilent Technologies Triple-  
424 Quad ICP-MS. Contaminant ratios (Al/Ca and Fe/Ca) were monitored in order to assess any  
425 clay and organic contaminations (Supplementary Fig. 4).

#### 426 **Estimating temperature and $\delta^{18}\text{O}_{\text{sw}}$**

427 The addition of a reductive step during foraminiferal trace element cleaning has been shown to  
428 reduce Mg/Ca values<sup>68</sup>. Following ref. 69, we apply a correction for a 10% reduction in Mg/Ca  
429 associated with the reductive method due to the chosen temperature calibration being based on  
430 analysis using only the oxidative step<sup>70</sup>. The Mg/Ca temperature calibration used was  
431 accordingly adjusted:

$$432 \quad \text{Mg/Ca} = 0.38(\pm 0.02) \exp((0.09 \pm 0.003) * T)^{70}$$

$$433 \quad \text{Adjusted Mg/Ca} = 0.342 \exp(0.09T)$$

434 An ice volume correction was applied to the calcite  $\delta^{18}\text{O}_c$  following the Red Sea Level Curve  
435 (95% probability maximum)<sup>71</sup> with a conversion factor  $\delta^{18}\text{O}$  enrichment of 0.008‰ per meter  
436 sea level lowering applied<sup>72</sup>:

$$437 \quad \delta^{18}\text{O}_{\text{IVC}}(t) = \delta^{18}\text{O}(t) + (\text{RSL}(t) * 0.008)$$

438 The temperature estimates derived from Mg/Ca and the measured calcite  $\delta^{18}\text{O}_c$  of planktic  
439 foraminifera allows for the derivation of seawater  $\delta^{18}\text{O}_{\text{sw}}$ :

$$440 \quad T^{\circ}\text{C} = 14.9(\pm 0.1) - 4.8(\pm 0.08) * (\delta^{18}\text{O}_c - \delta^{18}\text{O}_{\text{sw}}) - 0.27\text{‰}^{73}$$

441 The  $\delta^{18}\text{O}_{\text{sw}}$  has been shown to correlate strongly with salinity in the northern BoB. Factors  
442 controlling this relationship include precipitation, river runoff and evaporation thus during the  
443 summer monsoon months precipitation and runoff exceeds evaporation promoting a low  
444  $\delta^{18}\text{O}_{\text{sw}}$ -Salinity Slope<sup>74, 75</sup>. However, we do not convert U1446  $\delta^{18}\text{O}_{\text{sw}}$  to salinity using modern  
445 day calculated regressions due to observation of significant spatiotemporal variations and  
446 uncertainties in assumptions associated with extending these relationships into the past<sup>74</sup>.  
447 Furthermore, recent work has indicated the potential control salinity exerts on Mg-  
448 incorporation in foraminiferal calcite<sup>76</sup>. Low salinity during the warmer ISM season may  
449 potentially dampen our reconstructed SSTs based on Mg/Ca relative to actual SST however,  
450 there would be a limited overall effect on the reconstructed  $\delta^{18}\text{O}_{\text{sw}}$ .

451 *N. dutertrei* is typically inferred to represent thermocline conditions (~70-120m)  
452 accompanying the deep chlorophyll maximum<sup>77, 78</sup>. However, across the TII IS *N. dutertrei*  
453 shows more depleted  $\delta^{18}\text{O}_{\text{sw-IVC}}$  values than surface dwelling *G. ruber ss* (Fig. 2B n). We infer  
454 that this is associated with the unique hydrographic conditions that Site U1446 experiences and  
455 that *N. dutertrei* occupies a shallower depth, in the freshwater lens of the upper water column,  
456 than what is typically inferred. Additionally, available Mg/Ca calibrations based on upper  
457 thermocline habitat, and therefore a narrower temperature range, underestimates the  
458 temperature values for *N. dutertrei* thus resulting in more depleted  $\delta^{18}\text{O}_{\text{sw-IVC}}$  values as the  
459 calcite  $\delta^{18}\text{O}$  values are more enriched than *G. ruber ss* (Supplementary Fig. 5). During the TII  
460 IS, *G. ruber ss* and *N. dutertrei*  $\delta^{18}\text{O}_{\text{sw-IVC}}$  is decoupled by ~100 years (Fig. 2B n) highlighting  
461 the vertical flux of ISM induced freshening.

462 Error propagation of the temperature and  $\delta^{18}\text{O}_{\text{sw}}$  estimates was calculated using the following  
 463 equations<sup>79</sup> where Mg/Ca standard deviation is  $0.029\text{mmol/mol}^{-1}$  and  $\delta^{18}\text{O}_c$  is  $0.05\text{‰}$  based on  
 464 repeated analysis of internal standards. The error propagation is based on assumptions of no  
 465 covariance among a, b, T and  $\delta^{18}\text{O}_c$ <sup>79</sup>:

$$466 \quad \sigma_T^2 = \left(\frac{\partial T}{\partial a} \sigma_a\right)^2 + \left(\frac{\partial T}{\partial b} \sigma_b\right)^2 + \left(\frac{\partial T}{\partial \text{Mg/Ca}} \sigma_{\text{Mg/Ca}}\right)^2$$

467 where:

$$468 \quad a = 0.342(\pm 0.02)^{70}$$

$$469 \quad b = (0.09 \pm 0.003)^{70}$$

$$470 \quad \frac{\partial T}{\partial a} = -\frac{1}{a^2} \ln\left(\frac{\text{Mg/Ca}}{b}\right)$$

$$471 \quad \frac{\partial T}{\partial b} = -\frac{1}{ab}$$

$$472 \quad \frac{\partial T}{\partial \text{Mg/Ca}} = \frac{1}{a} \times \frac{1}{\text{Mg/Ca}}$$

$$473 \quad \sigma_{\delta^{18}\text{O}_{\text{sw}}}^2 = \left(\frac{\partial \delta^{18}\text{O}_{\text{sw}}}{\partial T} \sigma_T\right)^2 + \left(\frac{\partial \delta^{18}\text{O}_{\text{sw}}}{\partial a} \sigma_a\right)^2 + \left(\frac{\partial \delta^{18}\text{O}_{\text{sw}}}{\partial b} \sigma_b\right)^2 + \left(\frac{\partial \delta^{18}\text{O}_{\text{sw}}}{\partial \delta^{18}\text{O}_c} \sigma_{\delta^{18}\text{O}_c}\right)^2$$

474 where:

$$475 \quad a = 14.9(\pm 0.1)^{73}$$

$$476 \quad b = -4.8(\pm 0.08)^{73}$$

$$477 \quad \frac{\partial \delta^{18}\text{O}_{\text{sw}}}{\partial T} = -\frac{1}{b}$$

$$478 \quad \frac{\partial \delta^{18}\text{O}_{\text{sw}}}{\partial a} = \frac{1}{b}$$

479 
$$\frac{\partial \delta^{18}\text{O}_{\text{sw}}}{\partial b} = \frac{T}{b^2} - \frac{a}{b^2}$$

480 
$$\frac{\partial \delta^{18}\text{O}_{\text{sw}}}{\partial \delta^{18}\text{O}_{\text{c}}} = 1$$

481 To further constrain errors associated with calculating SST and  $\delta^{18}\text{O}_{\text{sw}}$  we used Paleo-Seawater  
482 Uncertainty Solver (PSUSolver)<sup>80</sup>. PSUSolver models uncertainties associated with age model,  
483 calibrations, analytical and sea level estimate errors by performing bootstrap Monte Carlo  
484 simulations<sup>80</sup>. Accounting for AICC2012 age model errors<sup>12</sup> we input an average age model  
485 error of 2 ka and analytical errors; Mg/Ca of 0.029mmol/mol<sup>-1</sup> and  $\delta^{18}\text{O}_{\text{c}}$  is 0.05‰, in order for  
486 PSUSolver to probabilistically constrain the median estimate and confidence intervals for SST  
487 and  $\delta^{18}\text{O}_{\text{sw}}$  (Supplementary Fig. 6a). To assess the influence age model error exerts on U1446  
488 SST and  $\delta^{18}\text{O}_{\text{sw}}$  we also input an age model error of 1 ka (Supplementary Fig. 6b) and 0 ka  
489 (Supplementary Fig. 6c). This indicates that age model errors exert the strongest influence on  
490 PSUSolver SST and  $\delta^{18}\text{O}_{\text{sw}}$ . An average age model error of 2 ka renders the TII IS  
491 inconspicuous. However, we have confidence in our original U1446 SST and  $\delta^{18}\text{O}_{\text{sw}}$   
492 interpretations despite the associated errors with the AICC2012 chronology owing to TII IS  
493 having been resolved in other independently dated records (Fig. 2B) and the coherence of  
494 U1446  $\delta^{18}\text{O}_{\text{sw}}$  with deglacial warming in western Mediterranean Sea SST records from ODP  
495 Site 976<sup>28</sup> (Fig. 3) that has a radiometrically constrained age model<sup>26</sup>.

#### 496 **Interpreting Mn/Ca, Nd/Ca & U/Ca as river runoff proxies**

497 Mn/Ca ratios measured in foraminifera are typically used as an indicator of contamination of  
498 foraminifer calcite from authigenic Mn-rich oxide coatings on the foraminifer shell. Our Mn/Ca  
499 data display no correlation with Mg/Ca ( $r^2=0.0894$ ), strongly arguing against the presence of  
500 Mn-rich oxide coatings on our foraminifera that would bias our Mg/Ca-derived SSTs. The  
501 foraminifera cleaning method applied in this study had the reductive cleaning step included,

502 which ensures removal of Fe-Mn coatings, added on the carbonate tests at the sediment-water  
503 interface<sup>19, 68</sup>. Mn/Ca correlates with Nd/Ca and U/Ca (Supplementary Fig. 7), reinforcing  
504 evidence that these elements are delivered to our study site via fluvial runoff and can thus be  
505 used as runoff proxies in this proximal setting. High fluvial fluxes in the BoB reflect the  
506 monsoon region's vigorous hydrological and concomitant weathering regime. This is  
507 expressed by the vast quantities of material discharged via the rivers; the Ganges-Brahmaputra  
508 systems contribute alone  $1.06 \times 10^9$  tonnes of sediment annually<sup>81</sup>. Such a unique hydrographic  
509 setting allows high concentrations of dissolved lithogenic elements (Mn, Nd, U) to be  
510 precipitated (either as authigenic or biogenic carbonate phases) upon mixing with seawater.  
511 The observed concentrations of these elements at Site U1446 are well beyond the  
512 concentrations that are typically found in planktic foraminifera<sup>20</sup>. Similarly, elevated levels of  
513 Mn/Ca, Nd/Ca and U/Ca ratios have been found in planktic foraminifera from Ceara Rise, ODP  
514 Site 926, receiving amazon fluvial fluxes<sup>82, 83</sup>. Furthermore, we generated trace element data  
515 for *G. ruber ss* from NBBT-05-S sediment trap from the northern BoB. The range of values  
516 exhibited by this runoff tracers record (Mn, Nd and U) overlaps with the range found in the  
517 NBBT-05-S sediment trap data (Fig. 2B m). Thus, we interpret Mn/Ca, Nd/Ca and U/Ca ratios  
518 in *G. ruber ss* (Supplementary Fig. 8) as a proxy for fluvial runoff at marginal sites and suggest  
519 that they could be further ground-truthed for application in other marginal marine settings.  
520 Owing to the similarity between Mn/Ca, Nd/Ca and U/Ca we normalise using the standard  
521 deviation<sup>21</sup>:

$$522 \quad /Ca(t)_{\text{norm}} = \frac{/Ca(t) - \overline{/Ca}}{\sigma(/Ca)}$$

523 Where:

524  $/Ca(t)$  (e.g. Mn/Ca) represents the trace element to Ca ratio at a given time.

525  $\overline{X}/Ca$  represents the mean of all the trace element to Ca ratios (e.g. Mn/Ca) across study  
526 interval.

527  $\sigma(X)/Ca$  represents the standard deviation of the trace element to Ca ratio across study interval.

528 Subsequently we average these values ( $X/Ca(t)_{norm}$ ) for each of the tracers to produce a factor  
529 representing *G. ruber ss* runoff tracers. Furthermore, there is a similar signature among these  
530 tracers with the data gained from pXRF (Supplementary Fig. 9).

### 531 **Discrete portable X-Ray Fluorescence Analysis**

532 Analysis of major and minor elements was performed using a Niton XL3t900 portable X-Ray  
533 Fluorescence (pXRF). Prior to analysis 5 grams of material was weighed, dried in an oven at  
534 40°C and subsequently homogenized into a fine powder through use of a pestle and mortar.  
535 The powdered material was transferred into 7ml vials, sealed tightly with non-PVC Clingfilm  
536 and placed flush over the aperture of the X-ray emitter (Saker-Clark, M., per comms).  
537 Calibration for each element of interest was performed through analysis of geochemical in-  
538 house and reference powdered rock standards with known concentrations. A set of internal and  
539 reference standards were run every 10<sup>th</sup> sample for quality control (Supplementary Table 1).  
540 Bulk sediment elemental geochemistry is controlled by detrital (i.e. terrigenous input via river  
541 runoff) and authigenic processes. Therefore, in order to reconstruct ISM derived river runoff a  
542 selection of inferred terrigenous derived elements were selected to represent increased fluvial  
543 runoff and detrital input to the site; Ti, K, Al and Rb (Supplementary Fig. 9). These elements  
544 were combined through normalising to unit variance (described in the above section for *G.*  
545 *ruber ss* runoff tracers) to produce a factor of pXRF runoff element variations<sup>21</sup> due to showing  
546 strong correlation with each other (Supplementary Fig. 10). In order to clarify the inconsistency  
547 in elements chosen to represent fluvial runoff between the pXRF element stack and the *G.*  
548 *ruber ss* tracers: i) Uranium concentrations in discrete U1446 samples were below detection



549 limit and Nd was not measured and ii) Mn concentrations in ocean sediments is complicated  
550 by redox processes and therefore, not a suitable candidate for representing the detrital phase in  
551 bulk sediment elemental profiles. We infer that due to increased terrigenous supply during a  
552 strengthened ISM, reduced bottom water conditions are established, resulting in Mn reduction  
553 and dissolution into pore waters due to the increased solubility of reduced Mn ( $Mn^{2+}$ )<sup>84-87</sup>. In  
554 contrast, during times of weaker ISM and reduced terrigenous supply, aerobic conditions  
555 promote formation of solid-phase Mn oxyhydroxides and thus increase in Mn concentrations in  
556 the bulk sediment (Supplementary Fig. 9)<sup>84-87</sup>. This reasoning is coherent with conditions found  
557 in the Cariaco Basin, proximal to high terrigenous fluxes via river runoff<sup>88</sup>.

### 558 **Detection of TII Change Points**

559 In order to empirically assess deglaciation onset during TII we employed the RAMPFIT<sup>35</sup>  
560 algorithm. RAMPFIT segments the data into three parts using a weighted least squares  
561 regression and brute force to find two breakpoints denoted as  $t1$  and  $t2$ <sup>35</sup>. RAMPFIT was used  
562 to estimate deglaciation onset ( $t1$ ) and duration ( $t2$ ) in the EASM speleothem  $\delta^{18}O$  record<sup>6</sup>,  
563 ODP 976 western Mediterranean Sea SST<sup>26,28</sup>, ODP 1063 % warm species<sup>53</sup>, ODP 983 %  
564 NPS<sup>47</sup>, EPICA Dome C  $\delta D$ <sup>52</sup> and U1446  $\delta^{18}O_{sw}$ , *G. ruber ss* runoff tracers and pXRF stack  
565 (Fig. 3). These records were chosen in order to identify the proliferation of deglaciation across  
566 the NH having propagated from the SH. 400 iterations of wild bootstrap with seed generator  
567 number of 400 was used to determine the uncertainties (Supplementary Table. 2).

### 568 **Comparison of TII with TI**

569 The same methods described above were employed to characterise deglaciation across TI  
570 (Supplementary Fig. 10). Our results for TII demonstrate the sequence of deglaciation having  
571 been driven from the SH, a lagged NH response and the ISM contributing to the inter-  
572 hemispheric transfer of heat and moisture. Furthermore, we highlight the out-of-phase  
573 behaviour between the EASM and ISM (Fig. 3). However, this is in contrast to the sequence

574 of events across TI in which the ISM appears to be in-phase with the EASM and other NH  
575 climate records (Supplementary Fig. 11). Our results from TII thus exemplify the heterogeneity  
576 between TI and TII that draws on previous work in which orbital preconditioning is regarded  
577 as the driver in dictating the internal climate feedback response<sup>89, 90</sup>. Furthermore, the  
578 behaviour of the ISM during TII may be a result of the anomalous orbital conditions which  
579 stray from classic Milankovitch theory<sup>91</sup>. The early rise in NH solar insolation during TI is  
580 thought to have initiated deglaciation with rapid NH ice sheet retreat occurring from ~19-20  
581 ka<sup>92</sup> resulting in AMOC shutdown and subsequent warming in the SH<sup>93</sup>. This is in contrast to  
582 TII where the earlier rise in SH summer insolation occurs 10 ka prior to NH solar insolation  
583 increase<sup>25, 94</sup>. We postulate based on the opposing hemispheric controls on the ISM during TI  
584 and TII that the ISM is not hemispherically biased but is governed by inter-hemispheric climate  
585 controls in comparison to the predominantly NH-forced EASM<sup>6</sup>.

## 586 **Methods References**

- 587 54. Clemens, S. C. *et al.* Indian Monsoon Rainfall. *Proceedings of the International Ocean*  
588 *Discovery Program*. **353**, (2016).
- 589 55. Shenoi, S. S. C., Shankar, D. & Shetye, S. R. Differences in heat budgets of the near-  
590 surface Arabian Sea and Bay of Bengal: implications for the summer monsoon. *J.*  
591 *Geophys. Res. Oceans*. **107**, 2052 (2002).
- 592 56. Varkey, M. J., Murty, V. S. N. & Suryanarayana, A. Physical oceanography of the Bay  
593 of Bengal and Andaman Sea. *Oceanogr. Mar. Biol. Annu. Rev.* **34**, 1-70 (1996).
- 594 57. Levitus, S. & Mishonov, A. *World Ocean Atlas 2013, Volume 2: Salinity* (NESDIS,  
595 Silver Spring, 2013).
- 596 58. Kemp, D. B. & Sexton, P. F. Time-scale uncertainty of abrupt events in the geologic  
597 record arising from unsteady sedimentation. *Geology*. **42**, 891-894 (2014).

- 598 59. Paillaird, D., Labeyrie, L. & Yiou, P. Macintosh Program performs time-series analysis.  
599 *EOS*. **77**, 379 (1996).
- 600 60. Ullermann, J. *et al.* Pacific-Atlantic Circumpolar Deep Water coupling during the last  
601 500 ka. *Paleoceanography*. **31**, 639-650 (2016).
- 602 61. Lamy, F. *et al.* Increased Dust Deposition in the Pacific Southern Ocean During Glacial  
603 Periods. *Science*. **343**, 403-407 (2014).
- 604 62. Lambert, F. *et al.* Dust-climate couplings over the past 800,000 years from the EPICA  
605 Dome C ice core. *Nature*. **452**, 616-619 (2008).
- 606 63. Caballero-Gill, R. P., Clemens, S. C. & Prell, W. L. Direct correlation of Chinese  
607 speleothem  $\delta^{18}\text{O}$  and South China Sea planktonic  $\delta^{18}\text{O}$ : Transferring a speleothem  
608 chronology to the benthic marine chronology. *Paleoceanography*. **27**, PA2203 (2012).
- 609 64. Lisiecki, L. E. & Raymo, M. E. A Pliocene-Pleistocene stack of 57 globally distributed  
610  $\delta^{18}\text{O}$  records. *Paleoceanography*. **20**, PA1003 (2005).
- 611 65. Haslett, J. & Parnell, A. A simple monotone process with application to radiocarbon-  
612 dated depth chronologies. *Stat Soc Ser C Appl Stat*. **57**, 339-418 (2008).
- 613 66. Wang, L. Isotopic signals in two morphotypes of *Globigerinoides ruber* (white) from  
614 the South China Sea: implications for monsoon climate change during the last glacial  
615 cycle. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **161**, 381-394 (2000).
- 616 67. Rosenthal, Y., Boyle, E. A. & Labeyrie, L. Last glacial maximum paleochemistry and  
617 deepwater circulation in the Southern Ocean: Evidence from foraminiferal cadmium.  
618 *Paleoceanography*. **12**, 787-796 (1997).
- 619 68. Barker, S., Greaves, M. & Elderfield, H. A study of cleaning procedures used for  
620 foraminiferal Mg/Ca paleothermometry. *Geochem. Geophys. Geosyst.* **4**, 8407 (2003).
- 621 69. Gibbons, F. T. *et al.* Deglacial  $\delta^{18}\text{O}$  and hydrologic variability in the tropical Pacific and  
622 Indian Oceans. *Earth Planet Sci. Lett.* **387**, 240-251 (2014).

- 623 70. Anand, P., Elderfield, H. & Conte, M. H. Calibration of Mg/Ca thermometry in  
624 planktonic foraminifera from a sediment trap time series. *Paleoceanography*. **18**, 1050  
625 (2003).
- 626 71. Grant, K. M. *et al.* Rapid coupling between ice volume and polar temperature over the  
627 past 150,000 years. *Nature*. **491**, 744-747 (2012).
- 628 72. Adkins, J. F., McIntyre, K. & Schrag, D. P. The Salinity, Temperature, and  $\delta^{18}\text{O}$  of the  
629 Glacial Deep Ocean. *Science*. **298**, 1769-1773 (2002).
- 630 73. Bemis, B. E., Spero, H. J., Bijma, J. & Lea, D. W. Reevaluation of the oxygen isotopic  
631 composition of planktonic foraminifera: Experimental results and revised  
632 paleotemperature equations. *Paleoceanography*. **13**, 150-160 (1998).
- 633 74. Singh, A., Jani, R. A. & Ramesh, R. Spatiotemporal variations of the  $\delta^{18}\text{O}$ -salinity  
634 relation in the northern Indian Ocean. *Deep-Sea. Res. I*. **57**, 1422-1431 (2010).
- 635 75. Delaygue, G. *et al.* Oxygen isotope/salinity relationship in the northern Indian Ocean.  
636 *J. Geophys. Res.* **106**, 4565-4574 (2001).
- 637 76. Gray, W. R. *et al.* The effects of temperature, salinity, and the carbonate system on  
638 Mg/Ca in *Globigerinoides ruber* (white): A global sediment trap calibration. *Earth*.  
639 *Planet. Sci. Lett.* **482**, 607-620 (2018).
- 640 77. Ravelo, A. C. & Fairbanks, R. G. Oxygen Isotopic Composition of Multiple Species of  
641 Planktonic Foraminifera: Records of the Modern Photic Zone Temperature Gradient.  
642 *Paleoceanography*. **7**, 815-831 (1992).
- 643 78. Mohtadi, M. *et al.* Reconstructing the thermal structure of the upper ocean: Insights  
644 from planktic foraminifera shell chemistry and alkenones in modern sediments of the  
645 tropical eastern Indian Ocean. *Paleoceanography*. **26**, PA3219 (2011).
- 646 79. Bevington, P. R. & Robinson, K. D. *Data Reduction and Error Analysis for the Physical*  
647 *Sciences 3<sup>rd</sup> edn* (McGraw-Hill, 2003).

- 648 80. Thirumalai, K., Quinn, T. M. & Marino, G. Constraining past seawater  $\delta^{18}\text{O}_{\text{sw}}$  and  
649 temperature records developed from foraminiferal geochemistry. *Paleoceanography*.  
650 **31**, 1409-1422 (2016).
- 651 81. Milliman, J. D. & Syvitski, J. P. M. Geomorphic/Tectonic Control of Sediment  
652 Discharge to the Ocean: The Importance of Small Mountainous Rivers. *J. Geol.* **100**,  
653 525-544 (1992).
- 654 82. Stewart, J. A., James, R. H., Anand, P. & Wilson, P. A. Silicate Weathering and Carbon  
655 Cycle Controls on the Oligocene-Miocene Glaciation. *Paleoceanography*. **32**, 1070-  
656 1085 (2017).
- 657 83. Stewart, J. A., Gutjahr, M., James, R. H., Anand, P. & Wilson, P. A. Influence of the  
658 Amazon River on the Nd isotope composition of deep water in the western equatorial  
659 Atlantic during the Oligocene-Miocene transition. *Earth. Planet. Sci. Lett.* **454**, 132-141  
660 (2016).
- 661 84. Calvert, S. E. & Price, N. B. Diffusion and reaction profiles of dissolved manganese in  
662 the pore waters of marine sediments. *Earth Planet. Sci. Lett.* **16**, 245-249 (1972).
- 663 85. Thomson, J., Higgs, N. C., Croudace, I. W., Colley, S. & Hydes, D. J. Redox zonation  
664 of elements at an oxic/post-oxic boundary in deep-sea sediments. *Geochim. Cosmochim.*  
665 *Acta.* **57**, 579-595 (1993).
- 666 86. Calvert, S. E. & Pedersen, T. F. Geochemistry of Recent oxic and anoxic marine  
667 sediments: Implications for the geological record. *Mar Geol.* **113**, 67-88 (1993).
- 668 87. Burdige, D. J. The biogeochemistry of manganese and iron reduction in marine  
669 sediments. *Earth. Sci. Rev.* **35**, 249-284 (1993).
- 670 88. Yarincik, K. M., Murray, R. W., Lyons, T. W., Peterson, L. C. & Haug, G. H.  
671 Oxygenation history of bottom waters in the Cariaco Basin, Venezuela, over the past

672 578,000 years: Results from redox-sensitive metals (Mo, V, Mn, and Fe).  
673 *Paleoceanography*. **15**, 593-604 (2000).

674 89. Carlson, A. E. Why there was not a Younger Dryas-like event during the Penultimate  
675 Deglaciation. *Quat. Sci. Rev.* **27**, 882-887 (2008).

676 90. Alley, R. B., Brook, E. J. & Anandakrishnan, S. A northern lead in the orbital band:  
677 north-south phasing of Ice-Age events. *Quat. Sci. Rev.* **21**, 431-441 (2002).

678 91. Hays, J. D., Imbrie, K. & Shackleton, N. J. Variations in the Earth's Orbit: Pacemaker  
679 of the Ice Ages. *Science*. **194**, 1121-1132 (1974).

680 92. Clark, P. U. *et al.* The Last Glacial Maximum. *Science*. **325**, 710-714 (2009).

681 93. He, F. *et al.* Northern Hemisphere forcing of Southern Hemisphere climate during the  
682 last deglaciation. *Nature*. **494**, 81-85 (2013).

683 94. Masson-Delmotte, V. *et al.* Abrupt change of Antarctic moisture origin at the end of  
684 Termination II. *Proc. Nat. Acad. Sci.* **107**, 12091-12094 (2010).

685

## 686 **Data Availability**

687 Data generated from this study (IODP Exp. 353, Site U1446) are available via the National  
688 Geoscience Data Centre (NGDC), DOI: 10.5285/061d77af-a805-4cf0-b969-0b8f042fae74.

689 Antarctic EDC ice-core records presented on AICC2012 chronology are available from:

690 <https://doi.pangaea.de/10.1594/PANGAEA.824883> and

691 <https://doi.pangaea.de/10.1594/PANGAEA.824891>

692 The EASM composite speleothem  $\delta^{18}\text{O}$  record is available from:

693 <https://www.ncdc.noaa.gov/paleo-search/study/20450>

694 Bittoo Cave speleothem  $\delta^{18}\text{O}$  record is available from:

695 <https://www.ncdc.noaa.gov/paleo-search/study/20449>

696 ODP 983 and 1063 data is available as a supplementary data set associated with Ref. 53.

697 ODP 976, western Mediterranean Sea SST data on Corchia radiometrically constrained  
698 chronology is available as a supplementary dataset associated with Ref. 26.

699 Benthic  $\delta^{18}\text{O}$  of PS75/059-2 is available at: <https://doi.org/10.1594/PANGAEA.833422>

700 PS75/059-2 on AICC2012 chronology at: <https://doi.org/10.1594/PANGAEA.826580>.

701

702

703

704

705

706

707

708

709

710

711

712

713

714

715

716