



## Open Research Online

### Citation

Butcher, Frances E.G.; Balme, Matthew R.; Gallagher, Colman; Storrar, Robert D.; Conway, Susan J.; Arnold, Neil S.; Lewis, Stephen R. and Hagermann, Axel (2019). Morphometries of eskers on Mars: comparisons to eskers on Earth and implications for sediment-discharge dynamics of subglacial drainage. In: 20th Congress of the International Union for Quaternary Research (INQUA), 25-31 Jul 2019, Dublin, Ireland.

### URL

<https://oro.open.ac.uk/62798/>

### License

(CC-BY-NC-ND 4.0)Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

### Policy

This document has been downloaded from Open Research Online, The Open University's repository of research publications. This version is being made available in accordance with Open Research Online policies available from [Open Research Online \(ORO\) Policies](#)

### Versions

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding

## Morphometries of eskers on Mars: comparisons to eskers on Earth and implications for sediment-discharge dynamics of subglacial drainage.

Frances E.G. Butcher<sup>1</sup>, Matt R. Balme<sup>1</sup>, Colman Gallagher<sup>2,3</sup>, Robert D. Storrar<sup>4</sup>, Susan J. Conway<sup>5</sup>, Neil S. Arnold<sup>6</sup>, Stephen R. Lewis<sup>1</sup>, Axel Hagermann<sup>7</sup>

<sup>1</sup>School of Physical Sciences, The Open University, Milton Keynes, United Kingdom.

<sup>2</sup>UCD School of Geography, University College Dublin, Dublin, Ireland. <sup>3</sup>UCD Earth

Institute, University College Dublin, Dublin, Ireland. <sup>4</sup>Department of the Natural and

Built Environment, Sheffield Hallam University, Sheffield, Ireland. <sup>5</sup>CNRS, UMR6122,

LPG Université de Nantes, Nantes, France. <sup>6</sup>Scott Polar Research Institute, University

of Cambridge, Cambridge, United Kingdom. <sup>7</sup>Biological and Environmental Sciences, University of Stirling, Stirling, United Kingdom

### Abstract

Mars is a glaciated planet with an extremely cold and hyper-arid climate. It hosts water ice within polar ice caps, ground ice, and thousands of mid-latitude debris-covered glaciers. Until recently, it was widely thought that mid-latitude glaciers on Mars' had been pervasively cold-based since their formation 10s–100s Myr ago.

However, our recent discoveries of eskers associated with ~110–150 Myr old glaciers in the Phlegra Montes [1] and Tempe Terra [2] regions of Mars' northern mid-latitudes indicate that localised wet-based glaciation has occurred during Mars' most recent geological period, possibly driven by locally-elevated geothermal heat flux and strain heating within the glacial ice [1–3]. Eskers are sinuous ridges comprising sediments deposited in glacial meltwater conduits. They are important tools for reconstructing the extent and dynamics of wet-based glaciation on Earth, and have similar potential for Mars.

We used 1–2 m/pixel digital elevation models derived from 25–50 cm/pixel High Resolution Imaging Science Experiment stereo-pair images to measure the planform and 3D morphometries of the mid-latitude Martian eskers, and compare them with the morphometries of Quaternary-aged eskers in Canada [4] and SW Finland [5]. We found that the Martian eskers have remarkably similar lengths, sinuosities and heights to terrestrial eskers, but that the Martian eskers are typically wider and have lower side slopes. Large width-height ratios of the Martian eskers are consistent with our previous measurements of ancient (~3.5 Ga) eskers close to Mars' south pole [6]. These large ratios may arise from differences in either esker degradation state or fundamental glacio-hydrological controls on esker formation between Mars and Earth.

We also used a novel morphometric approach to test evidence for spatio-temporal variations in sediment-discharge dynamics during formation of the Martian esker in Tempe Terra. Our analyses reveal that this esker has a 'stacked' morphology: the crest of a wide, round-crested underlying ridge is superposed by a narrow, sharp-crested to multi-crested ridge. The superposed ridge is aligned with a channel-like depression that incises the underlying ridge. Based on morpho-sedimentary relationships observed along terrestrial eskers [7], we interpret this stacked morphology to represent conduit adjustment as a result of waning meltwater discharge and sediment supply towards the end of the esker-forming drainage episode(s).

This work was funded by STFC grant ST/N50421X/1.

References: [1] Gallagher, C., and Balme, M.R., (2015), *Earth. Planet. Sci. Lett.* 431, 96-109, [2] Butcher, F.E.G., et al. (2017), *J. Geophys. Res. Planets.* 122(12), 2445-2468, [3] Butcher, F.E.G., et al. (2019), This Conference, [4] Storrar, R.D., et al. (2014) *Quat. Sci. Rev.* 105, 1-25, [5] Storrar, R.D., and Jones, A., Unpublished, [6] Butcher, F.E.G., et al. (2016), *Icarus* 275, 65-84, [7] Burke, M.J., et al. (2010) *Geol. Soc. Am. Bull.* 122, 1637-1645.