

Open Research Online

The Open University's repository of research publications and other research outputs

Modelling the survival of ejected martian biomarkers impacting Phobos

Conference or Workshop Item

How to cite:

Morland, Zoe; Halim, Samuel; Pearson, Victoria; Patel, Manish; Green, Simon and Ramkissoon, Nisha (2020). Modelling the survival of ejected martian biomarkers impacting Phobos. In: EPSC Abstracts, Europlanet Science Congress 2020, article no. 254.

For guidance on citations see [FAQs](#).

© 2020 The Authors



<https://creativecommons.org/licenses/by/4.0/>

Version: Version of Record

Link(s) to article on publisher's website:

<https://meetingorganizer.copernicus.org/EPSC2020/EPSC2020-254.html>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk



Modelling the survival of ejected martian biomarkers impacting Phobos

Zoe Morland¹, Samuel Halim², Victoria Pearson¹, Manish Patel¹, Simon Green¹, and Nisha Ramkissoon¹

¹AstrobiologyOU, School of Physical Sciences, The Open University, Milton Keynes MK7 6AA, United Kingdom of Great Britain and Northern Ireland (zoe.morland@open.ac.uk)

²Department of Earth and Planetary Sciences, Birkbeck, University of London, Malet St, London WC1E 7HX, United Kingdom of Great Britain and Northern Ireland

Motivation and Approach

Phobos' proximity to Mars and short orbital period has led to the hypothesis that Phobos could sweep up particles ejected from large impacts on the martian surface [1]; models suggest that Phobos' regolith could include up to ~250 ppm of martian ejecta material [2,3]. Considering that Mars' surface has many "Special Regions" that could have been habitable in the past [4], it is not unreasonable to suggest that life could have developed and left behind biomarkers. An impact into one of these areas could eject material containing biologically-significant compounds and deposit it on the surface of Phobos. Therefore, an appealing possibility is that samples collected from the surface of Phobos, by missions such as JAXA's Martian Moons eXploration (MMX) [5], could contain martian biomarkers. Further investigation into the feasibility of biomarker transfer from Mars to Phobos is hence necessary before returned samples and *in situ* spacecraft data are analysed.

In order to investigate this scenario, light gas gun impact and heating experiments will be used to simulate the conditions a martian rock, containing biologically-relevant material, would undergo throughout the transport process from Mars to Phobos [6,7]. However, prior to carrying out these experiments, the biologically-relevant material must be defined. It is important that this material represents plausible molecular fingerprints of the types of organisms that might have existed in the harsh radiation conditions of Mars' near-surface for billions of years [8]. Furthermore, they must survive the shock conditions experienced during ejection by a sizeable impact capable of delivering material to reach Phobos' orbit, and the subsequent deposition onto the surface of Phobos. Biomarkers that could satisfy these criteria include amino acids, polycyclic aromatic hydrocarbons (PAHs), fatty acids and sterols [e.g. 9], some of which may be precursors of chlorinated molecules detected in the Cumberland mudstone at Gale Crater [10].

Previous impact modelling has suggested that a small proportion of near-surface martian material can be ejected at high velocities, capable of reaching Phobos from Mars, and experience shock-pressures low enough to remain solid [11]. However, simulations of martian ejecta impacting Phobos are limited to mass-less [2,3] or "hard rock" projectiles [12] that neglect the reduced shock experienced at the trailing edge of a projectile during the early stages of contact and compression [13], which could aid biomarker survival [14]. Therefore, this study uses the iSALE-2D shock-physics code [15-17] to estimate the pressure and temperature regimes within martian ejecta as it impacts Phobos' surface. This will enable constraints to be placed on conditions necessary for

biomarker survival.

Model Parameters

Consistency between *in situ* [e.g. 18, 19] and remote sensing [e.g. 20] observations, and the global meteorite collection [21], indicate that Mars' global surface composition can be approximated as basaltic. Thus, it can be assumed that basalt is the most likely crustal rock composition to be ejected from Mars. However, this igneous composition is not the most likely rock type in which to find biomarkers. Sedimentary rocks, such as mudstones, indicate favourable past conditions for habitability on Mars and have an increased potential to preserve biomarkers [22]. Organics and bioessential elements have been detected within mudstones at Gale crater [e.g. 23, 24]. However, mudstones are spread sparsely over Mars' surface, so are less likely to be ejected by impacts. This work, therefore, considers both igneous and sedimentary ejecta, modelling basalt and mudstone projectiles.

For the target, the surface composition of Phobos is largely unknown, owing to the lack of direct sampling from the surface; spectral data indicate that the surface resembles D- or T-type asteroids or carbonaceous chondrites, rich in phyllosilicates [25]. Therefore, the closest match to Phobos' compositional and physical properties will be chosen for the target, with reference to Phobos regolith simulants [e.g. 26].

The simulations will be run over a range of realistic impact parameters, including impactor size ($\sim 0.01 - 10$ m [27]) and impact velocity (up to 5.3 km s^{-1} [3,12]).

Implications

The results from this modelling will provide insight into the survival of martian biomarkers during deposition onto Phobos. Furthermore, the results will inform the subsequent impact and heat experimental simulations.

References

- [1] Murray, J. B., (2011) *EPSC-DPS Joint Meeting*, 1003. [2] Ramsley, K. R. & Head, J. W., (2013) *Planet. & Space Sci.*, 86, 115-129. [3] Chappaz, L. et al., (2013) *Astrobiology*, 13(10), 963-980. [4] Rummel, J. D., et al., (2014) *Astrobiology*, 14(11), 887-968. [5] Usui, T. et al., (2018) *42nd COSPAR Scientific Assembly*, B4.2-7-18, 2018. [6] Morland, Z. S. et al., (2020) *Proc. 51st Lunar Planet. Sci. Conf.*, 1096. [7] Morland, Z. S. et al., (2019) *Proc. British Planet. Sci. Conf.*, 88. [8] Kminek, G. et al., (2006) *Earth Planet. Sci. Lett.*, 245(1-2), 1-5. [9] Vago, J. L. et al., (2018) in B, Cavalazzi, and F, Westall (Eds.) *Springer*, 283-300. [10] Szopa, C., et al., (2020) *Astrobiology*, 20(2), 292-306. [11] Artemieva, N. & Ivanov, B., (2004) *Icarus*, 171, 84-101. [12] Summers, D., (2019) *Life Sci. Space Res.*, 23, 101-111. [13] Pierazzo, E. & Melosh, H. J., (2000) *Meteorit. Planet. Sci.*, 35, 117-130. [14] Pierazzo, E. & Chyba, C. F., (1999) *Meteorit. Planet. Sci.*, 34, 909-918. [15] Amsden, A. et al., (1980) *LANL Report*, LA-8095. [16] Collins, G. S. et al., (2004) *MAPS*, 38, 217-231. [17] Wünnemann, K. et al., (2006) *Icarus*, 180, 514-527. [18] Filiberto J., (2017) *Chemical Geology*, 466, 1-14. [19] Cannon, K. M. et al., (2019) *Icarus*, 317, 470-478. [20] McSween Jr. H. Y. et al., (2003) *JGR Planets*, 108(E12), 5135. [21] Martian Meteorite Compendium, <https://curator.jsc.nasa.gov/antmet/mmc/> [22] Hays, L. E. et al., (2017) *Astrobiology*, 17(4), 363-400. [23] Stern, J. C. et al., (2015) *Proc. Nat. Acad. Sci.*, 112(14), 4245-4250. [24] Eigenbrode, J. L. et al., (2018) *Science*, 360(6393), 1096-1101. [25] Fraeman, A. A. et al., (2014) *Icarus*, 229, 196-205. [26] Morland, Z. S., (2019) *Proc. 50th Lunar Planet. Sci. Conf.*, 2132. [27] Ramsley, K. R. & Head, J. W., (2013) *Planet. & Space Sci.*, 75, 69-95.