



## Open Research Online

### Citation

Heinze, Gerhard; Sartorius, Norman; Guizar Sanchez, Diana Patricia; Bernard-Fuentes, Napoleón; Cawthorpe, David; Cimino, Larry; Cohen, Dan; Lecic-Tosevski, Dusica; Filipcic, Igor; Lloyd, Cathy; Mohan, Isaac; Ndetei, David; Poyurovsky, Michael; Rabbani, Golam; Starostina, Elena; Yifeng, Wei and EstefaníaLimon, Limón (2021). Integration of mental health comorbidity in medical specialty programs in 20 countries. *The International Journal of Psychiatry in Medicine*, 56(4) pp. 568–581.

### URL

<https://oro.open.ac.uk/76146/>

### License

(CC-BY-NC-ND 4.0) Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

<https://creativecommons.org/licenses/by-nc-nd/4.0/>

### Policy

This document has been downloaded from Open Research Online, The Open University's repository of research publications. This version is being made available in accordance with Open Research Online policies available from [Open Research Online \(ORO\) Policies](#)

### Versions

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding

## The International Journal of Psychiatry in Medicine

### Integration of mental health comorbidity in medical specialty programs in 20 countries

Journal:	<i>The International Journal of Psychiatry in Medicine</i>
Manuscript ID	IJPM-20-0291.R1
Manuscript Type:	Original Research Article
Keywords:	Curriculum development, Medical education, Comorbidity, Medical Illness, Mental Illness
Abstract:	<p>Methods:A systematic analysis was performed of the medical specialization academic programs of 20 different countries to establish which ones take into account mental health comorbidity.The criteria that were explored in the educational programs include:1)name of the medical specialties that take into account mental health comorbidity in curriculum design,2) name of the mental health content addressed by these programs. After independent review and data extraction, paired investigators compared the findings and reached consensus. Descriptive statistics evaluated the frequency of the data presented.</p> <p>Results:Internal medicine, family medicine, neurology, pediatrics and geriatrics were the specialties that included mental health topics in their programs. In four countries: Bangladesh, Serbia, the Netherlands and France, 50%of all graduate specialty training programs include mental health content. In ten countries:Germany, Sweden, the United Kingdom, Mexico, Belgium, India, Russia, Canada, Israel and Spain, between 20%and 49%of all graduate specialty training programs include mental health content. In six countries(Brazil, Chile, Colombia, Croatia, Kenya, and the United States)less than 20%of all graduate specialty training programs include mental health content.</p> <p>Discussion:The proposal that we have made in this article should be taken into account by decision-makers, in order to complement the different postgraduate training programs with mental health comorbidity that are frequently present with other physical symptoms.It is not our intention that the different specialists know how to treat psychiatric comorbidities, but rather pay attention to their existence and implications in the diagnosis, evolution and prognosis of many other diseases.</p>

SCHOLARONE™  
Manuscripts

## Integration of mental health comorbidity in medical specialty programs in 20 countries

### ABSTRACT

**Methods:** A systematic analysis was performed of the medical specialization academic programs of 20 different countries to establish which medical specialties take into account mental health issues in the specialty curricular design and which mental health content these programs address. The criteria that were explored in the educational programs include: 1) name of the medical specialties that take into account mental health content in curriculum design, 2) name of the mental health issues addressed by these programs. After independent review and data extraction, paired investigators compared the findings and reached consensus on all discrepancies before the final presentation of the data. Descriptive statistics evaluated the frequency of the data presented.

**Results:** Internal medicine, family medicine, neurology, pediatrics and geriatrics were the specialties that included mental health topics in their programs. In four countries: Bangladesh, Serbia, the Netherlands and France, 50% of all graduate specialty training programs include mental health content. In ten countries: Germany, Sweden, the United Kingdom, Mexico, Belgium, India, Russia, Canada, Israel and Spain, between 20% and 49% of all graduate specialty training programs include mental health content. In six countries - Brazil, Chile, Colombia, Croatia, Kenya, and the United States - less than 20% of all graduate specialty training programs include mental health content.

**Discussion:** The proposal that we have made in this article should be taken into account by decision-makers, in order to complement the different postgraduate training programs with mental health issues that are frequently present with other physical symptoms. It is not our intention that the different specialists know how to treat psychiatric comorbidities, but rather pay attention to their existence and implications in the diagnosis, evolution and prognosis of many other diseases. The current fragmentation of medicine into ever finer specialties makes the management of comorbidity ever more difficult: a reorientation of post-graduate training might improve the situation.

## 1.- Introduction

The management of comorbidity and emerging diseases are the main challenges for medicine in the 21st century. The first requires, as a first step, changes in the organization of health services and in the training of health workers. While the second should be addressed by improving research aimed at the prevention and treatment of emerging diseases

The increasing prevalence of chronic illnesses has been accompanied by a concomitant increase in comorbidity (co-occurrence of two or more chronic health conditions). The success in prolonging life expectancy did not greatly increase the number of disease-free years: the years gained are years in which people tend to suffer from a variety of chronic diseases, impairments and disabilities. The simultaneous presence of comorbid diseases usually predicts a poorer prognosis for both conditions; it can lead to a greater number of complications, and makes treatment more complex<sup>1-2</sup> and expensive<sup>3</sup>. It is important to point out the difference between comorbidity as the start of another clinical entity in the course of an existing disease, and multimorbidity, defined as the coexistence of several diseases with varying degrees of association in the same individual<sup>4</sup>.

In 2019, according to the Global Burden of Disease Collaborators, mental disorders are among the top ten factors that produced the highest number of disability-adjusted life years in the world in the last 30 years<sup>5</sup>. Despite the fact that diseases such as ischemic heart disease, diabetes, stroke, chronic kidney disease and lung cancer are in the first places; mental disorders are often comorbid with such noncommunicable diseases, which worsens their prognosis and the increasing costs associated with their treatment. In addition to the fact that people with mental disorders live 10 years less than people without mental disorders, and non-communicable diseases are the leading cause of their death.

Adults with medical multimorbidity, usually defined as  $\geq 2$  concurrent chronic conditions – for example cardiovascular disease and diabetes – (<sup>6-7</sup>, have high rates of healthcare utilization and associated costs<sup>8</sup>.

The comorbidity of mental-physical diseases has become a major problem in medical practice, as it is present in approximately 30% of individuals<sup>9</sup>. In persons over the age of 60, comorbidity has become the rule rather than an exception<sup>10-11</sup>. Scott et al<sup>12</sup> analyzed 17 countries from the World Mental Health survey and reported a consistent pattern of associations between mental disorders and subsequent development of chronic medical

1  
2  
3 conditions, with odds ratios ranging from 1.2 (95% CI, 1.0-1.5) to 3.6 (95% CI, 2.0-6.6).  
4 Recent research indicates a bidirectional relationship with common risk factors (eg, adverse  
5 experiences in early childhood) and shared etiopathogenesis, and the results of the others  
6 impact through neurohormonal and behavioral pathways among some of the  
7 noncommunicable diseases and mental disorders (eg, depression). The pathways linking  
8 mental and physical comorbidity are complex and bidirectional such that, a physical disorder  
9 may cause a mental one<sup>12-14</sup>, a mental disorder may place a person at risk for a physical one,  
10 and some mental and physical disorders share risk factors, such as chronic stressful social  
11 life situations, physical inactivity, overweight, smoking, substance use disorders and  
12 endocrine dysregulation. Additionally, social conditions like poverty and adverse life events  
13 are risk factors for both types of conditions, perhaps due to biological mechanisms and an  
14 unfavorable environment.  
15

16  
17 The relationship of mental-physical disorders holds the potential for a population health index  
18 that can gauge the effect of prevention programs at the community level. A study by  
19 Cawthorpe found the median cost of physical disorders to be 2.9 times higher in the group  
20 with any International Classification of Diseases psychiatric disorder, in comparison with the  
21 group without psychiatric disorder<sup>15</sup>. In the USA, a major study of 30,022 adults showed that  
22 the risk of functional disability in people with diabetes was 2.42 times higher than in people  
23 without diabetes; that in people with depression alone it was 3 times higher than in people  
24 without depression; and that the risk for those who had depression and diabetes, the risk was  
25 7.15 times higher than for people who had neither diagnosis<sup>16</sup>. When depression is present,  
26 the cost of treating diabetes is 4.5 times higher<sup>17</sup>.  
27

28  
29 Living with medical comorbidity often includes taking multiple prescription medications and  
30 careful management of individual illnesses. Comorbid substance use disorder (SUD) can  
31 have profound negative effects on pre-existing chronic diseases, since different substances  
32 of abuse have been associated with a wide range of physiological effects (cardiovascular,  
33 gastrointestinal, hematological, pulmonary and neurological) that complicate the evolution,  
34 clinical management and prognosis of chronic diseases. Furthermore, many adults who are  
35 in treatment for SUD receive fragmented medical care for their other chronic conditions<sup>18</sup>.  
36

37  
38 Chronic medical diseases, mental illness, and substance use frequently are interrelated and  
39 represent a compound multimorbidity placing individuals at even higher risk for adverse  
40

1  
2  
3 outcomes including increases in healthcare utilization. Therefore, such comorbidity must be  
4 taken into account when developing effective care for medical comorbidity<sup>19</sup>.

5  
6 While the association between physical and mental disorders continues to grow, medicine is  
7 fragmenting into increasingly limited specialties. There are few residency programs that train  
8 in evaluation and assessment of mental disorders (like internal medicine) and their training is  
9 often inadequate or incomplete<sup>20</sup>. While clinicians may be aware of comorbidity, they may  
10 also feel pressed for time and powerless to handle problems, as well as lacking confidence  
11 or strategies to address all problems. Training should take into account that doctors can  
12 control and limit the interview as a way to avoid emotional difficulties. In other words,  
13 physicians increasingly treat problems and / or disorders in which they received little training  
14 during residency.

15  
16 A solution from medical education consists of developing and implementing a comprehensive  
17 curriculum in physical and mental comorbidity during training as medical specialists<sup>20</sup>. The  
18 context of the current pandemic, followed by extensive restrictive public health measures  
19 (such as isolation, quarantine, and monitoring) can cause significant distress resulting in the  
20 emergence of mental disorders such as depression in previously unaffected populations.  
21 Therefore, the current context demands a call for a significant change in the education of  
22 residents in practical skills to manage common difficulties of patients, such as anxiety,  
23 depression, substance abuse and chronic pain<sup>21</sup>. In addition, there is a gap in the literature  
24 related to the evidence for the integration of medico-mental comorbidity into the medical  
25 residency program and how these curricula can meet training needs<sup>22</sup>.

26  
27 The only way to ensure an appropriate attitude towards comprehensive patient care, in any  
28 medical specialty, is to work in a cooperative and interdisciplinary manner. Therefore, the  
29 training must incorporate some explanation of the comorbidity and the potentially dangerous  
30 implications of not identifying it within the consultation.

31  
32 The biopsychosocial model of health care, awareness of comorbidity (physical-mental)  
33 inclusive approaches on promotion and prevention aspects in mental health, doctor-patient  
34 relationship and ethics should be included in the different training programs.

35  
36 The current fragmentation of medicine into increasingly fine specialties makes the  
37 management of comorbidity even more difficult: a reorientation of postgraduate training could  
38 improve the situation. In our current context, a reorientation in the curriculum is urgent, since  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 health care providers have an essential role in the treatment of emotional outcomes as part  
4 of the response to the pandemic COVID-19 and control measures (eg quarantine and  
5 distancing Social). Future specialists need to acquire sufficient knowledge to diagnose and  
6 treat (or refer) common psychiatric diseases. The main objective of this study was to establish  
7 which medical specialties take mental health into account in the design of their training  
8 programs, and what mental health topics they address.  
9  
10  
11  
12  
13  
14  
15  
16

## 17 2.- Methods.

18 This document analysis systematically analyzed the medical specialization academic  
19 programs of 20 different countries. The searching criteria that were explored in the  
20 educational programs include:  
21  
22

- 23 1. Which specialty programs addressing mental health comorbidity in their training  
24 program?
- 25 2. How many specialties include the following mental health comorbidity in their regular  
26 postgraduate program: affective disorders, anxiety disorders, substance use disorders,  
27 and alcohol use disorders?  
28
- 29 3. What other mental health topics are included in the different medical specialty  
30 programs?  
31  
32  
33  
34  
35

36 Operational definition. Mental health content was defined as the specific mention in the  
37 curriculum of each academic program of depressive, anxiety, substance use disorders  
38 comorbid with medical pathology. As well as the express mention of the study of the doctor-  
39 patient relationship in the practice of the specialty.  
40  
41  
42  
43

### 44 2.1. Document collection

45 For each of the 20 countries: 1. all medical schools in the country were searched, 2.- once all  
46 schools were identified by country, all graduate programs associated with each school were  
47 identified, and 3.- then their study plan for each of the programs was analyzed.  
48  
49

50 Study plan and medical specialty programs were requested through invitations, searches on  
51 institutional websites and communications from all medical schools in the participating  
52 countries. For countries with multiple medical schools, each website was independently  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 searched for relevant documents. To be included in the analysis, documents had to be  
4 retrievable from an institution. Otherwise, schools were excluded from the study.  
5  
6  
7  
8  
9

## 10 2.2 Data extraction process

11 The data extraction form was created in Excel and was designed to extract documented  
12 information. The majority of items on the form appeared as check- boxes to indicate the  
13 presence or absence of criteria. Those related to the proposed criteria were extracted for  
14 analysis. The initial draft of the data extraction form was created by three coauthors (GH,  
15 DPGS, NB), and was subsequently reviewed and revised by four groups of four investigators  
16 (16 investigators in total) extracted data from the available papers using the final form. After  
17 independently extracting the data, each group of investigators compared the entries, resolved  
18 discrepancies by consensus, and submitted a final data extraction form for each medical  
19 school reviewed.  
20  
21  
22  
23  
24  
25  
26

27 Data were organized and analyzed using IBM SPSS statistical software, version 26 (IBM  
28 Corporation, New York, NY, USA). Descriptive statistics evaluated the frequency of the data  
29 presented.  
30  
31  
32  
33

## 34 3.- Results

35 The materials describing 1036 postgraduate specialty programs that were examined showed  
36 that internal medicine, family medicine, neurology, pediatrics and geriatrics were the  
37 specialties that included mental health topics in their programs with some differences between  
38 the 20 countries, for example: Neurology include mental health comorbidity on 17 different  
39 countries vs Geriatrics that include mental health comorbidity on 8 different countries (See  
40 Figure 1).  
41  
42  
43  
44  
45  
46

47 INSERT FIGURE 1  
48  
49  
50

51 The percentages of mental health content found in the postgraduate programs from the 20  
52 countries included in this survey were: 1) In four countries: Bangladesh, Serbia, the  
53 Netherlands, and France, over 50% of all postgraduate specialty training programs included  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 mental health content; 2) In ten countries: Germany, Sweden, United Kingdom, Mexico,  
4 Belgium, India, Russia, Canada, Israel, and Spain, between 20%-49% of all postgraduate  
5 specialty training programs included mental health content; 3) In six countries: Brazil, Chile,  
6 Colombia, Croatia, Kenya, and the United States, 20% of all postgraduate specialty training  
7 programs included mental health content. (See Figure 2)  
8  
9

10  
11  
12 INSERT FIGURE 2  
13  
14

15 Croatia and Russia do not have medical specialties that include affective disorders in their  
16 academic programs. Belgium, Brazil and Kenya do not have medical specialties that include  
17 psychosomatic disorders in their academic programs, whereas Germany, France and Croatia  
18 do not include anxiety disorders in their academic specialty programs. Bangladesh, Mexico,  
19 the Netherlands, Israel, Serbia, and Spain include mental health issues in more than 10 of  
20 their medical specialty programs (see Figure 3).  
21  
22  
23  
24

25  
26 INSERT FIGURE 3  
27  
28

29 Brazil, Croatia and Russia do not have medical specialties that include substance use  
30 disorders (SUD) or alcohol use disorder (AUD) in their academic programs. None of the  
31 medical specialty programs in Kenya include AUD. Belgium, Chile, Colombia, England,  
32 France, Germany, and the United States do include AUD and SUD, but only in less than five  
33 medical specialty programs. (See Figure 4)  
34  
35  
36

37  
38 INSERT FIGURE 4  
39

40 In Serbia, Mexico, Bangladesh, the Netherlands, Russia, England, France, Sweden, and  
41 Spain, more than 10 different programs address other mental health topics, such as the  
42 doctor-patient relationship, violence, medical ethics, medical responsibility,  
43 neuropsychological syndromes, etc.  
44  
45  
46  
47

#### 48 4.- Discussion

49 It is of great concern that only four of the 20 countries in our study contemplate mental health  
50 in over 50% of their postgraduate specialty training programs, while this percentage falls  
51 below 20% in six countries.  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 In recent years, the increasing prevalence of chronic, often comorbid conditions requires that  
4 patients receive care from multiple providers in a variety of settings. Intensified specialization  
5 has sought to generate greater interdependence among clinicians and promote cross-  
6 fragment coordination, however, it has exacerbated fragmentation by increasing the number  
7 of narrowly trained specialists. "Fragmentation" in healthcare services means a systemic lack  
8 of coordination that spawns inefficient allocation of material and human resources.  
9 Fragmentation adversely impacts quality, cost, and outcomes<sup>14</sup>. The healthcare system is  
10 suffering from what George Halvorson calls "clinical linkage deficiencies"<sup>23</sup>. This systemic  
11 fragmentation is difficult to dislodge, because it is steeped in the history and culture of  
12 medicine and embedded population-wide within the current system — operationally,  
13 financially, and in the clinic.

14  
15 The comorbidity of a mental and physical condition may impact the mode of presentation,  
16 clinical severity, response to treatment, and burden of illness of both conditions. New insights  
17 in biology, physics, human organization, and other fields have led to understanding complex  
18 systems as more than the sum of their parts<sup>24</sup>. However, because of our fragmented  
19 understanding of the natural world, systems, and human interactions, health care has not  
20 kept up with these advances<sup>25</sup>. Specialized information has expanded without an  
21 accompanying expansion in our ability to integrate, prioritize, and personalize narrowly  
22 construed information.

23  
24 Fragmentation often involves multiple providers and organizations, with deficient coordination  
25 among the parties, that may lead to suboptimal care, including important healthcare issues  
26 being inadequately addressed, poor patient outcomes, and unnecessary or even harmful  
27 services that ultimately both raise costs and degrade quality<sup>26-27</sup>.

28  
29 Research evidence consistently demonstrates that people with chronic diseases are two to  
30 three times more likely to experience mental health problems than the general population<sup>18</sup>.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

Research evidence consistently demonstrates that people with chronic diseases are two to three times more likely to experience mental health problems than the general population<sup>18, 28-29</sup>. As a result of this comorbidity, both the prognosis for their chronic condition and the quality of life they experience can deteriorate markedly. In addition, the costs of providing care for this group of people are increased as a result of less effective self-care and other complicating factors related to poor mental health. The mechanisms underlying this type of comorbidity are complex, and evidence suggests that a combination of biological, psychosocial, environmental, and behavioral factors may be at play<sup>28</sup>.

1  
2  
3 Medical specialization (ME) plays a key role in contemporary medicine and the role of the  
4 specialist should not be questioned. ME emerged as a result of the shift to pathological  
5 anatomy (medical model divides the body into parts in which different types of doctors have  
6 to specialize), the emergence of medical technologies, and due to urbanization.  
7  
8 Specialization is a way to improve healthcare outcomes, by applying greater knowledge and  
9  
10 experience on specific diagnostic and therapeutic problems. This represented important  
11  
12 advances in the treatment of diseases previously considered incurable, as well as in  
13  
14 increasing life expectancy. Adequate communication between specialists is essential to avoid  
15  
16 distancing from integrated medicine  
17

18  
19 In particular, cardiology, endocrinology, pneumology, infectiology, rehabilitation medicine,  
20  
21 and orthopedics do not include mental health content in their academic programs. In the  
22  
23 majority of the surveyed countries, there is strong clinical evidence that these specialties may  
24  
25 benefit from teaching mental health topics to postgraduate residents, given their close  
26  
27 relationship with cardiovascular diseases, diabetes, chronic obstructive pulmonary disease  
28  
29 (COPD), and chronic musculoskeletal disorders<sup>30-36</sup>.

30  
31 Cardiovascular prognosis is less favorable for patients with comorbid mental health diseases,  
32  
33 for instance, an association with depression increases acute exacerbations per year by 50%,  
34  
35 as well as mortality rates<sup>31-34</sup>. People with comorbid diabetes-depression have an increased  
36  
37 risk (> 20%) of unfavorable health outcomes and premature mortality<sup>30, 34</sup>. Mental illness can  
38  
39 also have a major impact on symptoms and outcomes for people with COPD (regardless of it  
40  
41 severity): psychiatric comorbidity in these patients is associated with poorer health status,  
42  
43 increased breathlessness, and an increase in mortality rates<sup>30, 36-40</sup>. There is also evidence  
44  
45 for a higher prevalence of mental disorders among people with other conditions, including  
46  
47 asthma, arthritis, cancer, HIV/AIDS, and others<sup>37,38</sup>.

48  
49 Finally, it is also important to take into account child maltreatment (CM) as the most important  
50  
51 preventable cause of psychopathology. There is a palpable relationship between adversity  
52  
53 early in life and adult health status<sup>38</sup>.

54  
55 Studies show that comprehensive care that avoids diagnostic-therapeutic fragmentation  
56  
57 improves a patient's evolution, prognosis and quality of life<sup>4, 9-10</sup>. Appropriate management of  
58  
59 mental and somatic comorbidity, at the individual and at the public health level, will require a  
60  
significant reorientation of medical education.

1  
2  
3 Psychiatrists are no better than other specialists at identifying comorbid conditions. This is  
4 most clearly demonstrated in the creation of a sub-discipline called liaison psychiatry. No  
5 other medical discipline has a similar subspecialty; there is no liaison cardiology or  
6 pulmonology because any specialist is expected to deal with the physical problems of their  
7 patients. Psychiatrists often overlook or under-treat a comorbid physical illness, omitting a  
8 medical examination that could inform them of the presence of another illness. Given that  
9 cardiovascular, cancer, neonatal and musculoskeletal disorders rank above mental health  
10 morbidity statistics worldwide, it is expected that in the future psychiatrists will acquire  
11 sufficient knowledge to diagnose and treat (or refer) non-psychiatric diseases and that,  
12 therefore, the discipline liaison psychiatry is unnecessary. We are still far from this goal. The  
13 academic training program of a psychiatrist must be based on an epidemiological profile and  
14 include the most common chronic degenerative diseases. A future study regarding the  
15 inclusion of physical health content in psychiatric training would be interesting in the future.  
16 Worldwide efforts are needed to convince decision makers, educators, clinicians, healthcare  
17 institutions, and insurance providers, not only that mental-physical comorbidity is not an  
18 exception but nearly a rule, but also that it constitutes one of the most urgent challenges for  
19 public health authorities.

20  
21 Not only should undergraduate education be a target for the objective of integration, it is  
22 therefore unavoidable that specialists in all disciplines also assume some responsibility for  
23 dealing with the issue of mental comorbidity in the patients whom they treat.

24  
25 Education and training on psychosocial issues should be provided to health system leaders  
26 and health professionals on the front line of COVID care. Some patients will need a referral  
27 for formal mental health assessment and treatment, in some cases immediate (suicidality),  
28 while others may benefit from supportive interventions designed to promote well-being and  
29 improve coping. In addition, there should be no substantial differences in the treatment of  
30 COVID-19 infection between patients with a previous diagnosis of a mental disorder and  
31 those without a diagnosis of mental disorder.

## 51 5.- Conclusion

52  
53 Comorbidity is a term that has been borrowed from physical medicine. Its original meaning  
54 indicated the presence of at least two diseases. The co-occurrence of these diseases  
55  
56  
57  
58  
59  
60

1  
2  
3 complicates diagnosis and treatment; and raises questions about the etiology and the mutual  
4 interdependence of diseases<sup>36, 39</sup>. This in turn calls for the involvement of different medical  
5 specialties in patient care, in order to clarify the questions raised and provide comprehensive  
6 treatment.  
7  
8  
9

10 The challenges of responding effectively to compounded mental and physical health needs  
11 are part of a larger problem regarding the care for people with comorbidities of any kind.  
12 Current service models are often orientated around single diseases and fail to provide well-  
13 coordinated care to a great and growing number of people with multiple health problems.  
14  
15

16 For physicians it must be fundamental to treat each patient as a whole. At present, however,  
17 specialist doctors tend to focus only on their particular area of study, making it impossible to  
18 care adequately for patients who have other types of health problems. It thus becomes  
19 especially important for primary care physicians to understand the bases of attention in  
20 psychiatry, as they are the first point of contact for patients. Knowing how to identify  
21 psychiatric pathology would enable primary care physicians to detect it opportunely, prescribe  
22 initial treatment, and adequately refer cases that need more specialized attention.  
23  
24  
25  
26  
27  
28

29 No method can be developed to reliably determine the quality of psychiatric education and  
30 training worldwide because each country has its own epidemiology of mental illness.  
31 However, it is recommended that the institutions that regulate postgraduate curricula in each  
32 country consider the incidence and prevalence of these diseases in order to include them in  
33 the curricula of different specialties, paying special attention to psychiatric education for  
34 residents of primary care medicine. It is desirable to include mental health topics, as well as  
35 rotations in the psychiatric clinic area<sup>36</sup>, in postgraduate curricula as befits the needs of each  
36 country, so that psychiatrists and other specialists share the responsibility of detecting and  
37 treating mental illness opportunely.  
38  
39  
40  
41  
42  
43

44 Some specific mental health topics could be added to the curriculum in many countries  
45 carefully according to epidemiological needs, avoiding their impact on other aspects of the  
46 program. The curriculum is a zero-sum game, which means adding one thing may require  
47 removing another.  
48  
49  
50

51 Awareness of mental health must be integrated into all aspects of health and social policy,  
52 into the planning of the healthcare system and into the provision of primary & secondary  
53 healthcare. The proposal is to integrate (within medical training programs) common mental  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 health problems in the routine care of people affected by other chronic non-communicable  
4 diseases (such as cancer, diabetes and cardiovascular disorders), HIV / AIDS and maternal  
5 health care. Integrating physical and mental comorbidity into medical education programs  
6 offers the opportunity to treat " the whole patient ", with a better overall cost-effectiveness  
7 than an approach in which mental health, acute physical health and Chronic, reproductive  
8 health problems, infectious and chronic pain are addressed without effective knowledge and  
9 communication among health professionals. An integrated care model is often more attractive  
10 to patients and family members who are concerned about the stigma that is still associated  
11 with mental and substance abuse disorders.

12  
13 The proposal of the present article is to complement the pre-existing programs with  
14 annotations in mental health, since it is not intended that the different specialist doctors know  
15 how to treat all psychiatric diagnoses, but that they know their existence, implications in the  
16 diagnosis and prognosis of other diseases, as well as the changes in the clinical evolution  
17 generated by the medical and mental comorbidity.

18  
19 An integrated medical education approach to address mental health in for example: the  
20 context of HIV care, maternal mental health, and chronic non-communicable diseases is  
21 based on the conviction and growing evidence of the efficiency, effectiveness, and cost  
22 savings of an integrated medical care<sup>41</sup>. Prevention and early intervention also reduce the  
23 global burden of mental and physical comorbidity. An integrated approach that seeks to  
24 prevent conditions that affect mental and physical health would share common strategies; for  
25 example, reducing alcohol consumption and smoking and promote physically active  
26 lifestyles<sup>42</sup>.

27  
28 Finally, we must not forget that the mental health implications of the COVID-19 pandemic  
29 (protecting people with mental disorders from COVID-19, the anticipated increase in mental  
30 disorders throughout the population, and the neuropsychiatric consequences of the disease)  
31 require a greater focus on public interventions mental health. The notion that only  
32 psychiatrists can provide treatment must be abandoned. Other forms of collaborative and  
33 community-based approaches are required, through training of the general practitioner and  
34 specialists.

## 6.- Contributors

GH, NS, DGS, NBF were responsible for the literature search, figures, study design, data collection, data analysis, data interpretation, and writing. The rest of the authors contributed equally to data collection, data interpretation, and writing.

## 7.- Conflicts of interest

All the authors declare no conflicts of interest.

## 9.- References

1. Kissane DW, Maj M, Sartorius N. Depression and Cancer. 2011. Chichester, UK: Wiley-Blackwell.
2. Kurrle S, Brodaty H, Hogarth R. Physical Comorbidities of Dementia. 2012. Cambridge, UK: Cambridge University Press.
3. Sartorius N. Comorbidity of mental and physical disorders: a key problem for medicine in the 21st century. *Act Psychiatr Scand.* 2018; 137:369–70.
4. Barnett K, Mercer S, Norbury M, Watt G, Wyke S, Guthrie B. Epidemiology of multimorbidity and implications for health care, research, and medical education. *Lancet.* 2012; 380(9836):37–43.
5. Collaborators, G.B.D.N. Global, regional, and national burden of neurological disorders, 1990-2016: A systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol.* 2019, 18, 459–480.
6. Fortin M., Stewart M., Poitras M.-E., Almirall J., Maddocks H. A systematic review of prevalence studies on multimorbidity: toward a more uniform methodology. *Ann. Fam. Med.* 2012;10(2):142–151.
7. Wallace E., Salisbury C., Guthrie B. Managing patients with multimorbidity in primary care. *BMJ.* 2015;350:h176.
8. Lochner K.A., Cox C.S. Prevalence of multiple chronic conditions among Medicare beneficiaries, United States, 2010. *Prev. Chronic Dis.* 2013;10
9. Sartorius N. Comorbidity of mental and physical diseases: a main challenge for medicine of the 21st century. *Shanghai Arch Psychiatry.* 2013; 25(2):68-69.
10. Naylor C, Parsonage M, McDaid D, Knapp M, Fossey M, Galea A. Long-term conditions and mental health: the cost of co-morbidities. London: The King's Fund and Centre for

- 1  
2  
3 Mental Health. 2012. [accessed 2019 March].  
4 <https://www.kingsfund.org.uk/publications/long-term-conditions-and-mental-health>.  
5  
6  
7 11. Kasteridis P, Street A, Dolman M, Gallier L, Hudson K, Martin J, Wyer I. Who would most  
8 benefit from improved integrated care? Implementing an analytical strategy in South  
9 Somerset. *Int J Integr Care*. 2015; 15:1.  
10  
11 12. Scott KM, Lim C, Al-Hamzawi A, et al. Association of Mental Disorders with Subsequent  
12 Chronic Physical Conditions: World Mental Health Surveys From 17 Countries. *JAMA*  
13 *Psychiatry*. 2016;73(2):150-158. doi:10.1001/jamapsychiatry.2015.2688  
14  
15 13. Ryu E, Chamberlain AM, Pendegraft RS, Petterson TM, Bobo WV, Pathak J. Quantifying  
16 the impact of chronic conditions on a diagnosis of major depressive disorder in adults: a  
17 cohort study using linked electronic medical records. *BMC Psychiatry*. 2016;16:114.  
18 Published 2016 Apr 26. doi:10.1186/s12888-016-0821-x  
19  
20 14. Shih A, Davis K, Schoenbaum S, Gauthier A, Nuzuma R, McCarthy D. Organizing the  
21 U.S. Health Care Delivery System for High Performance. The Commission on a High-  
22 Performance Health System. 2018. Commonwealth Fund.  
23  
24 15. Cawthorpe D. A novel population-based health index for mental disorder. *Perm J*. 2013;  
25 17(2):50–54.  
26  
27 16. Egede LE. Diabetes, major depression and functional disability among U.S. adults.  
28 *Diabetes Care*. 2004;27(2):421-428.  
29  
30 17. Egede LE. Medical cost of depression and diabetes. In: Katon W, Maj M, Sartorius N,  
31 eds. *Depression and Diabetes*. Hoboken, NJ: Wiley Blackwell; 2010:63-79.  
32  
33 18. Gourevitch M.N., Chatterji P., Deb N., Schoenbaum M.D., Turner B.J. On-site medical  
34 care in methadone maintenance: associations with health care use and expenditures. *J*  
35 *Subst. Abus. Treat*. 2007;32:143–151.  
36  
37 19. Paez K.A., Zhao L., Hwang W. Rising out-of-pocket spending for chronic conditions: a  
38 ten-year trend. *Health Aff*. 2009;28:15–25.  
39  
40 20. Hemming P & Loeb D. Internal medicine residents' inadequate preparation in mental  
41 health. *SGIM Forum*. 2013; 36(12), 10-11.  
42  
43 21. Smith, R. C. Educating trainees about common mental health problems in primary care:  
44 A (not so) modest proposal. *Academic Medicine*. 2011; 86(11), e16.  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



22. Smith RC, Laird-Fick H, D'Mello D, Dwamena FC, Romain A, Olson J, Kent K, Blackman K, Solomon D, Spoolstra M, Fortin AH, Frey J, Ferenchick G, Freilich L, Meerschaert C, & Frankel R. Addressing mental health issues in primary care: An initial curriculum for medical residents. *Patient Education and Counseling*. 2014. 94(1), 33–42. <https://doi.org/10.1016/j.pec.2013.09.010>
23. Halvorson GC. 2007. *Health Care Reform Now! A Prescription for Change*. San Francisco, CA: Jossey-Bass.
24. Lenski RE, Barrick JE, Ofria C. Balancing robustness and evolvability. *PLoS Biol*. 2006; 4(12): e 428.
25. Borrell-Carrio F, Suchman AL, Epstein RM. The biopsychosocial model 25 years later: principles, practice, and scientific inquiry. *Ann Fam Med*. 2004; 2(6):576–582.
26. Fortin M, Soubhi H, Hudon C, Bayliss EA, van den Akker M. Multi-morbidity's many challenges. Time to focus on the needs of this vulnerable and growing population. *BMJ*. 2007; 334(7602):1016–1017.
27. Frandsen BR, Joynt KE, Rebitzer JB, Jha AK. Care fragmentation, quality, and costs among chronically ill patients. *Am J Manag Care*. 2015;21(5):355-62. PMID: 26167702.
28. Katon WJ, Lin EHB, Von Korff M, Ciechanowski P, Ludman EJ, Young B, Peterson D, Rutter CM, McGregor M, McCulloch D. Collaborative care for patients with depression and chronic illnesses. *N Engl J Med*. 2010; 363:2611–20.
29. Lloyd C, Nouwen A, Sartorius, N. Prevalence and correlates of depressive disorders in people with Type 2 diabetes: results from the International Prevalence and Treatment of Diabetes and Depression (INTERPRET-DD) study, a collaborative study carried out in 14 countries. *Diabetic Med*. 2018; 35(6):760-769.
30. Sheehy C, Murphy E, Barry M. Depression in rheumatoid arthritis – underscoring the problem. *Rheumatology (Oxford)*. 2006; 45(11):1325–7.
31. Welch CA, Czerwinski D, Ghimire B, Bertsimas D. Depression and costs of health care. *Psychosomatics*. 2009; 50(4):392–401.
32. Vamos EP, Mucsi I, Keszei A, Kopp MS, Novak M. Comorbid depression is associated with increased healthcare utilization and lost productivity in persons with diabetes: a large nationally representative Hungarian population survey. *Psychosomatic Medicine*. 2009; 71(5):501–7.

- 1  
2  
3 33. Gunn JM, Ayton DR, Densley K, Pallant JF, Chondros P, Herrman HE, Dowrick CF. The  
4 association between chronic illness, multimorbidity and depressive symptoms in an  
5 Australian primary care cohort. *Soc Psychiatry Psychiatr Epidemiol*. 2010; 47(3):175–84.  
6  
7  
8 34. Livermore N, Sharpe L, McKenzie D. Panic attacks and panic disorder in chronic  
9 obstructive pulmonary disease: A cognitive behavioral perspective. *Respir Med*. 2010;  
10 104(9):1246–53.  
11  
12  
13 35. Cimpean D, Drake RE. Treating comorbid medical conditions and anxiety/ depression.  
14 *Epidemiol Psychiatr Sci*. 2011; 20(2):141–50.  
15  
16  
17 36. Kaplan B, Crawford S, Cantell M, Kooistra L, Dewey D. 2006. Comorbidity, co-occurrence,  
18 continuum: what's in a name? *Child Care Health Dev* 32(6):723–731.  
19  
20  
21 37. Theis KA, Helmick CG, Hootman JM. Arthritis burden and impact are greater among U.S.  
22 women than men: Intervention opportunities. *J Womens Health (Larchmt)*. 2007;  
23 16(4):441–53.  
24  
25  
26 38. Sale J, Gignac M, Hawker G. The relationship between disease symptoms, life events,  
27 coping and treatment, and depression among older adults with osteoarthritis. *J Rheumatol*.  
28 2008; 35(2):335–42.  
29  
30  
31 39. Cowley D, Dunaway K. Teaching Psychiatry Residents to Work at the Interface of Mental,  
32 Health and Primary Care. *Acad Psychiatry*. 2014; 38:398–404.  
33  
34 40. Belsky M. *The Deepest Well: Healing the Long-Term Effects of Childhood Adversity*: by  
35 Nadine Burke Harris, MD. *Perm J*. 2019; 23:18-075.  
36  
37  
38 41. Sweeney S, Dayo Obure C, Maier CB, Greener R, Dehne K, et al. Costs and efficiency of  
39 integrating HIV/AIDS services with other health services: a systematic review of evidence  
40 and experience. *Sex Transm Infect*. 2011; 88: 85–99.  
41  
42  
43 42. Miranda JJ, Kinra S, Casas JP, Davey Smith G, Ebrahim S. Non-communicable diseases  
44 in low- and middle-income countries: context, determinants and health policy. *Trop Med*  
45 *Int Health*. 2008; 13: 1225–1234  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Figure 2. Number of medical specialties by country addressing mental health comorbidity: affective and anxiety disorders

338x190mm (54 x 54 DPI)



Figure 1. The main five specialty programs with mental health topics: distribution by countries.

338x190mm (54 x 54 DPI)




Figure 3. Number of medical specialties programs by country with comorbid mental health content: affective and anxiety disorders

338x190mm (54 x 54 DPI)




Figure 4. Number of medical specialties programs by country addressing mental health comorbid: substance and alcohol disorders, and Doctor-patient relationship

338x190mm (54 x 54 DPI)