


### CORRELATED ANALYSES OF D- AND $^{15}\text{N}$ -RICH CARBON GRAINS FROM CR2 CHONDRITE EET 92042.

H. Busemann<sup>1</sup>, C. M. O'D. Alexander<sup>1</sup>, L. R. Nittler<sup>1</sup>, T. J. Zega<sup>2</sup>, R. M. Stroud<sup>2</sup>, S. Bajt<sup>3</sup>, G. D. Cody<sup>4</sup> and H. Yabuta<sup>4</sup>.  
<sup>1</sup>DTM, Carnegie Institution of Washington DC. <sup>2</sup>Naval Research Laboratory Washington DC, <sup>3</sup>Lawrence Livermore National Laboratory, Livermore CA, <sup>4</sup>GL, Carnegie Institution of Washington DC, USA. (busemann@dtm.ciw.edu)

**Introduction:** Insoluble organic matter (IOM) and matrix from primitive carbonaceous chondrites carry isotope enrichments ( $\delta\text{D}\leq 20000\text{‰}$ ,  $\delta^{15}\text{N}\leq 3200\text{‰}$ ) that are comparable to those in interplanetary dust particles [1, this work]. Hence, primitive organics that formed in the protosolar cloud (PSC) – or maybe in the cold outer regions of the protoplanetary disk – survived accretion and planetary processing on the asteroids, the parent bodies of the chondrites. Most D and  $^{15}\text{N}$  anomalies are spatially uncorrelated, indicating that distinct processes produced them. While various reactions in the PSC can account for the D enrichments [2], the  $^{15}\text{N}$  anomalies cannot be explained by existing models [3]. Alternative mechanisms, possibly within the solar system [4], have to be considered. Identifying the isotopically anomalous carriers will help to understand the earliest evolution of organic matter from PSC to the solar system.

**Results:** SIMS analyses of CR2 chondrite EET 92042 IOM [5] revealed two isotopically anomalous, micron-size discrete carbon grains (“A” with  $\delta^{13}\text{C}\sim -113\text{‰}$  and  $\delta^{15}\text{N}\sim 1150\text{‰}$ ; “B” with  $\delta\text{D}\sim 6000\text{‰}$ , Fig.). Grains and intermediate IOM ( $\delta\text{D}\sim 2200\text{‰}$ ) were thinned and extracted by FIB-SEM [6] and examined by transmission electron microscopy (Fig.). EDS and electron diffraction patterns show that all analyzed matter is C-rich and amorphous.  $^{15}\text{N}$ -rich grain A is monolithic C with trace Si; D-rich grain B is porous organic C with traces of Si and S. The intermediate IOM is also porous organic C and contains nm-size Fe-Ni, chromite and Ca-rich grains. C- and N-XANES spectroscopy proves the hydro-carbonaceous, non-graphitic character of the IOM and indicates distinct N bonding states for grains A and B. On-going EELS, NanoSIMS, and synchrotron IR micro-spectroscopic analyses will characterize the isotopic compositions of additional elements, and the chemical structural variation and bonding states of the C-bearing molecules.



Part of this work was performed under the auspices of the U.S. DoE by the Univ. of Calif. Lawrence Livermore National Laboratory under Contract No. W-7405-ENG-48. The Advanced Light Source is supported by the Director, Office of Science, Office of Basic Energy Sciences, Materials Sciences Division, of the U.S. DoE under Contract No. DE-AC03-76F00098 at Lawrence Berkeley National Laboratory.

**References:** [1] Busemann H. et al. 2006. *Science* 90:1151–1154. [2] Sandford S. A. 2001. *Meteorit. & Planet. Sci.* 36:1117–1133. [3] Charnley S. B. & Rodgers S. D. 2002. *Astrophys. J.* 569:L133-L137. 27th LPSC. [4] Marty B. 2006. *Science* 90: 706–707. [5] Busemann H. et al. 2006. Abstract #2005. 37th LPSC. [6] Zega T. J. & Stroud R. M. 2006. Abstract #1441. 37th LPSC.