

Open Research Online

The Open University's repository of research publications and other research outputs

Trace element signatures of trapped KREEP in Olivine-rich clasts within lunar meteorite NWA773

Conference or Workshop Item

How to cite:

Bridges, John; Jeffries, T. E. and Grady, Monica (2002). Trace element signatures of trapped KREEP in Olivine-rich clasts within lunar meteorite NWA773. In: 65th Annual Meeting of the Meteoritical Society, 21-26 Jul 2002, Los Angeles, California, USA.

For guidance on citations see [FAQs](#).

© [\[not recorded\]](#)

Version: [\[not recorded\]](#)

Link(s) to article on publisher's website:

<http://www.lpi.usra.edu/meetings/metsoc2002/pdf/5137.pdf>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

TRACE ELEMENT SIGNATURES OF TRAPPED KREEP IN OLIVINE-RICH CLASTS WITHIN LUNAR METEORITE NWA773. J. C. Bridges, T. E. Jeffries and M. M. Grady, Dept. of Mineralogy, Natural History Museum, Cromwell Road, London SW7 5BD, UK. (j.bridges@nhm.ac.uk).

NWA773 (sample BM2001, M23) is a lunar regolith breccia [1] the major component of which is an unusual olivine-rich cumulate in clasts ≤ 2 cm across. We use laser-ICPMS trace element determinations on minerals in the clasts to describe their crystallisation history and melt compositions. They have a feldspathic-peridotite modal mineralogy: 66 vol% ol (Fo_{70-71}); 25.9% augite ($\text{En}_{50}\text{Wo}_{38}$) and pigeonite ($\text{En}_{61}\text{Wo}_{13}$); 8.2% plag (An_{88-91}) and minor constituents (from point counting of our 2 sections, although the clasts were called olivine gabbro-norite in [1]). The bulk composition is 41.2 wt% SiO_2 , 0.4 wt% TiO_2 , 5.7 wt% Al_2O_3 , 26.3 wt% MgO , 19.8 wt% FeO , 5.8 wt% CaO , $100\text{Mg}\# = 70.3$ (average of multiple EPMA point analyses). This composition suggests affinities with the Mg-suite of clasts identified within highland material [2] and in particular Apollo 16 sample 67667 [3].

The olivine, pyroxene and plagioclase have low siderophile and Rb ($<0.4 \times \text{CI}$) abundances (FIG. 1). Augite has the highest total REE abundances (La $5 \times \text{CI}$, Lu $18 \times \text{CI}$) and olivine the lowest (La $0.1 \times \text{CI}$, Lu $4.5 \times \text{CI}$). The bulk clasts have LREE $40 \times \text{CI}$, HREE $20 \times \text{CI}$ [1]. Melt trace element abundances were calculated using partition coefficients from [4,5] and are shown as open symbols in FIG. 1. The calculated melt compositions from the pigeonite and plagioclase ($200-400 \times \text{CI}$ LREE, $90-200 \times \text{CI}$ HREE, negative Eu anomalies) are close to KREEP compositions and trace element abundances suggested for the parental melts of lunar Mg-suite plutonic rocks [4]. However augite in the clasts crystallised from more enriched melt. Taking the original melt's composition as KREEP, and assuming Rayleigh fractionation, implies that augite crystallised from the last 6-10% of trapped melt. Olivine, pigeonite and plagioclase were derived from a primitive KREEP basaltic melt in a peridotitic cumulate pile, augite (together with some minor phases) crystallised from the final fractions of trapped melt.

References: [1] Fagan T. J. et al. (2001) *Meteorit. Planet. Sci.*, 36, A55. [2] Snyder G. A. et al. (1995) *JGR*, 100, 9365-9388. [3] Warren P. H. and Wasson J. T. (1979) *Proc. LPSC* 10, 583-610. [4] Papike J. J. et al. (1996) *GCA*, 60, 3967-3978. [5] <http://EarthRef.org/>




Fig. 1. Trace elements in minerals from NWA773 cumulate clasts.