



## Open Research Online

### Citation

Hathway, Kerry (2023). Comparing meso and macroinvertebrate abundance and taxa richness in soil between mechanically mown and unmown grass in a suburban garden. Student research project for the Open University module SXB390 Biology Project Module

### URL

<https://oro.open.ac.uk/94832/>

### License

(CC-BY-NC-ND 4.0) Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

<https://creativecommons.org/licenses/by-nc-nd/4.0/>

### Policy

This document has been downloaded from Open Research Online, The Open University's repository of research publications. This version is being made available in accordance with Open Research Online policies available from [Open Research Online \(ORO\) Policies](#)

### Versions

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding

**Supplementary Table 1:** No of individuals of each taxa appearing in samples from each plot at different time points in the no-mow treatment. The table also shows total number of individuals and total number of taxa in each sample as well as average per square metre values for each time point.

**Key:** Lumbricidae (**Lumb**): earthworms; Isopoda (**Iso**): woodlice; Formicidae (**Form**): ants; Araneae (**Ara**): spiders; Chilopoda (**Chil**): centipedes; Enchytraeidae (**Ench**): potworms; Acari (**Aca**): mites, ticks; Gastropoda (**Gast**): snails, slugs; Diplura (**Dip**): bristletails; Staphylinidae (**Stap**): rove beetles; Diplopoda (**Dipo**): millipedes; Curculionidea (**Curc**): weevils; Lepidoptera (**Lepi**): moth/butterfly caterpillars; Heteroptera (**Hete**): true bugs

**No-mow**

Date	Plot	Lumb	Iso	Form	Ara	Chil	Ench	Aca	Gast	Dip	Stap	Dipo	Curc	Lepi	Hete	Total no. ind. in sample	Total no. taxa in sample
April 8-14	1	7	0	0	1	0	0	0	2	0	0	0	0	0	0	10	3
	2	2	0	8	0	1	1	0	0	0	0	0	0	0	0	12	4
	3	2	0	0	0	0	0	0	0	0	0	0	0	0	0	2	1
	4	2	1	0	0	0	4	0	0	0	0	0	0	0	0	7	3
	5	4	0	0	0	0	0	0	0	0	0	0	0	0	0	4	1
	6	2	1	2	0	0	0	0	0	0	1	0	0	0	0	6	4
	7	0	1	2	0	0	1	0	0	0	0	0	0	0	0	4	3
	8	2	0	5	0	0	0	0	0	0	0	0	0	0	0	7	2
	9	4	0	0	0	0	1	0	0	0	0	0	0	0	0	5	2
	10	5	2	0	0	0	2	0	0	0	0	0	0	0	0	9	3
	<b>Average/sample</b>	3	0.5	1.7	0.1	0.4	0.6	0	0.2	0	0.1	0	0	0	0	6.6	2.7
	<b>m<sup>-2</sup> average</b>	300	50	170	10	40	60	0	20	0	10	0	0	0	0	660	2.7
April 22-28	1	6	1	0	1	0	0	0	1	0	0	1	0	0	0	10	5
	2	1	4	4	0	0	0	0	0	0	0	0	0	0	0	9	3
	3	0	0	2	0	1	0	0	0	0	0	1	0	0	0	4	3

	4	1	0	2	0	0	2	0	0	0	0	0	0	0	0	0	5	3
	5	4	2	4	0	0	0	0	0	0	0	0	0	0	0	0	10	3
	6	1	1	1	0	0	3	0	2	0	0	0	0	0	0	0	8	5
	7	2	2	2	0	0	0	0	1	0	1	0	0	0	0	0	8	5
	8	2	0	9	0	0	1	0	0	0	0	0	0	0	0	0	12	3
	9	5	0	2	0	0	2	0	0	0	0	0	0	0	0	0	9	3
	10	3	4	2	0	0	7	0	0	0	0	0	0	0	1	0	17	5
	<b>Average/ sample</b>	2.5	1.4	2.8	0.1	0.1	1.5	0	0.4	0	0.1	0.2	0	0.1	0		9.2	3.8
	<b>m<sup>2</sup> average</b>	250	140	280	10	10	150	0	40	0	10	20	0	10	0		920	3.8
May 6-12	1	3	0	0	0	0	2	0	1	1	2	1	0	0	0	0	10	6
	2	3	0	1	0	0	2	0	0	0	0	2	0	0	0	0	8	4
	3	0	0	4	0	3	0	0	0	0	2	0	0	0	0	0	9	3
	4	0	0	0	1	1	3	0	0	0	0	0	0	0	0	0	5	3
	5	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	2	2
	6	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	1
	7	2	0	5	0	0	0	0	0	0	0	0	0	0	0	0	7	2
	8	2	0	4	3	1	0	0	0	0	3	0	0	0	0	0	13	5
	9	3	0	13	0	0	1	0	1	0	2	0	0	0	0	0	20	5
	10	3	0	7	0	0	1	0	0	0	2	0	0	0	0	0	13	4
	<b>Average/ sample</b>	1.9	0	3.5	0.4	0.5	0.9	0	0.2	0.1	1.1	0.3	0	0	0		8.9	3.5
	<b>m<sup>2</sup> average</b>	190	0	350	40	50	90	0	20	10	110	30	0	0	0		890	3.5
May 20-26	1	1	1	7	0	0	0	0	0	0	0	0	0	0	0	0	9	3
	2	2	3	40	0	0	0	0	0	0	0	0	0	0	0	0	45	3
	3	0	1	15	0	1	1	0	0	0	0	0	0	0	0	0	18	4
	4	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	2	2

	5	2	0	4	1	0	0	0	0	0	0	1	0	0	0	8	4
	6	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	1
	7	4	0	0	1	0	1	0	0	0	0	0	0	0	1	7	4
	8	0	0	4	0	0	1	0	0	0	0	0	0	0	0	5	2
	9	2	1	13	0	1	0	0	0	0	0	0	0	0	0	17	4
	10	0	2	37	0	0	1	0	0	0	0	0	0	0	0	40	3
	<b>Average/ sample</b>	1.1	0.9	12	0.2	0.2	0.6	0	0	0	0	0.1	0	0	0.1	15.2	3
	<b>m<sup>-2</sup> average</b>	110	90	1200	20	20	60	0	0	0	0	10	0	0	10	1520	3
Jun 3-9	1	1	3	15	1	0	0	0	0	0	0	0	0	0	0	20	4
	2	0	1	29	1	0	2	0	0	0	0	0	0	0	0	33	4
	3	0	1	11	2	0	0	0	0	1	0	1	0	0	0	16	5
	4	0	0	4	0	0	0	0	0	0	0	0	0	0	0	4	1
	5	1	0	50	1	0	0	0	0	0	0	0	0	0	0	52	3
	6	0	1	5	1	1	0	0	0	0	0	0	0	0	0	8	4
	7	0	1	23	0	0	0	0	0	0	0	0	0	0	0	24	2
	8	0	1	7	0	0	0	0	0	0	0	0	0	0	0	8	2
	9	4	0	8	2	0	0	0	0	0	0	0	1	0	0	15	4
	10	1	1	4	0	0	0	0	0	0	0	0	0	0	0	6	3
	<b>Average/ sample</b>	0.7	0.9	15.6	0.8	0.1	0.2	0	0	0.1	0	0.2	0	0	0	18.6	3.2
	<b>m<sup>-2</sup> average</b>	70	90	1560	80	10	20	0	0	10	0	20	0	0	0	1860	3.2
	<b>Avg across study/m<sup>-2</sup></b>	<b>184</b>	<b>74</b>	<b>712</b>	<b>32</b>	<b>26</b>	<b>76</b>	<b>0</b>	<b>16</b>	<b>4</b>	<b>26</b>	<b>16</b>	<b>0</b>	<b>2</b>	<b>2</b>	<b>1170</b>	<b>3.22</b>

**Supplementary Table 2:** No of individuals of each taxa appearing in samples from each plot at different time points in the mown treatment. The table also shows total number of individuals and total number of taxa in each sample as well as average per square metre values for each time point.

**Key:** Lumbricidae (**Lumb**): earthworms; Isopoda (**Iso**): woodlice; Formicidae (**Form**): ants; Araneae (**Ara**): spiders; Chilopoda (**Chil**): centipedes; Enchytraeidae (**Ench**): potworms; Acari (**Aca**): mites, ticks; Gastropoda (**Gast**): snails, slugs; Diplura (**Dip**): bristletails; Staphylinidae (**Stap**): rove beetles; Diplopoda (**Dipo**): millipedes; Curculionidea (**Curc**): weevils; Lepidoptera (**Lepi**): moth/butterfly caterpillars; Heteroptera (**Hete**): true bugs

**Mown**

Date	Plot	Lumb	Iso	Form	Ara	Chil	Ench	Aca	Gast	Dip	Stap	Dipo	Curc	Lepi	Hete	Total no. ind. in sample	Total no. taxa in sample
April 8-14	1	4	0	0	0	0	1	0	0	0	0	0	0	0	0	5	2
	2	1	0	0	0	0	0	0	0	0	1	0	0	0	0	2	2
	3	0	0	14	0	0	0	0	0	0	0	0	0	1	0	15	2
	4	2	0	1	0	1	2	0	0	0	0	0	0	0	0	6	4
	5	2	3	11	0	1	0	0	0	0	0	0	0	0	0	17	4
	6	6	0	0	0	0	0	0	0	0	0	0	0	0	0	6	1
	7	4	0	5	0	0	0	0	0	0	0	0	0	0	0	9	2
	8	2	2	15	1	1	1	0	0	0	0	0	0	0	0	22	6
	9	4	0	0	0	0	0	0	0	0	0	0	0	0	0	4	1
	10	2	1	8	0	1	0	0	0	0	0	0	0	0	0	12	4
	<b>Average/sample</b>	2.7	0.6	5.4	0.1	0.4	0.4	0	0	0	0.1	0	0	0.1	0	9.8	2.8
	<b>m<sup>-2</sup> average</b>	270	60	540	10	40	40	0	0	0	10	0	0	10	0	980	2.8
April 22-28	1	2	0	0	0	0	0	0	0	0	0	0	0	0	0	2	1
	2	2	0	34	0	0	0	0	0	0	0	0	0	0	0	36	2

	3	0	0	11	0	0	0	0	0	0	0	0	0	0	0	0	11	1
	4	2	1	0	0	0	3	0	2	0	0	0	0	0	0	0	8	4
	5	6	7	4	0	2	2	1	0	0	0	0	1	0	0	0	23	7
	6	4	9	3	0	0	2	0	0	0	0	0	0	0	0	0	18	4
	7	4	1	5	1	0	1	0	0	0	0	0	0	1	0	0	13	6
	8	1	0	14	0	0	1	0	0	0	1	0	0	0	0	0	17	4
	9	3	0	0	0	0	1	0	0	0	0	1	0	0	0	0	5	3
	10	1	0	6	0	0	4	0	0	0	0	0	0	0	0	0	11	3
	<b>Average/ sample</b>	2.5	1.8	7.7	0.1	0.2	1.4	0.1	0.2	0	0.1	0.1	0.1	0.1	0	0	14.4	3.5
	<b>m<sup>2</sup> average</b>	250	180	770	10	20	140	10	20	0	10	10	10	10	0	0	1440	3.5
May 6-12	1	0	0	3	0	0	0	0	0	0	0	0	0	0	0	0	3	1
	2	3	0	0	1	1	0	0	0	0	0	0	0	0	0	0	5	3
	3	1	1	1	0	2	1	0	0	0	0	0	0	0	0	0	6	5
	4	0	0	1	0	1	2	0	0	0	0	0	0	0	0	0	4	3
	5	0	0	2	0	0	2	0	0	0	0	0	0	0	0	0	4	2
	6	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	2	2
	7	2	8	7	0	0	6	0	0	0	0	1	0	0	0	0	24	5
	8	1	0	2	0	3	3	0	0	0	0	0	0	0	0	0	9	4
	9	2	0	1	0	0	0	0	0	0	0	0	0	0	0	0	3	2
	10	4	1	0	0	1	6	0	0	0	0	0	0	0	0	0	12	4
	<b>Average/ sample</b>	1.4	1	1.7	0.1	0.9	2	0	0	0	0	0.1	0	0	0	0	7.2	3.1
	<b>m<sup>2</sup> average</b>	140	100	170	10	90	200	0	0	0	0	10	0	0	0	0	720	3.1
May 20-26	1	2	0	5	0	0	0	0	0	0	0	0	0	0	0	0	7	2
	2	3	0	7	0	0	0	0	0	0	0	0	0	0	0	0	10	2
	3	0	0	9	0	0	0	0	0	0	0	1	0	0	0	0	10	2

	4	2	0	3	0	0	2	0	0	0	0	0	0	0	0	0	7	3
	5	4	0	6	0	0	4	0	0	0	0	0	0	0	0	0	14	3
	6	0	0	9	0	0	1	0	0	0	0	0	0	0	0	0	10	2
	7	1	2	7	2	1	2	1	0	0	0	1	0	0	0	0	17	8
	8	0	0	4	0	1	2	0	0	0	0	0	0	0	0	0	7	3
	9	3	0	5	0	1	0	0	0	0	0	0	0	0	0	0	9	3
	10	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	2	2
	<b>Average/ sample</b>	1.5	0.2	5.6	0.2	0.3	1.2	0.1	0	0	0	0.2	0	0	0	0	9.3	3
	<b>m<sup>-2</sup> average</b>	150	20	560	20	30	120	10	0	0	0	20	0	0	0	0	930	3
Jun 3-9	1	1	0	1	0	0	0	0	0	0	0	2	0	0	0	0	4	3
	2	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	1
	3	0	0	13	0	0	0	1	0	0	0	0	0	0	0	0	14	2
	4	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	2	2
	5	0	0	7	0	1	0	0	0	0	0	1	0	0	0	0	9	3
	6	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	2	2
	7	2	3	6	1	0	2	0	0	0	0	0	0	0	0	0	14	5
	8	0	0	5	0	0	0	0	0	0	0	0	0	0	0	0	5	1
	9	1	3	2	0	3	0	0	0	0	0	1	2	0	1	0	13	7
	10	0	0	13	0	0	0	0	0	0	0	0	0	0	0	0	13	1
	<b>Average/ sample</b>	0.5	0.7	4.7	0.1	0.4	0.3	0.1	0	0	0.1	0.7	0	0.1	0	0	7.7	2.7
	<b>m<sup>-2</sup> average</b>	50	70	470	10	40	30	10	0	0	10	70	0	10	0	0	770	2.7
	<b>Average across study/ m<sup>-2</sup></b>	<b>172</b>	<b>86</b>	<b>502</b>	<b>12</b>	<b>44</b>	<b>106</b>	<b>6</b>	<b>4</b>	<b>0</b>	<b>6</b>	<b>22</b>	<b>2</b>	<b>6</b>	<b>0</b>	<b>968</b>	<b>3.02</b>	

**Supplementary Table 3:** Mean number of individuals of each taxa (per m<sup>2</sup>) at each dated survey point for no-mow and mown treatments (n = 10). Weather conditions, mows omitted or carried out, mean grass height (cm) also included.

<b>Date</b>	April 8-14	April 22-28	May 6-12	May 20-26	June 3-9
<b>Conditions</b>	Sun/winds	Light cloud/sun	Sun	Sun/winds	Sun/warm
<b>No-mow (n = 10)</b>					
Mows omitted	1	2	3	4	5
Mean grass height/cm	10	15	25	35	45
Average number of individuals of each taxa/per square metre					
Lumbricidae	300	250	190	110	70
Formicidae	170	280	350	1200	1560
Chilopoda	40	10	50	20	10
Araneae	10	10	40	20	80
Enchytraeidae	60	150	90	60	20
Isopoda	50	140	-	90	90
Staphylinidae	10	10	110	-	-
Gastropoda	20	40	20	-	-
Diplopoda	-	20	30	10	20
Lepidoptera	-	10	-	-	-
Diplura	-	-	10	-	10
Heteroptera	-	-	-	10	-
<b>Mown (n = 10)</b>					
No of mows since start	1	2	3	4	5
Days since last mow	8	4	5	4	4
Mean grass height/cm	3	3	3	3	3
Average number of individuals of each taxa/per square metre					
Lumbricidae	270	250	140	150	50
Formicidae	540	770	170	560	470
Chilopoda	40	20	90	30	40
Araneae	10	10	10	20	10
Isopoda	60	180	100	20	70
Staphylinidae	10	10	-	-	10
Lepidoptera	10	10	-	-	10
Enchytraeidae	-	140	200	120	30
Diplopoda	-	10	10	20	70
Acari	-	10	-	10	10
Gastropoda	-	20	-	-	-
Curculionoidea	-	10	-	-	-



**Supplementary Tables 4:** Results of two way ANOVA with replication for invertebrate abundance. Count, sum, mean and variance for each data collection period and total across study are shown. Table at bottom is data summary showing test statistics (F), p-values and critical values. Sample refers to comparison between time points (April 8-14, April 22-28, May 6-12, May 20-26, June 3-9), columns refers to comparison between treatments (mown, no-mow).

<i>April 8-14</i>	<b>Mown</b>	<b>No-mow</b>	<b>Total</b>	<i>April 22-28</i>	<b>Mown</b>	<b>No-mow</b>	<b>Total</b>
Count	10	10	20	Count	10	10	20
Sum	9800	6600	16400	Sum	14400	9200	23600
Average	980	660	820	Average	1440	920	1180
Variance	421777.8	93777.78	271157.9	Variance	964888.9	130666.7	590105.3

  

<i>May 6-12</i>	<b>Mown</b>	<b>No-mow</b>	<b>Total</b>	<i>May 20-26</i>	<b>Mown</b>	<b>No-mow</b>	<b>Total</b>
Count	10	10	20	Count	10	10	20
Sum	7200	8900	16100	Sum	9300	15200	24500
Average	720	890	805	Average	930	1520	1225
Variance	441777.8	303222.2	360500	Variance	169000	2390667	1304079

  

<i>June 3-9</i>	<b>Mown</b>	<b>No-mow</b>	<b>Total</b>	<i>Total</i>	<b>Mown</b>	<b>No-mow</b>
Count	10	10	20	Count	50	50
Sum	7700	18600	26300	Sum	48400	58500
Average	770	1860	1315	Average	968	1170
Variance	297888.9	2189333	1490816	Variance	487934.7	1142143

ANOVA

<i>Source of Variation</i>	<i>SS</i>	<i>df</i>	<i>MS</i>	<i>F</i>	<i>P-value</i>	<i>F crit</i>
Sample	4577400	4	1144350	1.545792	0.19575862	2.472927
Columns	1020100	1	1020100	1.377955	0.243546803	3.946876
Interaction	8669400	4	2167350	2.927664	0.025147357	2.472927
Within	66627000	90	740300			
Total	80893900	99				

**Supplementary Tables 5:** Results of two way ANOVA with replication for taxa richness. Count, sum, mean and variance for each data collection period and total across study are shown. Table at bottom is data summary showing test statistics (F), p-values and critical values. Sample refers to comparison between time points (April 8-14, April 22-28, May 6-12, May 20-26, June 3-9), columns refers to comparison between treatments (mown, no-mow).

<i>April 8-14</i>	<b>Mown</b>	<b>No-mow</b>	<b>Total</b>	<i>April 22-28</i>	<b>Mown</b>	<b>No-mow</b>	<b>Total</b>
Count	10	10	20	Count	10	10	20
Sum	28	26	54	Sum	35	38	73
Average	2.8	2.6	2.7	Average	3.5	3.8	3.65
Variance	2.622222	1.155556	1.8	Variance	3.833333	1.066667	2.344737

  

<i>May 6-12</i>	<b>Mown</b>	<b>No-mow</b>	<b>Total</b>	<i>May 20-26</i>	<b>Mown</b>	<b>No-mow</b>	<b>Total</b>
Count	10	10	20	Count	10	10	20
Sum	31	35	66	Sum	30	30	60
Average	3.1	3.5	3.3	Average	3	3	3
Variance	1.877778	2.5	2.115789	Variance	3.333333	1.111111	2.105263

  

<i>June 3-9</i>	<b>Mown</b>	<b>No-mow</b>	<b>Total</b>	<i>Total</i>	<b>Mown</b>	<b>No-mow</b>
Count	10	10	20	Count	50	50
Sum	27	32	59	Sum	151	161
Average	2.7	3.2	2.95	Average	3.02	3.22
Variance	3.788889	1.511111	2.576316	Variance	2.917959	1.522041

ANOVA

<i>Source of Variation</i>	<i>SS</i>	<i>df</i>	<i>MS</i>	<i>F</i>	<i>P-value</i>	<i>F crit</i>
Sample	10.66	4	2.665	1.16886	0.329910162	2.472927
Columns	1	1	1	0.438596	0.509492974	3.946876
Interaction	1.7	4	0.425	0.186404	0.944927123	2.472927
Within	205.2	90	2.28			
Total	218.56	99				

**Supplementary Tables 6:** Results of post hoc pairwise t-tests comparing invertebrate abundance between mown and no-mow plots for data collection periods May 20-26 (a) and June 3-9 (b).

(a)

	<b>Mown</b>	<b>No-mow</b>
Mean	930	1520
Variance	169000	2390667
Observations	10	10
Pearson Correlation	-0.34542	
Hypothesized Mean Difference	0	
df	9	
t Stat	-1.07741	
P(T<=t) one-tail	0.154664	
t Critical one-tail	2.262157	
P(T<=t) two-tail	0.309328	
t Critical two-tail	2.685011	

(b)

	<b>Mown</b>	<b>No-mow</b>
Mean	770	1860
Variance	297888.8889	2189333
Observations	10	10
Pearson Correlation	0.06301441	
Hypothesized Mean Difference	0	
df	9	
t Stat	-2.231730993	
P(T<=t) one-tail	0.026274033	
t Critical one-tail	2.262157163	
P(T<=t) two-tail	0.052548065	
t Critical two-tail	2.685010847	

**Supplementary Table 7:** Soil pH, temperature, qualitative moisture measurements and light levels in each plot during each data collection period for a) no-mow and b) mown treatments

a)

<i>No-mow</i>						
<i>Date</i>	<i>Plot</i>	<i>pH</i>	<i>Temp/°C</i>	<i>Moisture (v. dry, dry, norm, wet, v. wet)</i>	<i>Texture</i>	<i>Light (low, norm, high)</i>
<i>April 8-14</i>	1	6.5	8	v. wet	Clumpy, heavy clay	low
	2	6.5	11	v. wet		high
	3	6.5	10	v. wet		norm
	4	6	11	v. wet		norm
	5	6	11	v. wet		low
	6	6.5	12	v. wet		low
	7	5.5	12	v. wet		low
	8	6.5	12	v. wet		low
	9	6	12	v. wet		low
	10	6	12	v. wet		low
	Mean values (soil pH, temp)		6.2	11.1		
<i>April 22-28</i>	1	6	15	v. wet		low
	2	6	14	v. wet		low
	3	6.5	13	v. wet		low
	4	6	14	v. wet		low
	5	6.5	13	v. wet		low
	6	6.5	11	v. wet		low
	7	5.5	11	v. wet		low
	8	6.5	11	v. wet		low
	9	6	11	v. wet		low
	10	6.5	11	v. wet		low
	Mean values (soil pH, temp)		6.2	12.4		
<i>May 6-12</i>	1	6.5	16	v. wet		low
	2	5	17	v. wet		low
	3	6	16	v. wet		low
	4	5	16	wet	Crumblier	low
	5	6.5	15	v. wet	Heavy, rolls into sausage	low
	6	6.5	15	v. wet	Heavy texture	norm
	7	5.5	15	v. wet		low
	8	6.5	16	v. wet		norm

	9	6	15	v. wet		low
	10	7	15	v. wet		low
	Mean values (soil pH, temp)	6.05	15.6			
<i>May 20-26</i>	1	6.5	19	dry	Crumbly, drier	low
	2	7	20	norm		norm
	3	6.5	18	norm		norm
	4	6.5	17	norm		norm
	5	6.5	16	norm		low
	6	6.5	17	norm	Crumbly, drier	norm
	7	5.5	19	norm		low
	8	6	18	norm		high
	9	6.5	18	norm		high
	10	6	17	norm		norm
	Mean values (soil pH, temp)	6.35	17.9			
<i>June 3-9</i>	1	6.5	17	dry	Hard, dry	low
	2	7	17	dry		low
	3	6.5	18	dry	Crumbly	low
	4	7	17	dry	Crumbly	low
	5	7	16	dry		low (shaded)
	6	6.5	21	dry		norm
	7	6.5	21	dry		low
	8	6.5	20	dry		low
	9	7	21	dry	Crumbly, not as dry as mown plots	low
	10	6.5	21	dry	Crumbly, not as dry as mown plots	low
	Mean values (soil pH, temp)	6.7	18.9			

b)

<i>Mown</i>						
<i>Date</i>	<i>Plot</i>	<i>pH</i>	<i>Temp/°C</i>	<i>Moisture (v. dry, dry, wet, v. wet)</i>	<i>Texture</i>	<i>Light (low, norm, high)</i>
<i>April 8-14</i>	1	6.6	10	v. wet	Clumpy, heavy clay	low
	2	6.5	9	v. wet	Clumpy, heavy clay	low
	3	6	12	v. wet	Clumpy, heavy clay	low
	4	6	11	v. wet		low
	5	6	12	v. wet		high
	6	6.5	14	v. wet		norm
	7	6.5	14	v. wet		low
	8	6.5	12	v. wet		high
	9	6.5	13	v. wet		low
	10	6	13	v. wet		high
	Mean values (soil pH, temp)	6.31	12			
<i>April 22-28</i>	1	6	13	wet	Heavy clay, drier than last session	low
	2	6.5	12	wet		low
	3	6.5	13	wet		low
	4	6.5	12	wet		low
	5	6	12	wet		low
	6	6	14	v. wet		low
	7	6.5	16	v. wet		low
	8	6.5	15	v. wet		low
	9	6.5	16	v. wet		low
	10	6	16	v. wet		low
	Mean values (soil pH, temp)	6.3	13.9			
<i>May 6-12</i>	1	6.5	18	v. wet	Heavy clay	norm
	2	6.5	18	v. wet		low
	3	7	14	v. wet		norm
	4	6.5	14	v. wet	Sticky, heavy clay	norm
	5	6.5	15	v. wet		norm
	6	7	15	v. wet	Heavy, binds into sausage	high
	7	6.5	15	wet		norm
	8	6.5	15	wet		norm
	9	6.5	15	wet		norm

	10	6.5	45	wet		norm
	Mean values (soil pH, temp)	6.6	18.4			
<i>May 20-26</i>	1	6.5	22	dry	Crumbly	norm
	2	6.5	20	dry	Crumbly	norm
	3	6.5	20	dry	Crumbly	norm
	4	6.5	21	norm		norm
	5	6.5	21	norm		low (shaded)
	6	6.5	22	norm	Crumbly	high
	7	6	21	dry	Crumbly, dry	low (shaded)
	8	5	21	dry		high
	9	6.5	21	dry		low (shaded)
	10	6	21	norm		high
	Mean values (soil pH, temp)	6.25	21			
<i>June 3-9</i>	1	6.5	17	dry	Crumbly	low
	2	7	18	dry	Crumbly	low
	3	7	19	dry	Softer than plot 5	low
	4	6.5	18	dry	Softer than plot 5	low
	5	7	18	v. dry	Hard, crumbles with force	low
	6	6.5	22	v. dry	Crumbly	norm
	7	6.5	19	v. dry		norm (shaded)
	8	6.5	21	dry		norm
	9	6.5	22	dry	Crumbly, less dry than plot 10	low
	10	7	22	v. dry	V. crumbly	high
	Mean values (soil pH, temp)	6.7	19.6			